

ANGEL ACTION · DENTAL

Knocked-Out Tooth

Hold a knocked-out adult tooth by the crown, rinse gently with milk or saline if dirty, and reinsert into the socket if possible — otherwise store in milk or saliva and get to a dentist within 30 minutes.

Hold a knocked-out adult tooth by the crown, rinse gently with milk or saline if dirty, and reinsert into the socket if possible — otherwise store in milk or saliva and get to a dentist within 30 minutes.

Knocked-out adult tooth

1. **Hold the tooth by the crown (white part)** — never touch the root.
2. If dirty, rinse very gently with milk or saline (not water if you can avoid it). Do NOT scrub.
3. Try to **reinsert it into the socket** the right way around and have the person bite gently on a clean cloth.
4. If you cannot reinsert it, store the tooth in **milk** or in the person's **own saliva** (between cheek and gum, if old enough not to swallow it).
5. See a **dentist within 30 minutes** — the sooner the better.

Baby teeth

Do **not** reinsert a knocked-out baby tooth — it can damage the developing adult tooth. Still see a dentist.

Broken or chipped tooth

- Save any tooth fragments in milk or saliva.
- Rinse the mouth gently with warm water.
- Apply a cold pack to the cheek to reduce swelling.
- See a dentist as soon as possible.

Adapted from Australian First Aid 5th Edition. Get the full guide and live CPR metronome at firstaidangel.org.