

Elliptic Curves Over Finite Fields and Applications to Cryptography

2026 School on Arithmetic, Geometry and Algorithms

Juanita Duque-Rosero

Bernoulli Institute, University of Groningen.

j.duque.rosero@rug.nl

Introduction

These lecture notes accompany the lectures for the course on elliptic curves over finite fields and applications to cryptography for the 2026 School on Arithmetic, Geometry and Algorithms. The main references I used are [Kun26, Sil09]. Other useful references are [HPS14, AM26].

At the heart of public-key cryptography are functions that are easy to compute but are believed hard to reverse without secret information. In these notes, we first build the arithmetic of elliptic curves over finite fields from explicit formulas. We then use it to formulate the discrete logarithm problem and elliptic-curve Diffie–Hellman. At the end, we will discuss isogenies and their cryptographic context.

These notes were prepared with the help of ChatGPT 5.6 Terra. All errors are my own.

Explicit computations

The ideal way to follow these notes is to try examples in your favorite computer algebra system. I recommend you use the open source software [SageMath](#), but you are welcome to use whatever you prefer ([Magma](#), [Oscar](#), [PARI/GP](#), etc.).

Prerequisites

The required background includes arithmetic in $\mathbb{F}_p$: modular addition, multiplication, inverses, and polynomial evaluation. Basic group theory is also used in the course.

A note from the author

Despite my best efforts, these notes may contain typos. If you spot any, please feel free to email me; I appreciate your help!

Acknowledgments

These notes were inspired by the Arizona Winter School 2026: Computational Aspects of Arithmetic Geometry and Cryptography, and the Preliminary Arizona Winter School 2025: Mathematical cryptography and algorithms in number theory. Thank you to the speakers, Sarah Arpin, Kirsten Eisentraeger, David Harvey, Sabrina Kunzweiler, Chloe Martindale, and John Voight and to the organizers Renee Bell, Daniel Erman, Brandon Levin, Anna Medvedovsky, Padma Srinivasan, Anthony Várilly-Alvarado, Isabel Vogt, and Hang Xue.

Contents

1	Elliptic curves over finite fields	1
1.1	A short toolkit	1
1.1.1	Arithmetic in $\mathbb{F}_p$	1
1.2	Curves and their points	2
1.2.1	Points on elliptic curves	3
1.2.2	Counting points by hand	4
1.3	The group law	5
1.4	How many points can there be?	7
1.5	Practice problems	7
2	The elliptic-curve discrete logarithm problem	9
2.1	Public-key cryptography	9
2.2	Further structure of the points on elliptic curves	10
2.3	The discrete logarithm problem	11
2.4	Diffie–Hellman key agreement	12
2.5	What is assumed to be hard?	14
3	Isogenies of elliptic curves	15
3.1	Maps between elliptic curves	15
3.1.1	Isomorphisms and the j -invariant	16
3.2	Isogenies	16
3.2.1	Subgroups and isogenies	17
3.3	Isogenies in cryptography: SIDH	19
4	ℓ-Isogenies and ordinary/supersingular curves	20
4.1	ℓ -isogenies	20
4.2	Ordinary and supersingular elliptic curves	21
4.3	Isogenies and supersingular elliptic curves	22
5	Isogeny graphs and cryptography	24
5.1	Isogeny graphs	24
5.1.1	Properties of isogeny graphs	27
5.2	Cryptography and isogeny graphs	28
5.2.1	CSIDH	28
5.2.2	SQIsign	30
5.3	Extra: how does one describe isogenies?	30

Lecture 1: Elliptic curves over finite fields

This course starts with arithmetic in prime fields. Our immediate goal is to construct a finite abelian group from a polynomial equation and learn to compute in it. The cryptographic applications in later lectures use exactly this arithmetic, but with primes and subgroup orders too large to enumerate.

Throughout this lecture, $p > 3$ is a prime.

1.1 A short toolkit

1.1.1 Arithmetic in $\mathbb{F}_p$

The elements of $\mathbb{F}_p$ are $0, 1, \dots, p-1$, with addition and multiplication reduced modulo p . Every nonzero element $a \in \mathbb{F}_p$ has a unique inverse a^{-1} , so division by a means multiplication by a^{-1} . For example, in $\mathbb{F}_7$ we have $2^{-1} = 4$, since $2 \cdot 4 = 8 = 1$.

Definition 1.1. An element $r \in \mathbb{F}_p$ is a **quadratic residue** if $r = y^2$ for some $y \in \mathbb{F}_p$. Otherwise it is a **quadratic nonresidue**.

For p odd, every nonzero quadratic residue has exactly two square roots, y and $-y$, while 0 has one. Thus a table of squares is enough to count points on the curves in this lecture.

Example 1.2. The squares in $\mathbb{F}_7$ are

$$0^2 = 0, \quad 1^2 = 6^2 = 1, \quad 2^2 = 5^2 = 4, \quad 3^2 = 4^2 = 2.$$

Hence the quadratic residues in $\mathbb{F}_7$ are 0, 1, 2, 4.

 **Exercise 1.3.** In this problem, you will study quadratic residues in finite fields.

1. Find the quadratic residues in $\mathbb{F}_{11}$ and $\mathbb{F}_{13}$. How many are there in each case?
2. Can you find a pattern in the number of quadratic residues in $\mathbb{F}_p$ for odd primes p ?
3. Can you prove the formula you found? *Hint: consider the map $x \mapsto x^2$ on the nonzero elements of $\mathbb{F}_p$.*

1.2 Curves and their points

An elliptic curve over a field k is a geometric object described here by an equation in two variables. In this course we use the **short Weierstrass equation**

$$y^2 = x^3 + ax + b,$$

where a and b are chosen constants in k . Usually, k will denote the finite fields $\mathbb{F}_p$ or $\mathbb{F}_{p^2}$. A solution $(x, y) \in k^2$ is called an **affine point** of the curve, or simply a point of the curve.

For example, on $y^2 = x^3 + x + 1$ over $\mathbb{F}_7$, taking $x = 2$ gives

$$y^2 = 2^3 + 2 + 1 = 4.$$

In 1.2, we saw that 4 is a quadratic residue in $\mathbb{F}_7$, so there are two values of y that satisfy the equation. Since $2^2 = 4$ and $5^2 = 4$ in $\mathbb{F}_7$, this value $x = 2$ gives the two affine points $(2, 2)$ and $(2, 5)$. For a fixed x , there can similarly be zero, one, or two affine points, according as the right-hand side is a nonresidue, zero, or a nonzero quadratic residue.

Finally, not every equation of this shape gives an elliptic curve. We require that it have no singular point—no crossing or cusp in its geometric picture. For a short Weierstrass equation in characteristic different from 2 and 3, this is exactly the nonvanishing condition in the following definition. It is what makes the addition law well behaved.

Definition 1.4. Recall that $p > 3$. An elliptic curve in short Weierstrass form over $\mathbb{F}_p$ is an equation

$$E: y^2 = x^3 + ax + b, \quad a, b \in \mathbb{F}_p, \quad (1.5)$$

such that

$$4a^3 + 27b^2 \neq 0.$$

Its set of $\mathbb{F}_p$ -points is

$$E(\mathbb{F}_p) = \{(x, y) \in \mathbb{F}_p^2 : y^2 = x^3 + ax + b\} \cup \{O\}.$$

The extra symbol O , called the **point at infinity**, will be the identity element of the group law.

To see where O comes from, homogenize the equation:

$$Y^2Z = X^3 + aXZ^2 + bZ^3.$$

In projective coordinates the solutions with $Z = 0$ consist of the single point $[0 : 1 : 0]$, represented in the affine formulas by O . Including this point is what allows point addition to have an identity.

Remark 1.6. The restriction $p > 3$ is deliberate: it makes this short equation and the formulas below valid without exceptional characteristic-2 or characteristic-3 cases.

 **Exercise 1.7.** Graph the real solutions of $y^2 = f(x)$ for

$$f(x) = x^3 - 1, \quad f(x) = x^3 - x, \quad f(x) = x^3 - x^2.$$

Which of these equations are nonsingular and hence define elliptic curves over $\mathbb{R}$?

Definition 1.8. The discriminant of (1.5) is

$$\Delta(E) = -16(4a^3 + 27b^2).$$

The condition $\Delta(E) \neq 0$ says that the cubic on the right has no repeated root. In this case, we say that E is nonsingular. If $\Delta(E) = 0$, then E is singular and has a cusp or crossing point.

 **Exercise 1.9.** For which primes $p > 3$ does $y^2 = x^3 + x + 2$ fail to define an elliptic curve over $\mathbb{F}_p$? What about over $\mathbb{F}_{p^2}$?

 **Exercise 1.10.** Let k be a field of characteristic different from 2 and 3, and let $f(x) = x^3 + ax + b$.

1. Show that $4a^3 + 27b^2 = 0$ if and only if f has a repeated root.
2. Use the partial derivatives of $g(x, y) = y^2 - x^3 - ax - b$ to show that f having a repeated root is equivalent to the existence of a singular affine point on the curve.

1.2.1 Points on elliptic curves

The same equation can be studied over any field containing its coefficients.

Definition 1.11. Let k and K be fields with $k \subseteq K$. Then we say that K is an extension field of k , denoted by K/k .

If E is defined over k and K/k is a field extension, then

$$E(K) := \{(x, y) \in K^2 : y^2 = x^3 + ax + b\} \cup \{O\}.$$

The notation $\#E(K)$ denotes the number of these points when the set is finite.

Example 1.12 (One curve over several fields). Consider the nonsingular short Weierstrass equation

$$E: y^2 = x^3 - 27x + 2970,$$

Over $\mathbb{C}$, every value of x except a root of the cubic gives two values of y , so $E(\mathbb{C})$ is infinite. Over $\mathbb{R}$,

$$x^3 - 27x + 2970 = (x + 15)(x^2 - 15x + 198),$$


and $x^2 - 15x + 198 > 0$ for every real x . Thus $E(\mathbb{R})$ has affine points exactly when $x \geq -15$, including $(-15, 0)$. Over $\mathbb{Q}$, the points $(-15, 0)$, $(3, \pm 54)$, and $(21, \pm 108)$ are rational.

Over $\mathbb{F}_{11}$, direct enumeration gives

$$E(\mathbb{F}_{11}) = \{O, (0, 0), (2, \pm 3), (3, \pm 1), (4, 0), (5, \pm 1), (7, 0), (10, \pm 2)\},$$

so $\#E(\mathbb{F}_{11}) = 12$. Over the quadratic extension $\mathbb{F}_{11^2}$ the same curve has 144 points. The points over $\mathbb{F}_{11}$ are included among these extension-field points. Questions about integral points, obtained by asking for integer coordinates rather than field-valued coordinates, lead to deeper arithmetic geometry; see [Zag87].

A useful source for further examples and arithmetic data is the [LMFDB elliptic-curve database](#). The page for the curve in Example 1.12 is [this](#). It contains a wealth of information, including the number of points over many finite fields, the group structure of $E(\mathbb{F}_p)$ for many primes p , and the torsion subgroup over $\mathbb{Q}$.



Exercise 1.13. For each curve below, compute $\#E(\mathbb{F}_5)$. Then use SageMath to compute $\#E(\mathbb{F}_{5^2})$. Do you see a pattern?

$$y^2 = x^3 - x, \quad y^2 = x^3 + x + 1.$$

1.2.2 Counting points by hand

For each $x \in \mathbb{F}_p$, compute $x^3 + ax + b$ and look it up in the table of quadratic residues. A nonzero residue gives two points, 0 gives one, and a nonresidue gives none. Finally add O .

Example 1.14. Let $E/\mathbb{F}_7$ be

$$E: y^2 = x^3 + x + 1.$$

The values of $x^3 + x + 1$ for $x = 0, 1, \dots, 6$ are

$$1, 3, 4, 3, 6, 5, 6.$$

Only 1 and 4 are quadratic residues in this list. Therefore

$$E(\mathbb{F}_7) = \{O, (0, 1), (0, 6), (2, 2), (2, 5)\}, \quad \#E(\mathbb{F}_7) = 5.$$

 **Exercise 1.15.** Count the points on each curve over $\mathbb{F}_7$:


$$y^2 = x^3 - x, \quad y^2 = x^3 + 3.$$

Check that their point counts are 8 and 13, respectively.

1.3 The group law

Over the real numbers one can picture the group law by drawing a line through two points, finding the third intersection, and reflecting in the x -axis. Over $\mathbb{F}_p$ there is no picture, but the same algebraic formulas work using modular inverses.

Equivalently, three collinear points on a nonsingular cubic add to O (with intersection multiplicities counted). For the real curve in Example 1.12, the line through $P = (-15, 0)$ and $Q = (3, 54)$ also meets the curve at $R = (21, 108)$. Thus $P + Q + R = O$, and $P + Q = (21, -108)$.



Theorem 1.16 (Addition formulas). *Let $E : y^2 = x^3 + ax + b$ be an elliptic curve over $\mathbb{F}_p$.*

1. *O is the identity: $P + O = O + P = P$.*
2. *The inverse of (x, y) is $-(x, y) = (x, -y)$.*
3. *If $P = (x_1, y_1)$ and $Q = (x_2, y_2)$ with $x_1 \neq x_2$, put*

$$m = \frac{y_2 - y_1}{x_2 - x_1}, \quad x_3 = m^2 - x_1 - x_2, \quad y_3 = m(x_1 - x_3) - y_1.$$

Then $P + Q = (x_3, y_3)$.

4. *If $P = (x_1, y_1)$ with $y_1 \neq 0$, put*

$$m = \frac{3x_1^2 + a}{2y_1}, \quad x_3 = m^2 - 2x_1, \quad y_3 = m(x_1 - x_3) - y_1.$$

Then $[2]P = (x_3, y_3)$.

5. *If $Q = -P$, including the case $y_1 = y_2 = 0$, then $P + Q = O$.*

With these rules, $(E(\mathbb{F}_p), +)$ is a finite abelian group.

The associativity assertion is the hard part of this theorem; it follows from the projective geometry of a nonsingular cubic. A proof of the group law and the formulas appears in [Sil09, Ch. III, §§2–3]. We will use the theorem, not attempt that proof here.

Example 1.17. Continue with Example 1.14. For $P = (0, 1)$, doubling gives

$$m = \frac{1}{2} = 4 \pmod{7}, \quad [2]P = (2, 5).$$

Adding P to $(2, 5)$ gives $[3]P = (2, 2)$, then $[4]P = (0, 6)$, and finally $[5]P = O$. Thus P generates all of $E(\mathbb{F}_7)$.

Definition 1.18. The scalar multiplication $[n]P$ is computed efficiently by double-and-add: write n in binary, repeatedly double the current point, and add only at the binary digits equal to 1. This needs only about $\log_2 n$ doublings and additions.

 **Exercise 1.19.** Use double-and-add to compute $[13]P$ for $P = (0, 1)$ on the curve in Example 1.14. Explain why the answer could also have been found by reducing 13 modulo $\text{ord}(P)$.

Remark 1.20. If instead of $\mathbb{F}_p$ we work over the rational numbers $\mathbb{Q}$, then the group $E(\mathbb{Q})$ is finitely generated by Mordell-Weil's theorem. The torsion subgroup is classified by Mazur's theorem, and the rank of the free part is still the subject of current research. It is not even known whether the rank can be arbitrarily large. You can find a list of the largest known ranks at [this link](#).

1.4 How many points can there be?

The elementary bound $\#E(\mathbb{F}_q) \leq 2q + 1$ follows by counting the at most two y -values above each x . Hasse's theorem is vastly sharper; see [Sil09, Ch. V, §1].

Theorem 1.21 (Hasse). *If E is an elliptic curve over $\mathbb{F}_q$, then*

$$q + 1 - 2\sqrt{q} \leq \#E(\mathbb{F}_q) \leq q + 1 + 2\sqrt{q}.$$

For $q = 7$, the possible integer values lie between 3 and 13. This explains why the examples above can have dramatically different point counts, even though both are defined over the same field.

Example 1.22. The two ends of Hasse's interval already occur over $\mathbb{F}_7$:

$$E_1 : y^2 = x^3 - 3 \quad \text{has } \#E_1(\mathbb{F}_7) = 3, \quad E_2 : y^2 = x^3 + 3 \quad \text{has } \#E_2(\mathbb{F}_7) = 13.$$

Thus the variation in point counts is real, even for a very small field. The points are:

$$E_1(\mathbb{F}_7) = \{O, (0, 2), (0, 5)\},$$

$$E_2(\mathbb{F}_7) = \{O, (1, 2), (1, 5), (2, 2), (2, 5), (3, 3), (3, 4), (4, 2), (4, 5), (5, 3), (5, 4), (6, 3), (6, 4)\}.$$

 **Exercise 1.23.** Use your answer to Exercise 1.3 to compute a bound for $\#E(\mathbb{F}_q)$ better than the naive bound $\#E(\mathbb{F}_q) \leq 2q + 1$.

 **Exercise 1.24.** Enumerate the curves $y^2 = x^3 + ax + b$ over $\mathbb{F}_7$ with nonzero discriminant. Verify that the curves in Example 1.22 attain the smallest and largest possible point counts.

 **Exercise 1.25.** For $p = 5, 7, 11$, use SageMath to enumerate all pairs (a, b) with $4a^3 + 27b^2 \neq 0$ and record the possible values of $\#E(\mathbb{F}_p)$. Compare your data with Hasse's interval.

 **Exercise 1.26.** For several primes p , compute $\lfloor p + 1 + 2\sqrt{p} \rfloor$ and enumerate all nonsingular curves $y^2 = x^3 + ax + b$ over $\mathbb{F}_p$. Let $M(p)$ be the number of pairs (a, b) whose curve has this maximal permitted number of points. Record the values of $M(p)$ and look for a pattern.

1.5 Practice problems

The following problems collect the operations used throughout the lecture. Work the first three by hand before checking any computations with SageMath.

 **Exercise 1.27.** Over $\mathbb{F}_{11}$, consider

$$E: y^2 = x^3 + 2x + 7.$$

Compute the discriminant and verify that E is nonsingular. Make a table of the quadratic residues in $\mathbb{F}_{11}$, then use it to enumerate $E(\mathbb{F}_{11})$.

 **Exercise 1.28.** Use your point count for $E/\mathbb{F}_{11}$ to check Hasse's bound. Determine the group structure of $E(\mathbb{F}_{11})$.

These are the same operations implemented by an elliptic-curve cryptographic library. The only difference in practice is the size of the field and the subgroup: production parameters make direct enumeration and trial search infeasible.

Lecture 2: The elliptic-curve discrete logarithm problem

The group law turns an elliptic curve into a setting for public-key cryptography. In this lecture, we will study the theoretical foundations of the elliptic-curve discrete logarithm problem (ECDLP).

2.1 Public-key cryptography

Public-key cryptography lets people who have never met establish secure communication over a public channel. The channel might be the Internet, a wireless network, or any other medium on which an adversary can read the messages. We will call the two people who want to communicate Alice and Bob, and call an adversary Eve. A cryptographic protocol specifies what Alice and Bob send, what they keep secret, and what Eve is allowed to see.

The basic aim is to turn a secret key into something useful. A secret **key** is a short piece of information used by a cryptographic algorithm. For example, a symmetric encryption algorithm uses the *same* secret key to encrypt and decrypt:

$$\text{plaintext} \xrightarrow{\text{encrypt with } K} \text{ciphertext} \xrightarrow{\text{decrypt with } K} \text{plaintext}.$$

Symmetric encryption is fast and is what normally protects the contents of a long message. Its difficulty is key distribution: how can Alice and Bob obtain the common secret key K in the first place if Eve can observe their communication?

Public-key cryptography addresses this problem by giving each participant a pair of related keys:

$$\text{public key} \quad \text{and} \quad \text{private key}.$$

The public key may be published or sent over the public channel. The private key must be sampled randomly and kept only by its owner. The two keys are constructed so that a public key can be computed efficiently from its private key, while recovering the private key from the public key is believed to be computationally infeasible. Security does not rely on hiding the algorithm: the curve, the protocol, and all public parameters are assumed to be known to Eve.

Public-key techniques have two common roles. First, **key agreement** allows Alice and Bob to derive a common symmetric key; Diffie–Hellman is the example studied below. Second, **digital signatures** allow a sender to prove possession of a private key, so that anyone with the corresponding public key can verify the sender’s signature. In practice, a protocol often uses public-key cryptography to authenticate participants and agree on a temporary symmetric key, then uses symmetric encryption to protect the data.

For elliptic-curve cryptography, the relevant one-way-looking operation is *scalar multiplication*. Public parameters include an elliptic curve $E/\mathbb{F}_q$ and a publicly known point P .

Alice chooses a secret integer a (her private key) and computes

$$A = [a]P.$$

She may publish A as her public key. Repeatedly adding P to itself (using the fast double-and-add algorithm) computes $[a]P$ efficiently. The reverse task—given P and A , recover a —is the elliptic-curve discrete logarithm problem. The remainder of this lecture makes this statement precise and explains how it leads to Diffie–Hellman key agreement.

Warning 2.1. Public information is not necessarily harmless information. A public key is safe to reveal only because the associated mathematical problem is hard for the chosen parameters. Tiny examples are excellent for learning the protocols, but their private keys can be found by simply trying all possibilities. Moreover, mathematical hardness alone does not provide every security property: key agreement must be authenticated to prevent an active attacker from impersonating a participant.

2.2 Further structure of the points on elliptic curves

Before we start looking at the discrete logarithm problem, we need to understand the structure of the group of points on an elliptic curve over a finite field. Because $E(\mathbb{F}_q)$ is a finite abelian group, it has a particularly simple form:

Theorem 2.2. *For an elliptic curve $E/\mathbb{F}_q$, there are positive integers N_1, N_2 with $N_2 \mid N_1$ and $N_2 \mid (q - 1)$ such that*

$$E(\mathbb{F}_q) \simeq \mathbb{Z}/N_1\mathbb{Z} \times \mathbb{Z}/N_2\mathbb{Z}.$$

For an elliptic curve E/k and any positive integer N , the N -torsion subgroup is

$$E[N] = \{P \in E(\bar{k}) : [N]P = O\}.$$

If $\text{char}(k)$ does not divide N , then

$$E[N] \simeq \mathbb{Z}/N\mathbb{Z} \times \mathbb{Z}/N\mathbb{Z}.$$

In characteristic p , the p -power torsion is more subtle. For $n \geq 1$, as we will see later, over an algebraic closure $E[p^n]$ is trivial when E is *supersingular* and is isomorphic to $\mathbb{Z}/p^n\mathbb{Z}$ when E is *ordinary*.

 **Exercise 2.3.** Using Sage, compute the point sets and determine the group structures of the following curves over $\mathbb{F}_{23}$:

$$y^2 = x^3 + x + 1, \quad y^2 = x^3 - x, \quad y^2 = x^3 + 5x + 15.$$

 **Exercise 2.4.** Let $p > 3$, and let

$$E : y^2 = x^3 + ax + b, \quad \tilde{E} : y^2 = x^3 + ax - b$$

be elliptic curves over $\mathbb{F}_p$. Show that if $p \equiv 1 \pmod{4}$, then $\#E(\mathbb{F}_p) = \#\tilde{E}(\mathbb{F}_p)$; if $p \equiv 3 \pmod{4}$, show that $\#E(\mathbb{F}_p) + \#\tilde{E}(\mathbb{F}_p) = 2p + 2$. Hint: determine when -1 is a square in $\mathbb{F}_p$.

2.3 The discrete logarithm problem

Now we are ready to study the discrete logarithm problem, a problem that is easy to state, but believed to be hard to solve. The problem is defined in any finite group, but we will focus on the case of elliptic curves over finite fields.

Problem 2.5 (Elliptic-curve discrete logarithm problem, ECDLP). Given an elliptic curve E over a finite field $\mathbb{F}_q$, a point $P \in E(\mathbb{F}_q)$ of prime order n , and a point $Q \in \langle P \rangle$, find the unique $d \in \{0, \dots, n-1\}$ such that

$$Q = [d]P.$$

The integer d is called the discrete logarithm of Q to the base P .

Example 2.6. On the curve of Example 1.14, $P = (0, 1)$ has order 5 and $[3]P = (2, 2)$. Hence the discrete logarithm of $(2, 2)$ to the base P is 3.

 **Exercise 2.7.** Here is Sage Math code to create a larger example. Can you find the discrete logarithm of Q to the base P ?

```
p = Integer(
    "115792089237316195423570985008687907853269984665640"
    "564039457584007908834671663"
)
F = GF(p)
E = EllipticCurve(F, [0, 0, 0, 0, 7])
P = E(Integer("55066263022277343669578718895168534326250603453777594"
    "175500187360389116729240"),
    Integer("3267051002075881697808308513050704318447127338065924327"
    "5938904335757337482424"))
Q = E(Integer("28921421024367332478688415468105168944814824843786637"
    "14526246502832759657938"),
    Integer("9448041122529449226887754910982892170726235696258976831"
    "6934175098490464915373"))
```

The best general classical attacks, such as Pollard rho, use roughly $\sqrt{n}$ group operations and little memory. Therefore a subgroup of order about 2^{256} gives an approximately 128-bit classical security target. Special curves or poorly chosen parameters can be weaker, so practical ECC uses standardized, audited curves and algorithms.

 **Exercise 2.8.** Let P have order n . Show that $[a]P = [b]P$ if and only if $a \equiv b \pmod{n}$. Why does this prove uniqueness in the ECDLP definition?

 **Exercise 2.9.** On the curve and with the generator $P = (0, 1)$ from Example 1.14, write down the table

$$O, P, [2]P, [3]P, [4]P.$$

Use it to find the discrete logarithm of each nonidentity point to the base P . Which part of this method becomes infeasible when the subgroup order is about 2^{256} ?

 **Exercise 2.10** (Transporting a discrete logarithm). Let $\varphi : E \rightarrow E'$ be a group homomorphism, and suppose that P has prime order n and that $\varphi(P)$ also has order n . If $Q = [d]P$, prove that

$$\varphi(Q) = [d]\varphi(P).$$

Show that a solution to the discrete logarithm problem for $(\varphi(P), \varphi(Q))$ recovers d modulo n . Why is the assumption on the order of $\varphi(P)$ essential?

2.4 Diffie–Hellman key agreement

Diffie–Hellman is a *key-agreement* protocol, not an encryption protocol. It lets two parties derive the same shared secret over a public channel in the chosen subgroup.

1. Alice and Bob agree on public parameters $(E, \mathbb{F}_q, P, n)$, where P has prime order n .
2. Alice samples $a \in \{1, \dots, n-1\}$ uniformly and sends $A = [a]P$.
3. Bob samples $b \in \{1, \dots, n-1\}$ uniformly and sends $B = [b]P$.
4. Alice computes $S_A = [a]B$ and Bob computes $S_B = [b]A$.

Both obtain the same point

$$S_A = [a]([b]P) = [ab]P = [b]([a]P) = S_B.$$

The protocol uses a key-derivation function (KDF) on an agreed encoding of S_A , together with the public transcript, to obtain symmetric key material; the point P is not used directly as an encryption key.

Example 2.11 (A toy ECDH exchange). On the curve in Example 1.14, take the generator $P = (0, 1)$ of order 5. If Alice chooses $a = 2$ and Bob chooses $b = 3$, then

$$A = [2]P = (2, 5), \quad B = [3]P = (2, 2).$$

Alice computes $[2]B = [6]P = P$, while Bob computes $[3]A = [6]P = P$. Thus they obtain the same shared point. The calculation also makes clear why the scalars are read modulo 5: here $6 \equiv 1 \pmod{5}$.

Remark 2.12. One might try to imitate the protocol with a public nonzero integer p : Alice sends $A = ap$ and Bob sends $B = bp$, and they compute

$$aB = bA = abp.$$

This gives no secrecy. An eavesdropper who knows p , A , and B can recover $a = A/p$ and $b = B/p$, or compute the shared value directly as AB/p . Thus multiplication by a public integer is efficiently reversible by division. In elliptic-curve Diffie–Hellman, by contrast, recovering a from P and $[a]P$ is the presumed-hard discrete-logarithm problem.

 **Exercise 2.13.** Suppose an implementation accepts the point $B = O$ as Bob’s public message. What shared point does Alice compute, regardless of her secret scalar a ? Explain why checking subgroup membership alone does not rule out this case, and state an additional check the implementation should make.

 **Exercise 2.14.** In the subgroup generated by the point P of Example 1.14, let Alice choose $a = 2$ and Bob choose $b = 3$. Compute A , B , and the shared point.

 **Exercise 2.15.** Suppose that a generic discrete-logarithm attack requires approximately $\sqrt{n}$ group operations in a subgroup of prime order n .

1. Estimate the work factor when n is about 2^{256} .
2. If an attacker can carry out 2^{40} group operations per second, estimate, in years, how long 2^{128} operations would take.
3. Explain why this estimate motivates describing such a subgroup as offering roughly 128 bits of classical security.

Warning 2.16. Unauthenticated Diffie–Hellman does *not* authenticate Alice or Bob. An active attacker can replace A and B and mount a man-in-the-middle attack. Real protocols authenticate the exchange, for example with signatures, certificates, or a pre-shared authentication mechanism. Implementations must also validate received public points and handle cofactors according to the chosen curve protocol.

 **Exercise 2.17.** Alice sends $A = [a]P$ and Bob sends $B = [b]P$, but an active adversary Eve replaces Alice's message by $A' = [e_A]P$ and Bob's message by $B' = [e_B]P$, for scalars chosen by Eve.

1. Determine the point computed by Alice and the point computed by Bob.
2. Determine the two points Eve can compute from A , B , e_A , and e_B .
3. Explain how Eve can use these two shared points to read and alter messages if Alice and Bob do not authenticate the exchange.

2.5 What is assumed to be hard?

The computational Diffie–Hellman problem (CDH) in the subgroup $\langle P \rangle$ is: given

$$A = [a]P \quad \text{and} \quad B = [b]P,$$

compute $[ab]P$. The security of unauthenticated ECDH against a passive eavesdropper is modeled by the difficulty of this problem.

An algorithm for ECDLP solves CDH: recover a from (P, A) and then compute $[a]B$. The converse implication is not known in complete generality, so the two assumptions should not be identified. For well-chosen elliptic-curve groups, no general classical algorithm is known that is asymptotically faster than generic square-root search.

One curve that is widely used is the one recommended by the Internet Research Task Force (IRTF) given by the following equation over $\mathbb{F}_p$ with $p = 2^{255} - 19$:

$$E : y^2 = x^3 + 486662x^2 + x.$$

The base point recommended for use has coordinates $x = 9$ and $y =$

$$14781619447589544791020593568409986887264606134616475288964881837755586237401.$$

The order of this point is a prime number

$$n = 2^{252} + 27742317777372353535851937790883648493.$$

Moreover, NIST, the US National Institute of Standards and Technology, has a list of 15 elliptic curves that are recommended for use in cryptography. The curves are defined over prime fields $\mathbb{F}_p$ and binary fields $\mathbb{F}_{2^m}$, and have been chosen to have a prime number of points or a small cofactor.

Lecture 3: Isogenies of elliptic curves

The Diffie–Hellman key-agreement protocol studied in the previous lecture is not resistant to attacks by sufficiently powerful fault-tolerant quantum computers: Shor’s algorithm can solve the discrete logarithm problem on which it relies. This motivates the search for new post-quantum public-key encryption and key-agreement protocols. Isogenies provide one mathematical setting in which such protocols have been explored, and this lecture develops the basic language needed to study them.

The definitions in this section are motivated from the point arithmetic already developed, but some proofs use algebraic geometry and are stated without proof. The important new idea is that an isogeny is a structure-preserving map from one elliptic curve to another.

3.1 Maps between elliptic curves

Definition 3.1. Let E and E' be elliptic curves over a field $\mathbb{F}_p$. In these notes, a map between E and E' is a function $\varphi : E \rightarrow E'$ given by formulas

$$\varphi(x, y) = (f(x, y), g(x, y)),$$

where f and g are quotients of polynomials in x and y . Away from zeros of their denominators, the formulas must send points of E to points of E' . We also require that they extend to a well-defined value in E' at every affine point where a denominator vanishes, and that the map send the point at infinity O to O .

Example 3.2. Consider the curves $E : y^2 = x^3 - x$ and $E' : y^2 = x^3 + 4x$ over $\mathbb{F}_7$. For $x \neq 0$, put

$$\varphi(x, y) = \left(x - \frac{1}{x}, y \left(1 + \frac{1}{x^2} \right) \right),$$

and set $\varphi(O) = \varphi(0, 0) = O$. If $(x, y) \in E$ and $x \neq 0$, then

$$\left(y \left(1 + \frac{1}{x^2} \right) \right)^2 = \left(x - \frac{1}{x} \right)^3 + 4 \left(x - \frac{1}{x} \right),$$

so $\varphi(x, y)$ lies on E' . Thus these formulas define a map $E \rightarrow E'$. For instance, $(4, 2) \in E(\mathbb{F}_7)$ is sent to $(2, 3) \in E'(\mathbb{F}_7)$. We will recover this map from Vélu’s formulas in Example 3.12.

Example 3.3. An example of something that is not a map is the following. Let $E : y^2 = x^3 + x$ and $E' : y^2 = x^3 + 2x$ over $\mathbb{F}_7$. Consider the formulas

$$\varphi(x, y) = (2x, y).$$

This is not a map because it does not send points of E to points of E' . For instance, $(1, 3) \in E(\mathbb{F}_7)$ is sent to $(2, 3)$, which is not a point of $E'(\mathbb{F}_7)$.

3.1.1 Isomorphisms and the j -invariant

Definition 3.4. Let $E : y^2 = x^3 + ax + b$ and $E' : y^2 = x^3 + a'x + b'$ be short Weierstrass curves over $\mathbb{F}_p$. They are isomorphic over $\mathbb{F}_p$ if there is a $u \in \mathbb{F}_p^\times$ such that

$$a' = u^4 a, \quad b' = u^6 b.$$

The isomorphism sends (x, y) to $(u^2 x, u^3 y)$ and sends O to O .

Definition 3.5. The j -invariant of E is

$$j(E) = 1728 \frac{4a^3}{4a^3 + 27b^2} = \frac{256 \cdot 27a^3}{4a^3 + 27b^2}.$$

It is unchanged by isomorphism. Over an algebraic closure, two elliptic curves of characteristic different from 2 and 3 are isomorphic exactly when their j -invariants agree.

Example 3.6. Over $\mathbb{F}_7$, the curves $y^2 = x^3 + x$ and $y^2 = x^3 + 2x$ are isomorphic: take $u = 2$, for which $u^4 = 2$ in $\mathbb{F}_7$. The map is $(x, y) \mapsto (4x, y)$.

 **Exercise 3.7.** Verify directly that $(x, y) \mapsto (4x, y)$ sends points of $y^2 = x^3 + x$ over $\mathbb{F}_7$ to points of $y^2 = x^3 + 2x$. Find its inverse map, and compute the common j -invariant in $\mathbb{F}_7$.

 **Exercise 3.8.** Decide if the following statement is true or false: “Every isomorphism of elliptic curves is also a group isomorphism”.

3.2 Isogenies

Isomorphisms identify curves that have exactly the same elliptic-curve structure. We now broaden the comparison: an isogeny is a map between possibly different curves that preserves the group law, while allowing a finite kernel.

Definition 3.9. Let E and E' be elliptic curves over a field $\mathbb{F}_p$. An **isogeny** is a nonconstant such map of elliptic curves defined over $\mathbb{F}_p$ that respects the elliptic-curve group law:

$$\varphi(P + Q) = \varphi(P) + \varphi(Q).$$

Its **kernel** is $\ker \varphi = \{P \in E(\overline{\mathbb{F}_p}) : \varphi(P) = O\}$, where $\overline{\mathbb{F}_p}$ denotes an algebraic closure of $\mathbb{F}_p$. The **degree** of φ is its degree as a rational map. If φ is separable, then its degree is the size of its finite kernel.

For short Weierstrass models (in characteristic different from 2 and 3), write

$$E : y^2 = x^3 + ax + b, \quad E' : y'^2 = x'^3 + a'x' + b'.$$

Away from the finite set of points where a formula has a pole, an isogeny can be written in coordinates as

$$\varphi(x, y) = (r(x), y s(x))$$

for rational functions $r(x), s(x) \in \mathbb{F}_p(x)$, with $\varphi(O) = O$. The functions are not arbitrary: substituting into the equation of E' gives the identity

$$s(x)^2(x^3 + ax + b) = r(x)^3 + a'r(x) + b'.$$

Thus the coordinate formula sends points of E to points of E' . The additional group-law condition says that it also sends sums of points to sums of their images.

Example 3.10. Let $E: y^2 = x^3 - x$ defined over a prime field k of characteristic different from 2 and 3. When $y \neq 0$, we have the map $[2] : E \rightarrow E$ that sends P to $[2]P$. In coordinates, it is

$$[2](x, y) = (r(x), y s(x)),$$

where $a = -1$ and $b = 0$, and

$$r(x) = \frac{x^4 - 2ax^2 - 8bx + a^2}{4(x^3 + ax + b)}, \quad s(x) = -1 + \frac{(3x^2 + a)(x - r(x))}{2(x^3 + ax + b)} \in k(x).$$

Thus $[2]$ is algebraic: on this open set its coordinates are rational functions in x and y . Multiplication by 2 sends O to O . It is nonconstant because the numerator of $r(x)$ has degree 4, while its denominator has degree 3; hence $r(x)$ cannot be a constant rational function.

The map $[2] : E \rightarrow E$ is therefore an isogeny. Since the characteristic is different from 2, the map $[2]$ is separable. Its kernel is the set of points over $\bar{k}$ such that $[2]P = O$, namely the 2-torsion subgroup of E . The points $(0, 0)$, $(1, 0)$, and $(-1, 0)$ are the three nonidentity points of $E[2]$. Thus

$$E[2] = \{O, (0, 0), (1, 0), (-1, 0)\},$$

and $[2]$ has degree 4.

More generally, the nonidentity points in the kernel of $[2]$ on a short Weierstrass curve are the three points with $y = 0$ over $\bar{k}$; their x -coordinates are the roots of the cubic on the right-hand side. Even if none of those roots is in k , the kernel still has four points over $\bar{k}$. This is why the definition of a kernel uses an algebraic closure.

3.2.1 Subgroups and isogenies

Theorem 3.11. *If $G \subseteq E(\bar{k})$ is a finite subgroup stable under the relevant field automorphisms, then there is an elliptic curve E/G and an isogeny*

$$\pi_G : E \longrightarrow E/G$$

with kernel G . In explicit computations, Vélu's formulas construct E/G and π_G from the points of G .

Vélu's formulas make the preceding theorem explicit. Suppose that $\text{char}(k) \neq 2, 3$, that

$$E: y^2 = x^3 + ax + b,$$

and let $G \subseteq E(\bar{k})$ be a finite subgroup stable under the relevant field automorphisms. Put $G^* = G \setminus \{O\}$. Then a short Weierstrass model for the quotient is

$$E/G: Y^2 = X^3 + a_G X + b_G,$$

where

$$\begin{aligned} a_G &= a - 5 \sum_{Q \in G^*} (3x(Q)^2 + a), \\ b_G &= b - 7 \sum_{Q \in G^*} (5x(Q)^3 + 3a x(Q) + 2b). \end{aligned}$$

For an affine point P for which the expressions are finite, the quotient map is given by

$$\begin{aligned} \pi_G(P) &= (X(P), Y(P)), \\ X(P) &= x(P) + \sum_{Q \in G^*} (x(P+Q) - x(Q)), \\ Y(P) &= y(P) + \sum_{Q \in G^*} y(P+Q). \end{aligned}$$

These rational expressions extend to all of E , with every point of G mapping to O . Thus, given the points of the kernel, one can compute both the neighboring curve and the isogeny to it; see Vélu [V71].

Example 3.12. Let $E: y^2 = x^3 - x$ over $\mathbb{F}_7$. Its nonidentity 2-torsion points are $(0,0)$, $(1,0)$, and $(6,0)$. Take $G = \langle (0,0) \rangle$, so that $G^* = \{Q\}$ with $Q = (0,0)$. Here $a = -1$ and $b = 0$. Vélu's coefficient formulas give

$$a_G = -1 - 5(3 \cdot 0^2 - 1) = 4, \quad b_G = 0 - 7(5 \cdot 0^3 + 3(-1) \cdot 0 + 2 \cdot 0) = 0 \quad \text{in } \mathbb{F}_7.$$

Thus the quotient curve computed by Vélu's formulas is

$$E/G: y^2 = x^3 + 4x.$$

Now let $P = (x, y)$ with $x \neq 0$. The addition formula with $Q = (0,0)$ gives

$$x(P+Q) = \left(\frac{y}{x}\right)^2 - x = -\frac{1}{x}, \quad y(P+Q) = \frac{y}{x} \left(x + \frac{1}{x}\right) - y = \frac{y}{x^2}.$$

Since $x(Q) = y(Q) = 0$, Vélu's coordinate formulas therefore give

$$\pi_G(x, y) = \left(x + x(P+Q), y + y(P+Q)\right) = \left(x - \frac{1}{x}, y \left(1 + \frac{1}{x^2}\right)\right).$$

The formulas extend with $\pi_G(O) = \pi_G(0,0) = O$. This formula is visibly nonconstant, and direct substitution gives

$$y^2 \left(1 + \frac{1}{x^2}\right)^2 = \left(x - \frac{1}{x}\right)^3 + 4 \left(x - \frac{1}{x}\right),$$

so it sends points of E to points of E/G . Vélu's construction shows that it preserves the group law; hence it is a 2-isogeny with kernel $\{O, (0, 0)\}$. For instance,

$$\pi_G(4, 2) = (2, 3).$$

Remark 3.13. You can define isogenies in Sage as quotients. For example, the isogeny in Example 3.12 can be computed as follows:

```
F = GF(7)
E = EllipticCurve(F, [-1, 0])
P = E(0, 0) # Generator of the kernel
phi = E.isogeny([P])
```

The output is the isogeny π_G with kernel $\langle P \rangle$. You can evaluate it on points of E and check that it sends them to points of the codomain curve. To find the codomain curve, you can use `phi.codomain()`.

 **Exercise 3.14.** Let P have order ℓ . Why is $\langle P \rangle$ an allowable candidate for the kernel of an ℓ -isogeny when it is defined over the base field? What information must Vélu's formulas take as input?

 **Exercise 3.15.** Find all curves over $\mathbb{F}_7$ that are isogenous to $y^2 = x^3 - x$. For each, also find the degree of the isogeny.

3.3 Isogenies in cryptography: SIDH

SIDH (Supersingular Isogeny Diffie–Hellman) is a public-key key-agreement protocol. Historically, it began with a supersingular curve $E/\mathbb{F}_{p^2}$ carrying two public torsion bases (P_A, Q_A) and (P_B, Q_B) of coprime prime-power orders. Alice chooses a secret cyclic subgroup generated by $P_A + [a]Q_A$, computes the corresponding quotient isogeny $\varphi_A : E \rightarrow E_A$, and publishes E_A together with $\varphi_A(P_B)$ and $\varphi_A(Q_B)$; Bob does the analogous computation with his own secret subgroup. The published images let each party transport the other secret kernel through their own isogeny, so both obtain isomorphic quotient curves.

For roughly a decade after its introduction, no practical method was known to reconstruct the secret isogeny from the public curve and these auxiliary images, so SIDH was regarded as a promising post-quantum proposal. In 2022, Castryck and Decru showed that the images could be combined with a special endomorphism of the starting curve to recover the secret key efficiently; SIDH and its key-encapsulation mechanism SIKE are therefore broken and must not be used [CD22].

Lecture 4: ℓ -Isogenies and ordinary/supersingular curves

Now we are ready to study isogeny graphs, which are the main objects of interest in isogeny-based cryptography. We will see that they have a rich arithmetic structure and that they can be used to construct cryptographic protocols that are believed to be secure against quantum attacks.

4.1 ℓ -isogenies

We have already seen that isogenies are maps between elliptic curves that respect the group law. We now introduce a special class of isogenies that will be particularly useful in cryptography.

Definition 4.1. If ℓ is a prime different from the characteristic of $\mathbb{F}_q$, an ℓ -isogeny is an isogeny whose kernel

$$\ker \varphi = \{P \in E(\overline{\mathbb{F}}_q) : \varphi(P) = O\}$$

has exactly ℓ points. This is equivalent to saying that φ has degree ℓ .

Example 4.2. Consider the multiplication-by-2 isogeny $\varphi: E \rightarrow E$ from Example 3.10, where

$$E : y^2 = x^3 - x.$$

A point belongs to $\ker \varphi$ exactly when doubling it gives O ; in other words, it has order dividing 2. Besides O , these are precisely the points with $y = 0$. Since

$$x^3 - x = x(x-1)(x+1),$$

we obtain

$$\ker \varphi = \{O, (0, 0), (1, 0), (-1, 0)\}.$$

The kernel has four points, so φ has degree 4. In particular, multiplication by 2 is *not* a 2-isogeny: the 2 in φ describes the group operation, whereas a 2-isogeny has a kernel of size 2.

Example 4.3. Over $\mathbb{F}_7$, consider the curves

$$E : y^2 = x^3 - x, \quad E_1 : y^2 = x^3 + 4x.$$

Example 3.12 constructs an isogeny $\varphi_1 : E \rightarrow E_1$ given, for $x \neq 0$, by

$$\varphi_1(x, y) = \left(x - \frac{1}{x}, y \left(1 + \frac{1}{x^2} \right) \right),$$

together with $\varphi_1(O) = \varphi_1(0, 0) = O$.

Note that no point with $x \neq 0$ is sent to O . Hence

$$\ker \varphi_1 = \{O, (0, 0)\},$$

and φ_1 is a 2-isogeny.

 **Exercise 4.4.** Here are two more isogenies of the elliptic curve E from Example 4.3 (you do not need to prove that they are isogenies). Show that they both have degree 2.

$$\begin{aligned} \varphi_2: \quad E &\rightarrow E_2: y^2 = x^3 + 3x \\ (x, y) &\mapsto \left(\frac{x^2 + x + 2}{x + 1}, \frac{x^2 y + 2xy - y}{x^2 + 2x + 1} \right) \end{aligned}$$

$$\begin{aligned} \varphi_3: \quad E &\rightarrow E_2: y^2 = x^3 + 3x \\ (x, y) &\mapsto \left(\frac{x^2 - x + 2}{x - 1}, \frac{x^2 y - 2xy - y}{x^2 - 2x + 1} \right) \end{aligned}$$

Remark 4.5. It turns out that the isogenies from Example 4.3 and Exercise 4.4 are all of the 2-isogenies of E defined over $\mathbb{F}_7$.

 **Exercise 4.6.** Let E be an elliptic curve defined over $\mathbb{F}_p$ and φ be an isogeny of E of degree ℓ . If n is a positive integer coprime to ℓ , show that the restriction of φ to a subgroup of order n is injective.

4.2 Ordinary and supersingular elliptic curves

Not all curves are equally useful for isogeny-based constructions. The following definition distinguishes two important classes of curves in characteristic p .

Definition 4.7. For an elliptic curve E over $\mathbb{F}_p$, define its **Frobenius trace** by

$$a_p = p + 1 - \#E(\mathbb{F}_p).$$

The curve is **ordinary** if $p \nmid a_p$, and **supersingular** if $p \mid a_p$.

This is a practical definition: we can determine the type by counting points, which we already know how to do. Supersingular curves are central to several isogeny-based constructions because their endomorphisms and isogeny graphs have unusually rigid structure.

Example 4.8. Consider the elliptic curve $E: y^2 = x^3 - x$ over $\mathbb{F}_7$. We have computed that $\#E(\mathbb{F}_7) = 8$, so

$$a_7 = 7 + 1 - 8 = 0.$$

Since $7 \mid a_7$, the curve E is supersingular.

Example 4.9. Over $\mathbb{F}_5$, direct point counting gives

$$\#\{y^2 = x^3 + 1\}(\mathbb{F}_5) = 6, \quad \#\{y^2 = x^3 + x\}(\mathbb{F}_5) = 4.$$

The first curve is supersingular and the second is ordinary. Their Frobenius traces $a_5 = 5 + 1 - \#E(\mathbb{F}_5)$ are respectively 0 and 2. The point-counting criterion used here agrees with the standard p -torsion definition; see [Sil09, Ch. V, §4].

4.3 Isogenies and supersingular elliptic curves

An isogeny can connect curves given by quite different equations, so it is not obvious from the definition whether it preserves the ordinary/supersingular distinction. Our goal in this section is to show that it does: if two curves over $\mathbb{F}_p$ are isogenous over $\mathbb{F}_p$, then either both are ordinary or both are supersingular. With the point-counting criterion, it is enough to prove that isogenous curves have the same number of points over $\mathbb{F}_p$. The following theorem gives exactly this.

Theorem 4.10 (Tate). *Two elliptic curves over $\mathbb{F}_q$ are $\mathbb{F}_q$ -isogenous if and only if they have the same number of $\mathbb{F}_q$ -points.*

For $q = p$, this proves the claimed invariance: equal point counts give equal Frobenius traces, and hence the same ordinary/supersingular type.

The proof is beyond the scope of this course, but we can check an example. Try the following SageMath code to see which curves over $\mathbb{F}_7$ are isogenous to $y^2 = x^3 - x$.

```
F = GF(7)

E = EllipticCurve(F, [-1, 0]) # y^2 = x^3 - x

for a in F:
    for b in F:
        if 4*a^3 + 27*b^2 != 0:
            Eab = EllipticCurve(F, [a, b])

            if Eab.is_isogenous(E):
                print("Isogenous: a =", a, ", b =", b)
            else:
                print("Not isogenous: a =", a, ", b =", b)

print(Eab.cardinality(), "points in Eab;",
      E.cardinality(), "points in E")
print()
```

All supersingular elliptic curves over $\bar{\mathbb{F}}_p$ belong to one isogeny class. This does *not* mean that every two vertices are joined by a single ℓ -isogeny: generally they are connected by a path whose total degree is a power of ℓ .

Corollary 4.11. *Let $\varphi: E \rightarrow E'$ be an isogeny over $\mathbb{F}_p$ of elliptic curves. Then E is supersingular if and only if E' is supersingular (and E is ordinary if and only if E' is ordinary).*

Proof. By Theorem 4.10, we have

$$\#E(\mathbb{F}_p) = \#E'(\mathbb{F}_p).$$

Thus E and E' have the same Frobenius trace. Therefore p divides the trace of E if and only if it divides the trace of E' , proving the supersingular statement. The ordinary statement is its negation. $\square$

 **Exercise 4.12.** The following elliptic curves are all defined over $\mathbb{F}_7$. Determine which are ordinary and which are supersingular.

$$\begin{array}{ll} E_1 : y^2 = x^3 + 1, & E_2 : y^2 = x^3 + x + 3, \\ E_3 : y^2 = x^3 + 6x, & E_4 : y^2 = x^3 + 2x + 1, \\ E_5 : y^2 = x^3 + 3, & E_6 : y^2 = x^3 + 4x + 4. \end{array}$$

It might help you to compute the number of points on each curve for which the Sage command is `E.cardinality()` and to check for isogenies with the command `E.is_isogenous(E2)`.

Lecture 5: Isogeny graphs and cryptography

For cryptography, we need a setting in which individual operations are easy to compute but recovering hidden information is difficult. A small ℓ -isogeny can be computed from its kernel, so it is natural to regard it as one efficient step from its source curve to its target curve. A secret can then be represented by a sequence of such steps, while an adversary sees only the initial and final curves. Recording all possible small steps at once gives an isogeny graph.

5.1 Isogeny graphs


Definition 5.1. Fix primes p and $\ell \neq p$. The ℓ -isogeny graph over $\mathbb{F}_p$ is the directed multigraph whose vertices are the $\mathbb{F}_p$ -isomorphism classes of elliptic curves over $\mathbb{F}_p$. For every cyclic subgroup $G \subseteq E[\ell]$ of order ℓ that is the kernel of an $\mathbb{F}_p$ -defined ℓ -isogeny

$$\pi_G : E \longrightarrow E',$$

it has an edge from the class of E to the class of E' .

When convenient, a vertex is labelled by its j -invariant.

Example 5.2. Let $E : y^2 = x^3 - x$ over $\mathbb{F}_7$. Take vertices up to $\mathbb{F}_7$ -isomorphism and draw only 2-isogenies defined over $\mathbb{F}_7$. By Example 4.8, E is supersingular. Example 4.3 and Exercise 4.4 give its three 2-isogenies. The target $E_2 : y^2 = x^3 + 3x$ is $\mathbb{F}_7$ -isomorphic to E , whereas $E_1 : y^2 = x^3 + 4x$ is not. Thus φ_2 and φ_3 give two loops at the vertex E , while φ_1 gives an edge from E to E_1 . The 2-isogeny from E_1 with kernel $\{O, (0,0)\}$ has target $y^2 = x^3 + 5x$, which is $\mathbb{F}_7$ -isomorphic to E . The resulting $\mathbb{F}_7$ -rational graph is therefore



The two loops have kernels generated by $(1,0)$ and $(-1,0)$; the two directed edges between the vertices have kernel generated by $(0,0)$ on their respective source curves.

We can obtain this graph with Sage:

```

E = EllipticCurve(GF(7), [-1,0])
graph = E.isogeny_ell_graph(2)
graph.plot()

```

Example 5.3. The $\mathbb{F}_p$ -isogeny graph need not be connected. Take $p = 19$ and $\ell = 3$. There are four $\mathbb{F}_{19}$ -isomorphism classes of supersingular elliptic curves over $\mathbb{F}_{19}$, namely 7, 7', 18, and 18'. A prime on a label distinguishes the two classes with the same j -invariant.

In this case, one can show that there are no $\mathbb{F}_{19}$ -defined 3-isogenies between the supersingular curves. The resulting graph therefore consists of four isolated vertices:



In particular, this $\mathbb{F}_{19}$ -rational supersingular 3-isogeny graph is disconnected.

 **Exercise 5.4.** Write an algorithm to compute the 2-isogeny graph over $\mathbb{F}_p$ of an elliptic curve E over $\mathbb{F}_p$. For that, you need to respond to the following questions.

1. How to enumerate all of the relevant isomorphism classes?
2. Given an elliptic curve, how do you find all 2-isogenies from that elliptic curve?
3. Given a supersingular elliptic curve that you know is isomorphic to one of the vertices of your graph, how do you find which vertex it is isomorphic to?

Definition 5.5. Fix primes p and $\ell \neq p$. The **supersingular ℓ -isogeny graph** over $\mathbb{F}_p$ has one vertex for each $\mathbb{F}_p$ -isomorphism class of supersingular elliptic curves over $\mathbb{F}_p$. For every cyclic subgroup $G \subseteq E[\ell]$ of order ℓ that is the kernel of an $\mathbb{F}_p$ -defined ℓ -isogeny $\pi_G : E \rightarrow E'$, it has an edge from the class of E to the class of E' . Distinct kernels give distinct edges.

Remark 5.6. By Corollary 4.11, isogenies from supersingular elliptic curves also go to supersingular elliptic curves. This implies that the edges of the graph from Definition 5.5 are well-defined (since both endpoints are vertices of the graph).

Remark 5.7. We pick supersingular elliptic curves since for ordinary curves, the structure of the isogeny graphs is well-understood. This structure is known as isogeny volcanoes [Koh96].

Definition 5.8. Fix primes p and $\ell \neq p$. The **supersingular ℓ -isogeny graph** over $\mathbb{F}_{p^2}$ has one vertex for each supersingular j -invariant in $\mathbb{F}_{p^2}$. For every cyclic subgroup $G \subseteq E[\ell]$ of order ℓ , it has an edge from $j(E)$ to $j(E/G)$.

Every supersingular j -invariant in characteristic p lies in $\mathbb{F}_{p^2}$. Since $\ell \neq p$, there are $\ell + 1$ cyclic subgroups of $E[\ell]$ of order ℓ , so this graph has $\ell + 1$ outgoing edges at every vertex. The $\mathbb{F}_p$ -rational graph in Definition 5.5 need not have this property, because not every subgroup of $E[\ell]$ is defined over $\mathbb{F}_p$.

 **Exercise 5.9.** Let $E: y^2 = x^3 + ax + b$ be an elliptic curve defined over $\mathbb{F}_p$.

1. First suppose that $ab \neq 0$. Choose a nonquadratic residue $d \in \mathbb{F}_p^\times$ and define the quadratic twist of E by

$$E^{(d)}: y^2 = x^3 + d^2ax + d^3b.$$

Show that $E^{(d)}$ is an elliptic curve over $\mathbb{F}_p$ and that $j(E^{(d)}) = j(E)$.

2. Let $u \in \mathbb{F}_{p^2}$ satisfy $u^2 = d$. Verify that the following map is an isomorphism over $\mathbb{F}_{p^2}$:

$$E^{(d)} \longrightarrow E, \quad (x, y) \longmapsto \left(\frac{x}{u^2}, \frac{y}{u^3} \right).$$

3. Prove that E and $E^{(d)}$ are not isomorphic over $\mathbb{F}_p$. You may use that an isomorphism from E to $E^{(d)}$ between short Weierstrass equations in characteristic different from 2 and 3 has the form


$$(x, y) \longmapsto (v^2x, v^3y).$$

Compare the coefficients of x and of the constant term to show that $v^2 = d$, contradicting the choice of d .


4. The only cases left are those with $ab = 0$, or equivalently $j(E) \in \{0, 1728\}$. Treat the exceptional cases as follows. If $b = 0$, choose $c \in \mathbb{F}_p^\times$ that is not a fourth power and compare E with $E_c: y^2 = x^3 + cax$. If $a = 0$, choose $c \in \mathbb{F}_p^\times$ that is not a sixth power and compare E with $E_c: y^2 = x^3 + cb$.

By Exercise 5.9, every j -invariant gives at least two $\mathbb{F}_p$ -isomorphism classes of elliptic curves. When $j \notin \{0, 1728\}$, the second class can be represented by a quadratic twist. We can see this in the following example.

Example 5.10. For $p = 31$ and $\ell = 2$, there are three supersingular j -invariants, namely 23, 2, and 4. Each has two $\mathbb{F}_{31}$ -isomorphism classes. For $j = 2$ and $j = 4$, the two classes are represented by a curve and its quadratic twist. Since $23 = 1728$ in $\mathbb{F}_{31}$, the two classes with $j = 23$ are instead represented by a curve and a quartic twist. Thus the $\mathbb{F}_{31}$ -rational supersingular 2-isogeny graph has six vertices. A prime on a label distinguishes the second $\mathbb{F}_{31}$ -isomorphism class with the same j -invariant. Each pair of opposite arrows represents the two dual 2-isogenies defined over $\mathbb{F}_{31}$.



Example 5.11. We now pass from the $\mathbb{F}_p$ -rational graph to the supersingular ℓ -isogeny graph over $\mathbb{F}_{p^2}$. Its vertices are labelled by supersingular j -invariants. In these examples, the number of displayed vertices reduces.



For $p = 19$ and $\ell = 3$, the supersingular j -invariants are 18 and 7. There are four edges from 18 to 7, two edges from 7 to 18, and two loops at 7:

5.1.1 Properties of isogeny graphs

The field of definition and the vertex convention are both part of the data of an isogeny graph. We distinguish the $\mathbb{F}_p$ -rational graph from the supersingular ℓ -isogeny graph over $\mathbb{F}_{p^2}$ of Definition 5.8.

Over $\mathbb{F}_p$.

- Vertices are $\mathbb{F}_p$ -isomorphism classes. A curve and its quadratic twist therefore usually give two distinct vertices with the same j -invariant.
- An isogeny defined over $\mathbb{F}_p$ has a dual isogeny defined over $\mathbb{F}_p$, so every directed edge has a reverse directed edge.

Over $\mathbb{F}_{p^2}$.

- Vertices are the supersingular j -invariants, all of which lie in $\mathbb{F}_{p^2}$.
- Every cyclic subgroup of $E[\ell]$ of order ℓ gives an edge.
- Since $E[\ell] \simeq (\mathbb{Z}/\ell\mathbb{Z})^2$ when $\ell \neq p$, there are $\ell + 1$ such subgroups. Hence every vertex has $\ell + 1$ outgoing edges.
- The graph is connected.

 **Exercise 5.12.** Let ℓ be a prime number. Show that $(\mathbb{Z}/\ell\mathbb{Z})^2$ has $\ell + 1$ subgroups of cardinality ℓ .

For supersingular curves, every vertex has a j -invariant in $\mathbb{F}_{p^2}$. This graph is regular, with $\ell + 1$ outgoing edges at every vertex. It is more convenient than the $\mathbb{F}_p$ -rational graph, whose separate twist vertices can make the degrees of vertices vary. For this reason, the $\mathbb{F}_{p^2}$ graph is the usual setting for supersingular-isogeny algorithms and cryptographic constructions.

5.2 Cryptography and isogeny graphs

An isogeny graph supplies many efficiently computable maps between elliptic curves. This fact alone is not a cryptosystem: two arbitrary walks in a graph do not usually commute, so two parties cannot in general combine their secret walks into the same value. A cryptographic construction must specify the family of curves, the permitted isogenies, a way to represent and compose secrets, and a precise hardness assumption. CSIDH is the standard example where this additional structure is present.

5.2.1 CSIDH

CSIDH stands for *Commutative Supersingular Isogeny Diffie–Hellman*. It is a proposed key-agreement protocol. Its central idea is easier to see before discussing the number theory.

Imagine that the isogeny graph has several colours of edges, say colours $\ell_1, \dots, \ell_n$. For each colour, CSIDH gives a party two possible directions: “forward” and “backward”. A private key is a short list of such moves. For example, Alice might secretly choose

one ℓ_3 -move forward, two ℓ_5 -moves backward, and one ℓ_7 -move forward.

Starting from a public curve E_0 , she follows these moves and publishes only the endpoint E_A . Bob does the same with his own secret list and publishes E_B .




Figure 5.13: This is Figure 9 from [this](#) article.

For an arbitrary graph this does *not* give key agreement: Alice's moves followed by Bob's moves need not end at the same vertex as Bob's moves followed by Alice's. CSIDH chooses a very special family of curves for which the permitted moves commute. Consequently, Alice can apply her original secret list to Bob's public curve, and Bob can apply his original secret list to Alice's public curve. Both arrive at the same curve:

$$\begin{array}{ccc}
 E_0 & \xrightarrow{\text{Alice's secret}} & E_A \\
 \downarrow \text{Bob's secret} & & \downarrow \text{Bob's secret} \\
 E_B & \xrightarrow{\text{Alice's secret}} & E_{AB}.
 \end{array}$$

They use an invariant of E_{AB} , such as its j -invariant, as input to a key-derivation function. This is the direct analogue of ECDH: the secret scalars a, b are replaced by secret collections of isogeny moves, and the common point $[ab]P$ is replaced by the common curve E_{AB} .

Where do the commuting moves come from? CSIDH uses a carefully chosen prime

$$p = 4\ell_1 \cdots \ell_n - 1$$

and begins with the supersingular curve

$$E_0: y^2 = x^3 + x$$

over $\mathbb{F}_p$. It works in a special family X of supersingular curves over $\mathbb{F}_p$, usually written as

$$E_A: y^2 = x^3 + Ax^2 + x.$$

Each small prime ℓ_i supplies the two directions described above. Thus a secret can be stored as an exponent vector

$$\mathbf{e} = (e_1, \dots, e_n), \quad e_i \in [-B, B],$$

where $|e_i|$ says how many ℓ_i -moves to take and the sign says which direction. Each individual move is a small prime-degree isogeny, so it can be computed efficiently from a kernel using Vélu's formulas.

The theorem behind this picture is that the ideal class group

$$\text{Cl}(\mathcal{O}), \quad \mathcal{O} = \mathbb{Z}[\sqrt{-p}],$$

acts on X . The ideal class group is abelian, which is exactly the statement that the allowed moves commute. At this stage, students need not compute ideal classes: the important point is what the theorem provides—a consistent way to apply the same secret moves to *any* public curve in X .

Writing f_A and f_B for Alice's and Bob's secret actions, the preceding diagram says

$$f_A * (f_B * E_0) = (f_A f_B) * E_0 = (f_B f_A) * E_0 = f_B * (f_A * E_0).$$

The intended hard problem is the *group-action inverse* (or vectorization) problem: from E_0 and $E_A = f_A * E_0$, recover a suitable secret action f_A , or compute its effect on another public curve. The public keys are curves, not elements of the acting group, so the usual discrete-logarithm formulation does not apply directly. This is not a proof of post-quantum security: CSIDH has specialised classical and quantum attacks, and remains a research construction rather than a deployed standard. See [CLM⁺18] for the full construction.

5.2.2 SQIsign

CSIDH provides key agreement, whereas SQIsign is a signature scheme. SQIsign (“Short Quaternion and Isogeny Signature”) uses a different part of supersingular-isogeny arithmetic: the relation between supersingular curves, isogenies, and ideals in their quaternionic endomorphism rings. At a high level, a public key is a supersingular curve related to a fixed base curve by a hidden isogeny. Knowledge of the corresponding short quaternionic/isogeny data is the signing secret.

To sign, SQIsign applies the Fiat–Shamir transform to an identification protocol. The signer makes a random commitment; the hash of the message and commitment determines a challenge; and the response is short isogeny-and-ideal data whose endpoints and degrees can be checked publicly. Verification therefore uses explicit isogeny computations, while a forger would need to produce the required hidden relation in the supersingular graph. The compactness of the public key and signature is a principal attraction; the necessary endomorphism-ring and ideal arithmetic makes key generation and signing much more intricate than verification. Its security rests on carefully formulated supersingular-isogeny and endomorphism-ring assumptions, not on the CSIDH group-action assumption.

This distinction also matters historically: the 2022 attack that broke SIDH/SIKE exploited the auxiliary torsion-point images published by that key-exchange design; it did not by itself break CSIDH or SQIsign. SQIsign is an active research proposal, not a NIST standard. NIST selected it as one of nine third-round candidates in its additional post-quantum digital-signature process in 2026. See [DFKL⁺20, CD22] and [NIST’s digital-signature project](#) for current status and submission material.

5.3 Extra: how does one describe isogenies?

Fix a supersingular vertex E . The endomorphisms of E are the morphisms from E to itself. They can be added and composed; together they form a ring denoted by

$$\mathrm{End}(E) = \mathrm{End}_{\overline{\mathbb{F}}_p}(E).$$

Every nonzero endomorphism is an isogeny. A theorem of Deuring says that, for a supersingular curve in characteristic p ,

$$\mathrm{End}(E) \otimes_{\mathbb{Z}} \mathbb{Q} \simeq B_{p,\infty},$$

where $B_{p,\infty}$ is a *quaternion algebra* over $\mathbb{Q}$. This algebra is usually noncommutative, which is one source of the rich structure of supersingular isogeny graphs. At a more advanced level, suitable one-sided ideals in the endomorphism ring determine finite kernel subgroups and hence quotient isogenies from E . After the required equivalences, this gives a second description of supersingular curves and isogenies in terms of quaternionic data; see [Voi21, Chs. 13–16] for a rigorous treatment.

Bibliography

- [AM26] Sarah Arpin and Chloe Martindale. Arizona Winter School lectures, 2026. [swc-math.github.io](#). ↑[ii](#).
- [CD22] Wouter Castryck and Thomas Decru. An efficient key recovery attack on SIDH (preliminary version), 2022. Cryptology ePrint Archive, Paper 2022/975. ↑[19](#), [30](#).
- [CLM⁺18] Wouter Castryck, Tanja Lange, Chloe Martindale, Lorenz Panny, and Joost Renes. CSIDH: an efficient post-quantum commutative group action. In *Advances in Cryptology – ASIACRYPT 2018. Part III*, volume 11274 of *Lecture Notes in Computer Science*, pages 395–427. Springer, Cham, 2018. ↑[29](#).
- [DFKL⁺20] Luca De Feo, David Kohel, Antonin Leroux, Christophe Petit, and Benjamin Wesolowski. SQISign: compact post-quantum signatures from quaternions and isogenies. In *Advances in Cryptology – ASIACRYPT 2020. Part III*, volume 12493 of *Lecture Notes in Computer Science*, pages 64–93. Springer, Cham, 2020. ↑[30](#).
- [HPS14] Jeffrey Hoffstein, Jill Pipher, and Joseph H. Silverman. *An introduction to mathematical cryptography*. Undergraduate Texts in Mathematics. Springer, New York, second edition, 2014. ↑[ii](#).
- [Koh96] David Russell Kohel. *Endomorphism rings of elliptic curves over finite fields*. ProQuest LLC, Ann Arbor, MI, 1996. Thesis (Ph.D.)—University of California, Berkeley. ↑[25](#).
- [Kun26] Sabrina Kunzweiler. Preliminary Arizona Winter School (paws) lectures, 2026. [swc-math.github.io](#). ↑[ii](#).
- [Sil09] Joseph H. Silverman. *The arithmetic of elliptic curves*, volume 106 of *Graduate Texts in Mathematics*. Springer, Dordrecht, second edition, 2009. ↑[ii](#), [6](#), [7](#), [22](#).
- [V71] Jacques Vélú. Isogénies entre courbes elliptiques. *C. R. Acad. Sci. Paris Sér. A-B*, 273:A238–A241, 1971. ↑[18](#).
- [Voi21] John Voight. *Quaternion algebras*, volume 288 of *Graduate Texts in Mathematics*. Springer, Cham, 2021. ↑[30](#).

- [Zag87] Don Zagier. Large integral points on elliptic curves. *Math. Comp.*, 48(177):425–436, 1987. [↑4](#).