





ANNEX I
SUMMARY OF PRODUCT CHARACTERISTICS

 = End of contiguous region
 = Header
 = Not in parsing
 = Not in xml

1. NAME OF THE MEDICINAL PRODUCT

elmiron 100 mg hard capsules

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each hard capsule contains 100 mg of pentosan polysulfate sodium.

For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

αmlns → head 4 → text

Hard capsule.

White opaque capsules size 2.

4. CLINICAL PARTICULARS

αmlns → h6 → text

4.1 Therapeutic indications

elmiron is indicated for the treatment of bladder pain syndrome characterized by either glomerulations or Hunner's lesions in adults with moderate to severe pain, urgency and frequency of micturition (see section 4.4).

4.2 Posology and method of administration

Posology

Adults

The recommended dose of pentosan polysulfate sodium is 300 mg/day taken as one 100 mg capsule orally three times daily.

Response to treatment with pentosan polysulfate sodium should be reassessed every 6 months. In case no improvement is reached 6 months after treatment initiation, treatment with pentosan polysulfate sodium should be stopped. In responders pentosan polysulfate sodium treatment should be continued chronically as long as the response is maintained.

Special populations

Pentosan polysulfate sodium has not been specifically studied in special patient populations like elderly or patients with renal or hepatic impairment (see section 4.4). No dose adjustment is recommended for these patients.

Paediatric population

The safety and efficacy of pentosan polysulfate sodium in children and adolescent below 18 years has not been established.
No data are available.

Method of administration

The capsules should be taken with water at least 1 hour before meals or 2 hours after meals.

4.3 Contraindications

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

Due to the weak anticoagulant effect of pentosan polysulfate sodium, elmiron must not be used in patients who actively bleed. Menstruation is no contraindication.

4.4 Special warnings and precautions for use

Bladder pain syndrome is a diagnosis of exclusion and other urologic disorders should be eliminated by the prescriber, such as urinary tract infection or bladder cancer.

Pentosan polysulfate sodium is a weak anticoagulant. Patients undergoing invasive procedures or having signs/symptoms of underlying coagulopathy or other increased risk of bleeding (due to treatment with other medicinal products influencing coagulation such as anticoagulants, heparin derivatives, thrombolytic or antiplatelet agents including acetylsalicylic acid, or nonsteroidal anti-inflammatory medicinal products (see section 4.5)) should be evaluated for haemorrhagic events. Patients who have a history of heparin or pentosan polysulfate sodium induced thrombocytopenia should be carefully monitored when treated with pentosan polysulfate sodium.

Hepatic or renal insufficiency

elmiron has not been studied in patients with hepatic or renal insufficiency. Because there is evidence of hepatic and renal contribution to the elimination of pentosan polysulfate sodium, hepatic or renal impairment may have an impact on the pharmacokinetics of pentosan polysulfate sodium. Patients with relevant hepatic or renal insufficiency should be carefully monitored when treated with pentosan polysulfate sodium.

Rare cases of pigmentary maculopathy have been reported with use of pentosan polysulfate sodium (PPS), especially after long term use. Visual symptoms might include complaints of reading difficulty and slow adjustment to low or reduced light environments. All patients should have regular ophthalmic examinations for early detection of pigmentary maculopathy, particularly those with long-term use of PPS. In such situations, treatment cessation should be considered.

4.5 Interaction with other medicinal products and other forms of interaction

A study in healthy subjects revealed no pharmacokinetic or pharmacodynamic interactions between therapeutic doses of warfarin and pentosan polysulfate sodium. No further interaction studies have been performed.

Due to the weak anticoagulant effect of pentosan polysulfate sodium, patients, who are concomitantly treated with anticoagulants, heparin derivatives, thrombolytic or antiplatelet agents including acetylsalicylic acid, or nonsteroidal anti-inflammatory medicinal products should be evaluated for any haemorrhagic event in order to adapt the dose if needed (see section 4.4).

4.6 Fertility, pregnancy and lactation

Pregnancy

There are no data from the use of pentosan polysulfate sodium in pregnant women. Animal studies with respect to reproductive toxicity were not conducted.

elmiron is not recommended during pregnancy.

Breast-feeding

It is unknown whether pentosane polysulfate sodium or metabolites are excreted in human milk.

A risk to the newborns/infants cannot be excluded.

Therefore, pentosan polysulfate sodium should not be used during breast-feeding.

Fertility

No information on a potential impact of pentosane polysulfate sodium on fertility is available.

4.7 Effects on ability to drive and use machines

Pentosan polysulfate sodium has no or negligible influence on the ability to drive and use machines.

4.8 Undesirable effects

Summary of the safety profile

The following section lists adverse events reported in the literature from clinical studies with pentosan polysulfate sodium. The potential relatedness between these adverse events and the treatment with pentosan polysulfate sodium was not discussed in the respective publications.

The most common adverse events reported from the clinical studies are headache, dizziness and gastro-intestinal events like diarrhoea, nausea, abdominal pain and rectal bleeding.

The adverse events reported under treatment with pentosan polysulfate sodium were comparable to those reported under treatment with placebo in regards to quality and quantity.

Tabulated summary of adverse events

Adverse events are listed below by MedDRA body system organ class and by frequency. Very common ($\geq 1/10$); common ($\geq 1/100$ to $< 1/10$); uncommon ($\geq 1/1,000$ to $< 1/100$); rare ($\geq 1/10,000$ to $< 1/1,000$); very rare ($< 1/10,000$); not known (cannot be estimated from available data).

Infections and infestations	Common	Infections, influenza
Blood and lymphatic system disorders	Uncommon	Anaemia, ecchymosis, haemorrhage, leukopenia, thrombocytopenia
	Not known	Coagulation disorders
Immune system disorder	Uncommon	Photosensitivity
	Not known	Allergic reactions
Metabolism and nutrition disorders	Uncommon	Anorexia, weight gain, weight loss
Psychiatric disorders	Uncommon	Severe Emotional Lability/Depression
Nervous system disorders	Common	Headache, dizziness
	Uncommon	Increased sweating, insomnia, hyperkinesia, paraesthesia
Eye disorders	Uncommon	Lacrimation, amblyopia
Ear disorders	Uncommon	Tinnitus
Respiratory, thoracic and mediastinal disorders	Uncommon	Dyspnoea
Gastrointestinal disorders	Common	Nausea, diarrhoea, dyspepsia, abdominal pain, abdomen enlarged, rectal haemorrhage
	Uncommon	Indigestion, vomiting, mouth ulcer, flatulence, constipation
Skin and subcutaneous tissue disorders	Common	Peripheral oedema, alopecia
	Uncommon	Rash, increased mole size
Musculoskeletal and connective tissue disorders	Common	Back pain
	Uncommon	Myalgia, Arthralgia
Renal and urinary disorders	Common	Urinary frequency
General disorders and administration site conditions	Common	Asthenia, pelvic pain
Investigation	Not known	Liver function abnormalities

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system listed in [Appendix V](#).

4.9 Overdose

In the case of an accidental overdose, patients should be evaluated for potential adverse effects of pentosan polysulfate sodium like gastrointestinal symptoms or bleeding. In case of adverse reactions, treatment might be paused until the symptoms abate and treatment should be continued at the recommended dose after a critical balancing of the risks thereafter.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Urologicals, other urologicals, ATC code: G04BX15.

Mechanism of action

The hypothetical mechanism of action of pentosan polysulfate sodium includes a local effect in the bladder after systemic administration and excretion into the urine by binding of glycosaminoglycans to the deficient mucous of the bladder. This binding of glycosaminoglycans to the bladder mucous reduces bacterial adherence to the inner surface of the bladder and in consequence the incidence of infections is reduced as well. It is hypothesized, that a potential barrier function of pentosan polysulfate sodium instead of the damaged urothelial mucus might play a role as well the anti-inflammatory activity of pentosan polysulfate sodium.

Clinical efficacy and safety

A total of four randomised placebo-controlled, double-blind clinical studies prospectively enrolling patients with bladder pain syndrome diagnosed via cystoscopic examination with or without bladder hydrodistension evaluating the efficacy of oral treatment with pentosan polysulfate sodium were published in scientific literature. In all of these studies, patients reported a better subjective improvement of bladder pain syndrome under treatment with pentosan polysulfate sodium compared to placebo. In three studies, the observed difference was clearly statistically significant. The first study was a double-blind, randomized, placebo-controlled study with a planned cross-over design evaluating pentosan polysulfate sodium versus placebo. Depending on which institution the patients attended they were treated with either 3x100 mg or 2x200 mg PPS per day. 75 patients were randomised into the study and 62 of those completed the study. Efficacy of treatment was evaluated based on the patient reported improvement on four typical symptoms of bladder pain syndrome: pain, urgency, frequency, and nocturia, no primary endpoint was defined. A patient was counted as a responder to treatment in case a 50% improvement compared to baseline was reported for a specific symptom after 3 months of treatment. An evaluation of all data generated in the study showed that for all four symptoms statistically significant more patients responded to pentosan polysulfate sodium treatment compared to placebo:

	PPS	Placebo	P-value
Pain			
No. responders / total (%)	19/42 (45)	7/38 (18)	0.02
Av. % improvement*	33.0 ± 35	15.8 ± 26	0.01
Urgency			
No. responders / total (%)	21/42 (50)	9/48 (19)	0.03
Av. % improvement*	27.6 ± 31	14.0 ± 24	0.01
Frequency			
No. responders / total (%)	33/52 (63)	16/41 (39)	0.005
Av. improvement	-5.1	-0.4	0.002
Nocturia			

Av. improvement*	-1.5 ± 2.9	-0.5 ± 0.5	0.04
------------------	------------	------------	------

(*Mean ± SD)

The following two studies were conducted following very comparable double-blind, randomized, placebo-controlled multicentre study designs. The patients in both studies were treated for three months with either 3x100 mg pentosan polysulfate sodium or placebo. The primary efficacy endpoint of the study was the overall improvement as self-reported by the patient after three months of treatment. The patients were asked whether they felt improved overall since the start of treatment, and if so, whether the improvement was “slight” 25%, “moderate” 50%, “great” 75% or “complete cure” 100%. Patients who reported at least moderate (50%) improvement were counted as responders. The secondary efficacy endpoints included the investigators evaluation of improvement. The used scale for the investigators assessment included the categories “worse”, “no change”, “fair”, “good”, “very good”, and “excellent”. A responder was defined as a patient assessed to be at least “good” compared to baseline. Furthermore volume voiding profiles over three days and the impact of treatment on pain and urgency were evaluated as secondary endpoints. The impact on pain and urgency was evaluated via the same questionnaire as the primary endpoint with a responder defined as a patient experiencing an at least moderate (50%) improvement compared to baseline. In addition the impact on pain and urgency was evaluated via a 5 score scale, where a responder was defined as a patient experiencing at least a 1-point improvement compared to baseline.

110 patients were enrolled and treated for three months in the first of the two very comparable studies. A statistically significant benefit of pentosan polysulfate sodium over placebo was demonstrated over the primary endpoint, the patients overall-assessment of improvement as well as on the investigators overall assessment. Furthermore a trend for better efficacy of pentosane polysulfate sodium was observed for the patients self-assessment of an improvement of pain and urgency, despite a deviating effect observed for the evaluation of urgency via the scale. In addition positive effects were observed on the voiding profile, although the observed differences were not statistically significant:

	PPS	Placebo	P-value
Responders based on patients self-evaluation of overall improvement	28%	13%	0.04
Responders based on investigators evaluation of overall improvement	26%	11%	0.03
Responders regarding pain and urgency			
Pain (moderate/50% improvement)	27%	14%	0.08
Pain scale (1-point improvement)	46%	29%	0.07
Pressure to urinate (moderate/50% improvement)	22%	11%	0.08
Urgency scale (1-point improvement)	39%	46%	ns
Mean reduction in pain score from baseline	0.5	0.2	ns
Changes from baseline voiding characteristics			
Mean volume per void (cc)	9.8	7.6	ns
Increase of ≥ 20 cc (% pts)	30	20	ns
Total daily urine volume (cc)	+60	-20	ns
Voids per day	-1	-1	ns
3 voids less per day (% pts)	32	24	ns
Nocturia	-0.8	-0.5	ns

The second of the two very comparable studies enrolled 148 patients and demonstrated a statistically significant benefit pentosan polysulfate sodium over placebo was demonstrated on the patient reported overall improvement evaluated as primary endpoint and an the investigator-assessed overall improvement, all evaluations on pain and urgency. A trend for better efficacy under pentosan polysulfate sodium was observed for improved sexual intercourse:

	PPS	Placebo	P-value
Responders based on patients self-evaluation of overall improvement	32%	16%	0.01
Responders based on investigators evaluation of overall improvement	36%	15%	0.002
Responders regarding pain and urgency			
Pain (moderate/50% improvement)	38%	18%	0.005
Pain scale (1-point improvement)	66%	51%	0.04
Pressure to urinate (moderate/50% improvement)	30%	18%	0.04
Responders regarding pain and urgency	61%	43%	0.01
Improved sexual intercourse	31%	18%	0.06
Changes from baseline voided volume			
Mean volume per void (cc)	+20.4	-2.1	ns
Increase of ≥ 20 cc (% pts)	40	24	0.02
Total daily urine volume (cc)	+3	-42	ns

The fourth study was following a double-blind, double-dummy, multifactorial design and evaluated the effects of pentosan polysulfate sodium and hydroxyzine in one study. Patients were randomized to four treatment group and were treated for six months with 3x100 mg pentosane polysulfate sodium, 1x50 mg hydroxyzine, both active treatments or placebo. A responder analysis based on a patient-reported Global Response Assessment (GRA) after 24 weeks of treatment was defined as primary endpoint. The GRA assessment was evaluated via a 7-point centred scale, in which the patients can assess their global response compared to baseline as markedly worse, moderately worse, slightly worse, no change, slightly improved, moderately improved or markedly improved. Participants who reported either of the latter two categories were defined as treatment responders. Secondary outcome measures included the O'Leary-Sant IC Symptom and Problem Index, the University of Wisconsin Symptom score, patient reported symptoms of pain/discomfort and urgency, and results of a 24-hour voiding diary. Comparison of those patients receiving pentosan polysulfate sodium with those not receiving pentosan polysulfate sodium (irrespective of treatment with oral hydroxyzine) revealed no statistically significant difference between the two group, but a trend for better efficacy was observed for the primary endpoint in those patients treated with pentosan polysulfate sodium (either alone or in combination with hydroxyzine) (20 of 59, 34%) compared to the those patients not receiving pentosan polysulfate sodium, but who might receive hydroxyzine (11 of 62, 18%, p0.064):

	PPS	Placebo
No. randomized	59	62
No. responders (%)	20 (34)	11 (18)
No. complete secondary end point data (%)	49 (83)	47 (76)
Mean pain score \pm SD (0-9)	-1.2 \pm 1.9	-0.7 \pm 1.8
Mean urgency score \pm SD (0-9)	-1.2 \pm 1.6	-0.9 \pm 1.6
Mean 24-hr frequency \pm SD	-0.7 \pm 4.8	-0.9 \pm 6.3
Mean IC symptom index \pm SD (0-20)	-2.6 \pm 3.4	-1.7 \pm 3.5
Mean IC problem index \pm SD (0-16)	-2.6 \pm 3.5	-1.9 \pm 2.8
Mean Wisconsin IC score \pm SD (0-42)	-6.2 \pm 8.9	-6.7 \pm 8.2

A pooled analysis of the data described above from placebo-controlled clinical studies was conducted to evaluate, whether patients taking oral pentosane polysulfate sodium have clear benefit from the treatment. This pooled analysis showed that the percentage of patients responding to treatment with pentosan polysulfate sodium with a clinically relevant improvement in their overall assessment, pain and urgency was approximately 2-fold higher than the respective responder rates under placebo:

	PPS	Placebo
GRA (95% CI)	33.0% (27.1% - 39.4%)	15.8% (11.6% - 21.2%)
Pain (95% CI)	32.7% (26.0% - 40.3%)	14.2% (9.6% - 20.6%)
Urgency (95% CI)	27.4% (21.1% - 34.8%)	14.2% (9.6% - 20.6%)

5.2 Pharmacokinetic properties

Absorption

Less than 10% of orally administered pentosan polysulfate sodium are slowly absorbed from the gastrointestinal tract and are available in systemic circulation in the form of unchanged pentosan polysulfate sodium or its metabolites. All studies describe very low systemic availability of unchanged pentosan polysulfate sodium after oral administration. Overall, the reported systemic bioavailability after oral administration of pentosan polysulfate sodium is below 1%.

Distribution

In healthy volunteers, a single parenteral administration of radioactively labelled pentosan polysulfate sodium leads to a progressive up-take of total radioactivity by the liver, spleen, and kidney (50 min after 1 mg/kg i.v.: 60 % of the dose in the liver, 7.7 % in the spleen; 3 h post dosing: 60 % in the liver plus spleen, and 13 % in the bladder).

Biotransformation

Pentosan polysulfate sodium is metabolised extensively by desulfation in liver and spleen and depolymerisation in the kidney.

Elimination

The apparent plasma half-life of pentosan polysulfate sodium depends on the route of administration. While pentosan polysulfate sodium is rapidly cleared from circulation of i.v. administration, the apparent plasma half-life after oral administration is in the range of 24-34 hours. Accordingly, oral administration of pentosan polysulfate sodium 3-times daily is expected to lead to accumulation of pentosan polysulfate sodium over the first 7 days of administration (accumulation factor 5-6.7). After oral administration unabsorbed pentosan polysulfate sodium is excreted predominantly unchanged in the faeces. About 6% of the administered dose of pentosan polysulfate sodium were excreted via urine after desulfation and depolymerisation.

5.3 Preclinical safety data

Non-clinical data reveal no special hazard for humans based on conventional repeated dose toxicity, genotoxicity and long-term carcinogenicity studies.

The effect of pentosan polysulfate sodium on reproductive and developmental toxicity has not been investigated.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Capsule content

Microcrystalline cellulose
Magnesium stearate

Capsule shell

Gelatin
Titanium dioxide (E171)

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

Bottle

3 years

After first opening: use within 45 days.

Blister

21 months

6.4 Special precautions for storage

Bottle

Keep the bottle tightly closed in order to protect from moisture.

For storage conditions after first opening of the bottle, see section 6.3.

Blister

Do not store above 30°C.

6.5 Nature and contents of container

HDPE bottle with a tamper-evident child resistant closure of PP with 90 capsules.

HDPE bottle with a tamper-evident child resistant closure of PP with 100 capsules.

PVC/Aclar-Aluminium blister with 90 (9x10) capsules.

Bottle

Pack size of 90 capsules.

Pack size of 300 (3 bottles x 100) capsules.

Blister

Pack size of 90 capsules.

Not all pack sizes may be marketed.

← somewhere
else in
doc

6.6 Special precautions for disposal

No special requirements.

7. MARKETING AUTHORISATION HOLDER

bene-Arzneimittel GmbH

Herterichstrasse 1-3

D-81479 Munich

tel: ++49 (0) 89 / 7 49 87-0

fax: ++49 (0) 89 / 7 49 87-142

e-mail: contact@bene-arzneimittel.de

8. MARKETING AUTHORISATION NUMBER(S)

EU/1/17/1189/001

EU/1/17/1189/002

9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of first authorisation: 2 June 2017

10. DATE OF REVISION OF THE TEXT

Detailed information on this medicinal product is available on the website of the European Medicines Agency <http://www.ema.europa.eu>

ANNEX II

- A. MANUFACTURER(S) RESPONSIBLE FOR BATCH RELEASE**
- B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**
- C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**
- D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

A. MANUFACTURER RESPONSIBLE FOR BATCH RELEASE

Name and address of the manufacturer responsible for batch release

bene-Arzneimittel GmbH
Herterichstr. 1 - 3
81479 Munich
GERMANY

B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE

Medicinal product subject to medical prescription.

C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION

- **Periodic Safety Update Reports**

The requirements for submission of periodic safety update reports for this medicinal product are set out in the list of Union reference dates (EURD list) provided for under Article 107c(7) of Directive 2001/83/EC and any subsequent updates published on the European medicines web-portal.

The marketing authorisation holder shall submit the first periodic safety update report for this product within 6 months following authorisation.

D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT

- **Risk Management Plan (RMP)**

The MAH shall perform the required pharmacovigilance activities and interventions detailed in the agreed RMP presented in Module 1.8.2 of the Marketing Authorisation and any agreed subsequent updates of the RMP.

An updated RMP should be submitted:

- At the request of the European Medicines Agency;
- Whenever the risk management system is modified, especially as the result of new information being received that may lead to a significant change to the benefit/risk profile or as the result of an important (pharmacovigilance or risk minimisation) milestone being reached.

ANNEX III
LABELLING AND PACKAGE LEAFLET

A. LABELLING

PARTICULARS TO APPEAR ON THE OUTER PACKAGING

CARTON FOR BOTTLE

1. NAME OF THE MEDICINAL PRODUCT

elmiron 100 mg hard capsules
pentosan polysulfate sodium

2. STATEMENT OF ACTIVE SUBSTANCE(S)

Each capsule contains 100 mg of pentosan polysulfate sodium.

3. LIST OF EXCIPIENTS

4. PHARMACEUTICAL FORM AND CONTENTS

90 hard capsules
3 bottles
300 (3x100) hard capsules

5. METHOD AND ROUTE(S) OF ADMINISTRATION

Read the package leaflet before use.

Oral use

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY

8. EXPIRY DATE

EXP
After first opening: use within 45 days.

9. SPECIAL STORAGE CONDITIONS

Keep the bottle tightly closed in order to protect from moisture.

PARTICULARS TO APPEAR ON THE OUTER PACKAGING

CARTON FOR BLISTERS

1. NAME OF THE MEDICINAL PRODUCT

elmiron 100 mg hard capsules
pentosan polysulfate sodium

2. STATEMENT OF ACTIVE SUBSTANCE(S)

Each capsule contains 100 mg of pentosan polysulfate sodium.

3. LIST OF EXCIPIENTS

4. PHARMACEUTICAL FORM AND CONTENTS

90 hard capsules

5. METHOD AND ROUTE(S) OF ADMINISTRATION

Read the package leaflet before use.

Oral use

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY

8. EXPIRY DATE

EXP

9. SPECIAL STORAGE CONDITIONS

Do not store above 30°C.

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

bene-Arzneimittel GmbH, PO Box 710269, 81452 Munich, Germany

12. MARKETING AUTHORISATION NUMBER(S)

EU/1/17/1189/002

13. BATCH NUMBER

Lot

14. GENERAL CLASSIFICATION FOR SUPPLY

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

• • • • •

17. UNIQUE IDENTIFIER – 2D BARCODE

2D barcode carrying the unique identifier included.

18. UNIQUE IDENTIFIER - HUMAN READABLE DATA

PC: {number}

SN: {number}

NN: {number}

PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGING

BOTTLE LABEL

1. NAME OF THE MEDICINAL PRODUCT

elmiron 100 mg hard capsules
pentosan polysulfate sodium

2. STATEMENT OF ACTIVE SUBSTANCE(S)

Each capsule contains 100 mg of pentosan polysulfate sodium.

3. LIST OF EXCIPIENTS

4. PHARMACEUTICAL FORM AND CONTENTS

90 hard capsules
100 hard capsules

5. METHOD AND ROUTE(S) OF ADMINISTRATION

Read the package leaflet before use.

Oral use.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY

8. EXPIRY DATE

EXP

After first opening: use within 45 days.

Open date:

9. SPECIAL STORAGE CONDITIONS

Keep the bottle tightly closed in order to protect from moisture.

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

bene-Arzneimittel GmbH, PO Box 710269, 81452 Munich, Germany

12. MARKETING AUTHORISATION NUMBER(S)

EU/1/17/1189/001 90 hard capsules

EU/1/17/1189/003 300 (3 bottles x 100) hard capsules

13. BATCH NUMBER<, DONATION AND PRODUCT CODES>

Lot

14. GENERAL CLASSIFICATION FOR SUPPLY

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS
--

BLISTER

1. NAME OF THE MEDICINAL PRODUCT

elmiron 100 mg hard capsules
pentosan polysulfate sodium

2. NAME OF THE MARKETING AUTHORISATION HOLDER
--

bene-Arzneimittel GmbH

3. EXPIRY DATE

EXP

4. BATCH NUMBER<, DONATION AND PRODUCT CODES>
--

Lot

5. OTHER

B. PACKAGE LEAFLET

Package leaflet: Information for the patient

elmiron 100 mg hard capsules pentosan polysulfate sodium

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What elmiron is and what it is used for
2. What you need to know before you take elmiron
3. How to take elmiron
4. Possible side effects
5. How to store elmiron
6. Contents of the pack and other information

1. What elmiron is and what it is used for

elmiron is a medicine that contains the active substance pentosan polysulfate sodium. After taking the medicine it passes into the urine and attaches to the lining of the bladder, helping to form a protective layer.

elmiron is used in adults to treat **bladder pain syndrome** characterised by many tiny bleeds or distinctive lesions on the bladder wall and moderate to severe pain and a frequent urge to urinate.

2. What you need to know before you take elmiron

Do not take elmiron if you are

- **allergic** to pentosan polysulfate sodium or any of the other ingredients of this medicine (listed in section 6)
- **bleeding** (other than menstrual bleeding)

Warnings and precautions

Talk to your doctor or pharmacist before taking elmiron if you have:

- to undergo surgery
- a blood clotting disorder or increased risks of bleeding, such as using a medicine that inhibits blood clotting
- ever had a reduced number of blood platelets caused by the medicine called heparin
- reduced liver or kidney function

Rare cases of retinal disorders (pigmentary maculopathy) have been reported with use of elmiron (especially after long term use). Tell your doctor immediately if you experience visual changes such as reading difficulty and slower adjustment to low or reduced light. Your doctor will discuss with you whether the treatment should be continued. For early detection of retinal disorders, eye examination will be performed regularly.

Children and adolescents

elmiron **is not recommended** in children under 18 years as safety and efficacy have not been established in this group.

Other medicines and elmiron

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

Inform your doctor or pharmacist, particularly if you use medicines that prevent blood clotting, or painkillers that reduce blood clotting.

Pregnancy and breast-feeding

elmiron is **not recommended** during pregnancy or breast-feeding.

Driving and using machines

elmiron has no or negligible influence on the ability to drive and use machines.

3. How to take elmiron

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

The recommended dose is:

1 capsule, 3 times daily

Your doctor will assess your response to elmiron every 6 months.

Method of use

Take the capsules whole with one glass of water, at least 1 hour before or 2 hours after meals.

If you take more elmiron than you should

Inform your doctor, in the case of an overdose. Stop taking elmiron if side effects occur, until they disappear.

If you forget to take elmiron

Do not take a double dose to make up for a forgotten capsule.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Side effects have been observed with the following frequencies:

Common: may affect up to 1 in 10 people

- infections, flu

- headache, back pain
- dizziness
- nausea, indigestion, diarrhoea, abdominal pain, abdomen enlarged
- rectal bleeding
- accumulation of fluid in arms or legs
- hair loss
- weakness, pelvic (lower abdomen) pain
- need to urinate more frequently than usual
- abnormal liver function

Uncommon: may affect up to 1 in 100 people

- lack of blood platelets, red or white blood cells
- bleeding, including small bleeding beneath the skin
- blood clotting disorders
- allergic reactions, increased sensitivity to light
- loss of appetite, weight gain or loss
- severe mood swings or depression
- increased sweating, sleeplessness
- restlessness
- abnormal sensation such as prickling, tingling and itchiness
- flow of tears, lazy eye
- ringing or buzzing in the ears
- breathing difficulties
- indigestion, vomiting, wind, difficulty passing stools
- mouth ulcer
- skin rash, increased mole size
- joint or muscle pain

Not known: frequency cannot be estimated from the available data

- blood clotting disorders
- allergic reactions
- abnormal liver function

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via [the national reporting system](#) listed in [Appendix V](#). By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store elmiron

Keep this medicine out of the sight and reach of children.

• **bottle**

Do not use this medicine after the expiry date which is stated on the label and carton after EXP. The expiry date refers to the last day of that month.

Keep the bottle tightly closed in order to protect from moisture.

After first opening: use within 45 days. Dispose any remaining capsules after this period.

• **blister**

Do not use this medicine after the expiry date which is stated on the carton and blister after EXP. The expiry date refers to the last day of that month.

Do not store above 30 C.

Do not throw away any medicines via wastewater. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What elmiron contains

- The active substance is pentosan polysulfate sodium.
One hard capsule contains 100 mg pentosan polysulfate sodium.
- The other ingredients are microcrystalline cellulose, magnesium stearate, gelatin, titanium dioxide (E171).

What elmiron looks like and contents of the pack

The hard capsules are white and non-transparent, provided in a plastic bottle with child resistant closure or plastic/aluminium blisters, packed in a carton.

- **bottle**

Each carton contains 90 capsules.

Each carton contains 300 (3 bottles x 100) capsules.

- **blister**

Each carton contains 90 capsules.

Not all pack sizes may be marketed.

Marketing Authorisation Holder and Manufacturer

bene-Arzneimittel GmbH
Herterichstrasse 1-3
D-81479 Munich
tel: +49 (0)89 749870
fax: +49 (0)89 74987142
e-mail: contact@bene-arzneimittel.de

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder:

BG, CZ, EE, EL, IS, IT, CY, LV, LT, HU, MT, PL, PT, RO, SK

bene-Arzneimittel GmbH, D-81479 Munich,

Германия / Německo / Saksamaa / Γερμανία / Germany / Þýskaland / Germania / Vācija / Vokietija /
Németország / Il-Ġermanja / Niemcy / Alemanha / Nemecko,

Tel / Τел. / Τηλ / Sími / Tel.: +49 (0)89 749870, contact@bene-arzneimittel.de

AT

SIGMAPHARM Arzneimittel GmbH, Leystraße 129, A-1200 Wien,
Österreich,

Tel.: +43 (0) 1 330 06 71-0, mail@sigmapharm.at

BE, LU, NL

Lamepro B.V., Burgemeester Guljélaan 2, NL-4837 CZ Breda,
Pays-Bas, Nederland, Niederlande,

Tél/Tel: +31 (0)76 5600030, lamepro@lamepro.nl

DE

Dr. Pflieger Arzneimittel GmbH, D-96045 Bamberg,
Deutschland,
Tel.: +49 (0)951 6043-0, info@dr-pfleger.de

DK, NO

Navamedic AB, Krokslätts Parkgata 4, PO Box 24032, S-400 22 Göteborg,
Sverige,
Tlf: +46 (0)31 3351190, infose@navamedic.com

ES

Lacer S.A., Sardenya 350, 08025 Barcelona,
España,
Tel: +34 (0)934465300, info@lacer.es

FI, SE

Navamedic AB, Krokslätts Parkgata 4, PO Box 24032, S-400 22 Göteborg,
Sverige,
Puh/Tel: +46 (0)31 3351190, infose@navamedic.com

FR

Inresa SAS, 1 rue Jean Monnet, F-68870 Bartenheim,
France,
Tél: +33 (0)389 707660, info@inresa.fr

HR

MEDICOPHARMACIA d.o.o., Pere Budmanija 5, 10000 Zagreb,
Hrvatska,
Tel: + 385 1 55 84 604

IE

Consilient Health, Block 2A Richview Office Park, Clonskeagh, Dublin 14, D14 Y0A5
Ireland,
Tel: +353 (0) 1 2057760, irishoffice@consilienthealth.com

SI

Lenis d.o.o., Litostrojska cesta 52, 1000 Ljubljana,
Slovenija,
Tel: +386(0) 1 235 07 00, info@lenis.si

UK

Consilient Health Ltd., 1 Church Road, Richmond upon Thames, Surrey, TW9 2QE
United Kingdom,
Tel: +44 (0)20 3751 1888, info@consilienthealth.com

This leaflet was last revised in <{MM/YYYY}>.

Detailed information on this medicine is available on the European Medicines Agency web site
<http://www.ema.europa.eu>.