
BRAZIL

Cutaneous and mucosal leishmaniasis

2023



12 910 new cases reported
Cutaneous: 12 137 (94%)
Mucosal: 770 (6%)

12.4 cases per 100 000 population

Transmission risk:
VERY INTENSE
(CL composite indicator, 2021–2023 triennium)

***LEISHMANIA* SPECIES:**
L. amazonensis, L. braziliensis, L. guyanensis, L. lainsoni, L. lindenbergi, L. naiffi, and *L. shawi*

VECTORS - *LUTZOMYIA*:
Lu. amazonensis, Lu. ayrozai, Lu. complexa, Lu. davisi, Lu. fischeri, Lu. flaviscutellata, Lu. intermedia, Lu. migonei, Lu. neivai, Lu. ovallesi, Lu. paraensis, Lu. pessoai, Lu. salesi, Lu. shawi, Lu. ubiquitalis, Lu. umbratillis y *Lu. wellcomei,* and *Lu. whitmani*

AVAILABLE MEDICATIONS:
meglumine antimoniate, liposomal amphotericin B, amphotericin B deoxycholate, pentamidine isethionate, miltefosine, and local treatment

AGE GROUP
5.2% of cases in children under 10 years old

SEX
73% of cases in men

DIAGNOSIS
81% of cases diagnosed by laboratory

EVOLUTION
71% of cured patients
29% with no information on patient evolution

SURVEILLANCE
51% of the population in transmission areas

NOTIFICATION
114% of detected and reported cases, among those expected

Note: The Ministry of Health reported 1153 cases from unidentified municipalities that were not included in the leishmaniasis composite indicator analysis.
CL: cutaneous leishmaniasis

Source: Pan American Health Organization. Regional Information System on Leishmaniasis (SisLeish). Washington, D.C.: PAHO; 2024 [cited 8 October 2024]. Limited access.


Pan American Health Organization
World Health Organization
Americas Region

PAHO

BRAZIL

Visceral leishmaniasis

2023



1461 new cases reported

2.7 cases per 100 000 population

Transmission risk:
VERY INTENSE
(Regional VL composite indicator, triennium 2021–2023)

***LEISHMANIA* SPECIES:**
L. infantum

VECTORS - *LUTZOMYIA*:
Lu. cruzi, Lu. longipalpis, and *Lu. migonei*

DOMESTIC RESERVOIR: dog

FIRST-LINE MEDICATIONS:
meglumine antimoniate and liposomal amphotericin B

AGE GROUP
18% of cases in children under 5 years old and 29% in people over 50 years old

SEX
68% of cases in men

DIAGNOSIS
87% of cases diagnosed by laboratory

VL–HIV coinfection
20% of VL–HIV coinfected patients
11% with **no information** on HIV status

EVOLUTION
65% of cured patients
8% fatality by VL
23% **with no information** on patient evolution

BORDERS
3% of cases at international borders

Note: The Ministry of Health reported three cases from unidentified municipalities that were not included in the leishmaniasis composite indicator analysis.
VL: visceral leishmaniasis

Source: Pan American Health Organization. Regional Information System on Leishmaniasis (SisLeish). Washington, D.C.: PAHO; 2024 [cited 8 October 2024]. Limited access.

Pan American Health Organization
World Health Organization
Americas Region
PAHO