

Draft Genome Sequence of “*Candidatus Halobonum tyrrellensis*” Strain G22, Isolated from the Hypersaline Waters of Lake Tyrrell, Australia

Juan A. Ugalde,^a Priya Narasingarao,^a Sidney Kuo,^b Sheila Podell,^a Eric E. Allen^{a,b}

Marine Biology Research Division, Scripps Institution of Oceanography, University of California, San Diego, La Jolla, California, USA^a; Division of Biological Sciences, University of California, San Diego, La Jolla, California, USA^b

We report the draft 3.675-Mbp genome sequence of “*Candidatus Halobonum tyrrellensis*” strain G22, a novel halophilic archaeon isolated from the surface hypersaline waters of Lake Tyrrell, Australia. The availability of the first genome from the “*Candidatus Halobonum*” genus provides a new genomic resource for the comparative genomic analysis of halophilic *Archaea*.

Received 25 October 2013 Accepted 11 November 2013 Published 12 December 2013

Citation Ugalde JA, Narasingarao P, Kuo S, Podell S, Allen EE. 2013. Draft genome sequence of “*Candidatus Halobonum tyrrellensis*” strain G22, isolated from the hypersaline waters of Lake Tyrrell, Australia. *Genome Announc.* 1(6):e01001-13. doi:10.1128/genomeA.01001-13.

Copyright © 2013 Ugalde et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](http://creativecommons.org/licenses/by/3.0/).

Address correspondence to Eric E. Allen, eallen@ucsd.edu.

Halophilic *Archaea* of the class *Halobacteria* (phylum *Euryarchaeota*) are dominant members of extreme hypersaline environments worldwide (1). Numerous genera have been isolated from diverse hypersaline habitats, and many representative genome sequences are available (1, 2). However, recent metagenomic analyses of hypersaline ecosystems have revealed that these reference halophiles are not adequately representative of the dominant microbial populations present in many natural hypersaline habitats (3–5). Here, we report the genome sequence of a novel member of the class *Halobacteria* isolated from the hypersaline surface waters of Lake Tyrrell, Victoria, Australia.

Surface water samples were plated onto minimal medium containing a 23% salt solution amended with various carbon substrates, including glycerol, acetate, or glucose, and incubated at room temperature under aerobic conditions. After 3 to 4 weeks of incubation, the colonies were restreaked for purity and characterized via 16S rRNA gene amplification and sequencing to screen for novel species/strains. “*Candidatus Halobonum tyrrellensis*” strain G22 was isolated from minimal medium containing glycerol as the sole carbon source, incubated aerobically at room temperature. Genomic DNA was sequenced using 454 Titanium chemistry at the J. Craig Venter Institute (Rockville, MD). The total number of reads generated was 568,949, with an average length of 428 bp. The sequences were assembled using Newbler (version 2.7), resulting in a total of 72 contigs (N_{50} , 119,067 bp; mean contig length, 45,962 bp; maximum contig length, 303,316 bp), with an estimated genome size of 3,675,087 bp and a G+C content of 70.1%. Functional annotation of predicted gene sequences was performed using the IMG-ER platform (6). A total of 3,525 predicted coding sequences were identified, including 47 tRNAs and a single copy of the rRNA operon.

A phylogenetic tree based on 16S rRNA genes (<http://dx.doi.org/10.6084/m9.figshare.830514>) suggests that “*Ca. Halobonum tyrrellensis*” is a member of the *Halobacteriaceae* family and a sister group of the *Halobaculum* genus, sharing 92% 16S rRNA gene sequence identity with *Halobaculum gomorrense* (7). A com-

parison of the “*Ca. Halobonum tyrrellensis*” genome with the partial genome sequence available for *H. gomorrense* (632,433 bp) (8) revealed an average nucleotide identity (ANI) of $81.56 \pm 1.18\%$. A phylogenomic approach using multiple amino acid markers (9) supports the placement of “*Ca. Halobonum tyrrellensis*” as a new genus (<http://dx.doi.org/10.6084/m9.figshare.830514>). A detailed characterization of the physiology and metabolism of “*Ca. Halobonum tyrrellensis*” and a formal description of this strain are currently in progress.

The features found in the genome include the presence of a putative sensory rhodopsin, a high number of ABC transporters and carbon metabolism genes, including trehalose utilization genes, and the absence of conserved haloarchaeal genes encoding a flagellar system or gas vesicle synthesis proteins.

The “*Ca. Halobonum tyrrellensis*” genome represents the first high-quality draft sequence for a member of the new candidate genus *Halobonum*. These data expand the breadth of the reference genome sequence information for halophilic *Archaea*, providing a new resource for comparative genomic analyses and the phylogenetic binning of metagenomic sequence data recovered from hypersaline environments.

Nucleotide sequence accession number. The draft genome sequence of “*Ca. Halobonum tyrrellensis*” strain G22 is deposited at DDBJ/EMBL/Genbank databases under the accession no. [ASGZ000000000](https://www.ncbi.nlm.nih.gov/nuclseq/ASGZ000000000).

ACKNOWLEDGMENTS

Funding for this work was provided by NSF award no. 0626526 (to J. F. Banfield, K. B. Heidelberg, and E.E.A.) and NIH award R21HG005107-02 (E.E.A.). J.A.U. was supported by a Fulbright-Conicyt fellowship.

REFERENCES

1. Andrei AS, Banciu HL, Oren A. 2012. Living with salt: metabolic and phylogenetic diversity of archaea inhabiting saline ecosystems. *FEMS Microbiol. Lett.* 330:1–9.
2. Lynch EA, Langille MG, Darling A, Wilbanks EG, Haltiner C, Shao KS, Starr MO, Teiling C, Harkins TT, Edwards RA, Eisen JA, Facciotti MT.

2012. Sequencing of seven Haloarchaeal genomes reveals patterns of genomic flux. PLoS One 7:e41389. doi:[10.1371/journal.pone.0041389](https://doi.org/10.1371/journal.pone.0041389).
3. Ghai R, Pasić L, Fernández AB, Martín-Cuadrado A-B, Mizuno CM, McMahon KD, Papke RT, Stepanauskas R, Rodríguez-Brito B, Rohwer F, Sánchez-Porro C, Ventosa A, Rodríguez-Valera F. 2011. New abundant microbial groups in aquatic hypersaline environments. Sci. Rep. 1:135.
4. Podell S, Ugalde JA, Narasingarao P, Banfield JF, Heidelberg KB, Allen EE. 2013. Assembly-driven community genomics of a hypersaline microbial ecosystem. PLoS One 8:e61692. doi:[10.1371/journal.pone.0061692](https://doi.org/10.1371/journal.pone.0061692).
5. Narasingarao P, Podell S, Ugalde JA, Brochier-Armanet C, Emerson JB, Brocks JJ, Heidelberg KB, Banfield JF, Allen EE. 2012. *De novo* metagenomic assembly reveals abundant novel major lineage of Archaea in hypersaline microbial communities. ISME J. 6:81–93.
6. Markowitz VM, Chen IMA, Palaniappan K, Chu K, Szeto E, Grechkin Y, Ratner A, Jacob B, Huang J, Williams P, Huntemann M, Anderson I, Mavromatis K, Ivanova NN, Kyrpides NC. 2011. IMG: the integrated microbial genomes database and comparative analysis system. Nucleic Acids Res. 40:D115–D122. doi:[10.1093/nar/gkr1044](https://doi.org/10.1093/nar/gkr1044).
7. Oren A, Gurevich P, Gemmell RT, Teske A. 1995. *Halobaculum gomorense* gen. nov., sp. nov., a novel extremely halophilic archaeon from the Dead Sea. Int. J. Syst. Bacteriol. 45:747–754.
8. Goo YA, Roach J, Glusman G, Baliga NS, Deutsch K, Pan M, Kennedy S, DasSarma S, Ng WV, Hood L. 2004. Low-pass sequencing for microbial comparative genomics. BMC Genomics 5:3. doi:[10.1186/1471-2164-5-3](https://doi.org/10.1186/1471-2164-5-3).
9. Segata N, Börnigen D, Morgan XC, Huttenhower C. 2013. PhyloPhlAn is a new method for improved phylogenetic and taxonomic placement of microbes. Nat. Commun. 4:2304. doi:[10.1038/ncomms3304](https://doi.org/10.1038/ncomms3304).