

# FUNDAMENTAL THEOREM OF CALCULUS

JUAN VALLEJO

## PART 1

The *integral* from  $a$  to  $x$  represents the area under the curve  $y = f(t)$ . For a given  $f(t)$ , we define the function  $F(x)$  as

$$F(x) = \int_a^x f(t)dt, \quad \text{where } a \leq x \leq b$$

The above equation does not define or depend on knowledge of *derivatives*. We define the *derivative* of a  $F(x)$  as follows:

$$F'(x) = \lim_{\Delta x \rightarrow 0} \frac{F(x + \Delta x) - F(x)}{\Delta x} = \lim_{\Delta x \rightarrow 0} \frac{\int_a^{x+\Delta x} f(t)dt - \int_a^x f(t)dt}{\Delta x}$$

**Definition 1.** *The derivative of a function  $F(x)$  is equal to the quotient between the area under the curve  $f(t)$  from  $a$  to  $x + \Delta x$  minus the area under the curve  $f(t)$  from  $a$  to  $x$ , and  $\Delta x$ .*

$$\lim_{\Delta x \rightarrow 0} \frac{1}{\Delta x} \int_x^{x+\Delta x} f(t)dt$$

**Definition 2.** *Knowing the above, it can be stated that in the area under the curve  $f(t)$  from  $x$  to  $x + \Delta x$ , there exists a point  $c$  such that  $f(c)\Delta x = \int_x^{x+\Delta x} f(t)dt$ . This function  $f(c)$  is therefore known as the mean value function over the integral.*

$$f(c) = \frac{1}{\Delta x} \int_x^{x+\Delta x} f(t)dt$$

In other words, there exists a value  $c$  in  $[x, x + \Delta x]$  where

$$F'(x) = \lim_{\Delta x \rightarrow 0} f(c)$$

Knowing this, it can then be stated that  $f(c) \rightarrow f(x)$  as  $\Delta x \rightarrow 0$ .

**Theorem 3.** *Let there be two functions  $f_-(x)$  and  $f_+(x)$  such that  $f(x)$  is squeezed between the two,*

$$f_-(x) \leq f(x) \leq f_+(x)$$

*if*

$$r = \lim_{x \rightarrow a} f_-(x) = \lim_{x \rightarrow a} f_+(x)$$

*then,*

$$\lim_{x \rightarrow a} f(x) = r$$

Taking the *squeeze theorem* (defined above) into account, and assuming that

$$x \leq c(\Delta x) \leq x + \Delta x$$

we can then state that if

$$\lim_{\Delta x \rightarrow 0} x = x$$

and

$$\lim_{\Delta x \rightarrow 0} x + \Delta x = x$$

then,

$$\lim_{\Delta x \rightarrow 0} C(\Delta x) = x$$

The function  $f$  is continuous at  $c$ , so the limit can be taken inside of the function. Therefore

$$F'(x) = f(x)$$

Thus, proving that any continuous function has an anti-derivative.

*Q.E.D*

PART 2

SOURCES