

Least squares

Sample Subtitle

Juan V. Vía

August 14, 2021

Example

Showing why least squares

We have a variable y . We know that it's dependent of another variable x in some way. But we don't know how, exactly. So we go to the field and measure certain points. Those that we can reach. Six of them.


$$(2, 5), (5, 5), (7, 8), (11, 7), (14, 9), (18, 7)$$

That is: at $x = 2$ we measure $y = 5$, at $x = 5$ we measure $y = 5$ again, but at $x = 7$ we got $y = 8$, and so on.

Example

Showing why least squares

Back to desk we plot these points.




There is a model lurking in these points? A line perhaps?

Example

Showing why least squares

Let's start by tracing a line with "best fit" that data




Why a line? To warm the modeling machine in our minds we are considering a line, a one degree polynomial.

Example

Showing why least squares

What line to plot? How to choose that line?



Enter the math...

Example

Showing why least squares

A very common way to express a line in the plane is isolating y as a polynomial function of x :

$$y = mx + b$$

From this equation we can see that m is the *slope* of the line and b is the *y-intercept* (the value of y when $x = 0$).

Will be handy for us to write this equation using vector notation:

$$y = \begin{bmatrix} x & 1 \end{bmatrix} \begin{bmatrix} m \\ b \end{bmatrix}$$

And switching terms:

$$\begin{bmatrix} x & 1 \end{bmatrix} \begin{bmatrix} m \\ b \end{bmatrix} = y$$

Example

Showing why least squares

Ok. Fine. This form

$$\begin{bmatrix} x & 1 \end{bmatrix} \begin{bmatrix} m \\ b \end{bmatrix} = y$$

will be a useful one because we know x and y in six points. For example take the first point $(2, 5)$

$$\begin{bmatrix} 2 & 1 \end{bmatrix} \begin{bmatrix} m \\ b \end{bmatrix} = 5$$

Example

Showing why least squares

And yes, your guess is true. We can incorporate the second point (5, 5) and get

$$\begin{bmatrix} 2 & 1 \\ 5 & 1 \end{bmatrix} \begin{bmatrix} m \\ b \end{bmatrix} = \begin{bmatrix} 5 \\ 5 \end{bmatrix}$$

Doing the product you can workout m and b from this equation. $m = 0$ and $b = 5$. Because the matrix is square. And invertible by the way.

Example

Showing why least squares

Going further, incorporate the third point.

$$\begin{bmatrix} 2 & 1 \\ 5 & 1 \\ 7 & 1 \end{bmatrix} \begin{bmatrix} m \\ b \end{bmatrix} = \begin{bmatrix} 5 \\ 5 \\ 8 \end{bmatrix}$$

This invalidate the previous result. m cannot be 0 more. Neither b with 5.

Actually, henceforth the equation is over-determined and have no solution.

Example

Showing why least squares

Fourth point.

$$\begin{bmatrix} 2 & 1 \\ 5 & 1 \\ 7 & 1 \\ 11 & 1 \end{bmatrix} \begin{bmatrix} m \\ b \end{bmatrix} = \begin{bmatrix} 5 \\ 5 \\ 8 \\ 7 \end{bmatrix}$$

Example

Showing why least squares

Fifth point.

$$\begin{bmatrix} 2 & 1 \\ 5 & 1 \\ 7 & 1 \\ 11 & 1 \\ 14 & 1 \end{bmatrix} \begin{bmatrix} m \\ b \end{bmatrix} = \begin{bmatrix} 5 \\ 5 \\ 8 \\ 7 \\ 9 \end{bmatrix}$$

Example

Showing why least squares

Sixth and last point.

$$\begin{bmatrix} 2 & 1 \\ 5 & 1 \\ 7 & 1 \\ 11 & 1 \\ 14 & 1 \\ 18 & 1 \end{bmatrix} \begin{bmatrix} m \\ b \end{bmatrix} = \begin{bmatrix} 5 \\ 5 \\ 8 \\ 7 \\ 9 \\ 7 \end{bmatrix}$$

This is the equation derived from our measurement.

Example

Showing why least squares

Time to name things.

Example

Showing why least squares

Call A to the matrix

$$A = \begin{bmatrix} 2 & 1 \\ 5 & 1 \\ 7 & 1 \\ 11 & 1 \\ 14 & 1 \\ 18 & 1 \end{bmatrix}$$

Example

Showing why least squares

Call x to the column vector

$$x = \begin{bmatrix} m \\ b \end{bmatrix}$$

Example

Showing why least squares

Call b to the column vector

$$b = \begin{bmatrix} 5 \\ 5 \\ 8 \\ 7 \\ 9 \\ 7 \end{bmatrix}$$

Example

Showing why least squares

Thus, the equation

$$\begin{bmatrix} 2 & 1 \\ 5 & 1 \\ 7 & 1 \\ 11 & 1 \\ 14 & 1 \\ 18 & 1 \end{bmatrix} \begin{bmatrix} m \\ b \end{bmatrix} = \begin{bmatrix} 5 \\ 5 \\ 8 \\ 7 \\ 9 \\ 7 \end{bmatrix}$$

becomes

$$Ax = b$$

Example

Showing why least squares

In other words: if you want to know the values m and b in the equation $y = mx + b$ of the line we are searching find the x such $Ax = b$.

Example

Showing why least squares

In other words: if you want to know the values m and b in the equation $y = mx + b$ of the line we are searching find the x such $Ax = b$, but **that equation have no solution!**

Example

Showing why least squares

In other words: if you want to know the values m and b in the equation $y = mx + b$ of the line we are searching find the x such $Ax = b$, but **that equation have no solution!** because the matrix A is tall, so $Ax = b$ is over-determined, there are more equations than variables to choose. There is no x such the norm of the difference between Ax and b equals zero

$$\|Ax - b\| \neq 0$$

Example

Showing why least squares

In other words: if you want to know the values m and b in the equation $y = mx + b$ of the line we are searching find the x such $Ax = b$, but **that equation have no solution!** because the matrix A is tall, so $Ax = b$ is over-determined, there are more equations than variables to choose. There is no x such the norm of the difference between Ax and b equals zero

$$\|Ax - b\| \neq 0$$

We need the least squares approximate solution

Example

Showing why least squares

In the least squares approximate solution of $Ax = b$ we, provided that $\|Ax - b\| \neq 0$, select x that the *norm* of the *residual* $r = Ax - b$ is minimal.

Example

Showing why least squares

In the least squares approximate solution of $Ax = b$ we, provided that $\|Ax - b\| \neq 0$, select x that the *norm* of the *residual* $r = Ax - b$ is minimal.

It's all there: <http://vmls-book.stanford.edu/> and in many other sources. The least squares approximate solution is very known.

Example

Showing why least squares

In the least squares approximate solution of $Ax = b$ we, provided that $\|Ax - b\| \neq 0$, select x that the *norm* of the *residual* $r = Ax - b$ is minimal.

It's all there: <http://vmls-book.stanford.edu/> and in many other sources. The least squares approximate solution is very known.

Let's see it.

Example

Showing why least squares

All start with our equation $Ax = b$.

Example

Showing why least squares

Take the original matrix A :

$$A = \begin{bmatrix} 2 & 1 \\ 5 & 1 \\ 7 & 1 \\ 11 & 1 \\ 14 & 1 \\ 18 & 1 \end{bmatrix}$$

Example

Showing why least squares

Take the original matrix A :

$$A = \begin{bmatrix} 2 & 1 \\ 5 & 1 \\ 7 & 1 \\ 11 & 1 \\ 14 & 1 \\ 18 & 1 \end{bmatrix}$$

Transpose it, and get A^T :

$$A^T = \begin{bmatrix} 2 & 5 & 7 & 11 & 14 & 18 \\ 1 & 1 & 1 & 1 & 1 & 1 \end{bmatrix}$$

Example

Showing why least squares

Multiply A^T by A

$$A^T A = \begin{bmatrix} 2 & 5 & 7 & 11 & 14 & 18 \\ 1 & 1 & 1 & 1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 2 & 1 \\ 5 & 1 \\ 7 & 1 \\ 11 & 1 \\ 14 & 1 \\ 18 & 1 \end{bmatrix} = \begin{bmatrix} 719 & 57 \\ 57 & 6 \end{bmatrix}$$

Example

Showing why least squares

Invert the product

$$(A^T A)^{-1} = \begin{bmatrix} 719 & 57 \\ 57 & 6 \end{bmatrix}^{-1} = \begin{bmatrix} \frac{2}{355} & -\frac{19}{355} \\ -\frac{19}{355} & \frac{719}{1065} \end{bmatrix}$$

Example

Showing why least squares

Multiply inverse and transpose

$$\begin{aligned}(A^T A)^{-1} A^T &= \begin{bmatrix} \frac{2}{355} & -\frac{19}{355} \\ -\frac{19}{355} & \frac{719}{1065} \end{bmatrix} \begin{bmatrix} 2 & 5 & 7 & 11 & 14 & 18 \\ 1 & 1 & 1 & 1 & 1 & 1 \end{bmatrix} \\ &= \begin{bmatrix} -\frac{3}{71} & -\frac{9}{355} & -\frac{1}{71} & \frac{3}{355} & \frac{9}{355} & \frac{17}{355} \\ \frac{121}{213} & \frac{434}{1065} & \frac{64}{213} & \frac{92}{1065} & -\frac{79}{1065} & -\frac{307}{1065} \end{bmatrix}\end{aligned}$$

Let's call this product $(A^T A)^{-1} A^T$ the *pseudo-inverse*

Example

Showing why least squares

Finally multiply the pseudo-inverse by b

$$x = (A^T A)^{-1} A^T b \quad (1)$$

$$= \begin{bmatrix} -\frac{3}{\frac{71}{121}} & -\frac{9}{\frac{355}{434}} & -\frac{1}{\frac{64}{213}} & \frac{3}{\frac{355}{92}} & \frac{9}{\frac{355}{79}} & -\frac{17}{\frac{355}{307}} \end{bmatrix} \begin{bmatrix} 5 \\ 5 \\ 8 \\ 7 \\ 9 \\ 7 \end{bmatrix} \quad (2)$$

$$= \begin{bmatrix} \frac{61}{355} \\ \frac{5539}{1065} \end{bmatrix} \quad (3)$$

$$= \begin{bmatrix} 0.171831 \\ 5.20094 \end{bmatrix} \quad (4)$$

Example

Showing why least squares

That's it

$$x = \begin{bmatrix} m \\ b \end{bmatrix} = \begin{bmatrix} \frac{61}{355} \\ \frac{5539}{1065} \end{bmatrix} \text{ or } \begin{bmatrix} 0.171831 \\ 5.20094 \end{bmatrix}$$

and $m = 0.171831$

and $b = 5.20094$

and the line is


$$y = 0.171831x + 5.20094$$

and **that** is the line which **best fit** our data points.

Example

Showing why least squares

And the best fit is



Least squares

Recapitulation

You have a bunch of data points and suspect that a polynomial function $y = Ax + B$ from this points is a good model.

You build a matrix A and a vector b from the polynomial and the points.

The polynomial coefficients are a vector x so $Ax = b$

Then $x = (A^T A)^{-1} A^T b$

Least squares

Don't worry, its XXI century

So the key is calculate $x = (A^T A)^{-1} A^T b$.

But you don't need to do this by hand like Gauss did.

See the code, in this case Javascript code, using the mathjs package.

```
const { multiply, transpose, inv, matrix, format } =  
  require( "mathjs")
```

```
const solve = (A, b) => {  
  const transposed = transpose(A)  
  const product    = multiply(transposed, A)  
  const inverse     = inv(product)  
  const pseudoInverse = multiply(inverse,transposed)  
  const x           = multiply(pseudoInverse, b)  
  return x  
}
```

```
const A = matrix([[2,1],[5,1],[7,1],[11,1],[14,1],[18,1]])  
const b = matrix([5,5,8,7,9,7])
```

```
console.log(format(solve(A, b),5))
```

Least squares

The polynomial function

Get a terminal and do the magic

Least squares

The polynomial function

Get a terminal and do the magic

```
$ node ls.js
```

```
[0.17183, 5.2009]
```

Least squares

Going further

Mind blowing is coming

To avert catastrophic damage in our brains it's time to rename things

In the example we found a line $y = mx + b$, now:

m becomes a_1

b becomes a_0

x becomes x_1

y remains the same

The found line, then, becomes

$$y = a_1x_1 + a_0 = 0.17183x_1 + 5.2009$$

Nothing has changed, only the names.

Least squares

The polynomial function

In the example we picked a line for regression.

A line in a plane can be expressed as a polynomial function $P(x_1)$ where y (the dependent variable, a real number) is a function of x_1 (the independent variable, a real number)

$$y: \mathbb{R} \rightarrow \mathbb{R} = P(x_1) = a_1 x_1 + a_0$$

But there are others $P(x_1)$ s, for example:

$$y: \mathbb{R} \rightarrow \mathbb{R} = P(x_1) = a_2 x_1^2 + a_1 x_1 + a_0$$

What about using that quadratic function as the model for the measured points?

Let's see...

Least squares

A curve as a model

That is the curve: $y = a_2x_1^2 + a_1x_1 + a_0$

And those are the points: (2, 5), (5, 5), (7, 8), (11, 7), (14, 9), (18, 7)

Replacing x_1 and y in each point, and switching terms:

$$a_2x_1^2 + a_1x_1 + a_0 = 5$$