Jude Shin
CSC 4888-S01
Lab 11.2: Exam Questions
November 7, 2026

Module 7 Exam Questions: 3D Vision

1. Definition Question:

Suppose we have detected 2D projections of known 3D points. Suppose also the camera intrinsics for a particular set of images is known. What is the least number of 2D-to-3D correspondences needed to estimate the projection matrix?

(a) 1

(b) 2

(c) 3

(d) 4

Reasoning: *The rotation and translation matrix are the only unknowns in the projection matrix P. Because of this, we can ignore the intrinsics matrix K, eliminating the need to figure out focal length (f) and the camera center (cx, cy). The only variables remaining are rotation and translation. There are 3 degrees of freedom in the rotation matrix, and 3 degrees of freedom in the translation matrix.*

2. Conceptual Question:

Suppose you come across two pairs of stereo images, A and B, that were taken from the same cameras. (Stereo pair A has two images, and Stereo pair B also has two images, resulting in four images total). You find different keypoints in each pair and find that the disparity from the keypoints in pair A is *smaller* than the keypoints in pair B. Which of the following is true about the depth of those keypoints?

(a) The real world point in pair A is *further* from the cameras than in pair B.

(b) The real world point in pair A is *closer* from the cameras than in pair B.

(c) Both real world points in pairs A and B are the *same depth* from the cameras.

(d) It is impossible to tell which real world point is further from the cameras with the given information.

Reasoning: *Intuitively, "things" that are far away from us move less than "things" close to us. (i.e. mountains vs your outstretched hand). In stereo, the disparity is inversely related to the depth of a point.*

3. Computational Question:

A candidate 3D point X is seen by two cameras. Let u_i be the measured pixel location, and let $proj(P_iX)$ be the pixel location predicted from camera P_i.

Suppose the following are the measured pixel locations and the associated projections.

$$u_1 = (4, 3), \qquad proj(P_1X) = (3, 3)$$
$$u_2 = (7, 6), \qquad proj(P_2X) = (7, 2)$$

Calculate the error given the following error model used for triangulation. Assume we are using Euclidean distance.

$$\text{err}(X) = \sum_{i=1}^{n} \|u_i - \text{proj}(P_iX)\|^2$$

(a) -23
(b) 0
(c) 12
(d) 17

Reasoning:

For camera 1:

$$\begin{aligned} err(cam1) &= \|u_1 - proj(P_1X)\|^2 \\ &= \|(4, 3) - (3, 3)\|^2 \\ &= \|(1, 0)\|^2 \\ &= 1^2 + 0^2 \\ &= 1 \end{aligned}$$

For camera 2:

$$\begin{aligned} err(cam2) &= \|u_2 - proj(P_2X)\|^2 \\ &= \|(7, 6) - (7, 2)\|^2 \\ &= \|(0, 4)\|^2 \\ &= 0^2 + 4^2 \\ &= 16 \end{aligned}$$

Squared error for camera 1: 1.

Squared error for camera 2: 16.

Total error: $1 + 16 = 17$