



0025-5408(95)00116-6

STRUCTURE OF $(VO)_2P_2O_7$

P.T. Nguyen, R.D. Hoffman, and A.W. Sleight
Department of Chemistry, Oregon State University,
Corvallis, Oregon 97331-4003

(Received June 21, 1995; Communicated by A.J. Jacobson)

ABSTRACT

The crystal structure of the vanadium (IV) pyrophosphate that shows the remarkable ability to selectively catalyze the oxidation of butane to maleic anhydride has been accurately determined for the first time. This was facilitated by the growth of high quality $(VO)_2P_2O_7$ crystals under controlled oxygen fugacity. Single crystal X-ray diffraction indicated monoclinic symmetry rather than the orthorhombic symmetry assumed in earlier studies. The space group is $P2_1$ with $a = 7.28 \text{ \AA}$, $b = 16.588 \text{ \AA}$, $c = 9.58 \text{ \AA}$, $\beta = 89.975^\circ$, and $Z = 4$. A final R of 3.1% was obtained. Typical vanadyl environments are found for all eight crystallographically distinct vanadium atoms, and normal bond angles and distances are found for the P_2O_7 groups.

MATERIALS INDEX: vanadium, phosphate, catalyst

INTRODUCTION

One of the most remarkable heterogeneous catalysts is $(VO)_2P_2O_7$. If prepared according to certain procedures, this compound can catalyze the oxidation of *n*-butane to maleic anhydride at selectivities exceeding 80% (1). No other compound is known to catalyze this commercially important reaction at selectivities approaching 80%. No other partial oxidation of a saturated hydrocarbon is known to be so selective. Despite the uniqueness and importance

of $(VO)_2P_2O_7$ as a catalyst, it is poorly understood. Three previous crystal structure studies (2-4) did not lead to a good definition of the structure.

The first single crystal X-ray diffraction study (2) on $(VO)_2P_2O_7$ resulted in a final R of 19.8%. A second study (3) yielded an R of 8.9%, but some of the interatomic distances seemed unreasonable. Both of these studies reported essentially the same structural arrangement (Fig. 1). The $(VO)_2P_2O_7$ structure has recently been redetermined by Thompson and Ebner (4). They obtained good agreement factors ($R = 3.4\%$); however, they could obtain this result only with a disordered model. Streaking effects in their diffraction patterns also indicated disorder.




FIG 1.

The structure of $(VO)_2P_2O_7$, where the vanadyl groups are shown as square pyramids and the pyrophosphate groups are shown as a pair of corner sharing tetrahedra.

We have now obtained single crystals of $(VO)_2P_2O_7$ which give diffraction patterns free from any indications of disorder. No streaking was observed in our diffraction patterns, and a good refinement was obtained based on a fully ordered structure.

PREPARATION OF CRYSTALS

Polycrystalline $(VO)_2P_2O_7$ was prepared by dehydration of $(VO)_2P_2O_7 \cdot xH_2O$. Approximately 5 g of $(VO)_2P_2O_7$ was pressed into a 10-mm-diameter pellet. This pellet was placed in a Pt crucible and heated in a furnace where oxygen fugacity was controlled and continuously monitored. After rapid heating to 500°C, the sample was heated to 910°C at 50°/hour. After a hold period of 15 hours, the sample was cooled at 2°/hour to 890°C and then cooled at 4°/hour to 855°C. Power to the furnace was terminated, and the sample was cooled to room temperature in the furnace. This procedure was repeated at several oxygen fugacities between 0.001 atm and 0.003 atm.

Crystals prepared when using the lower oxygen pressure (~0.001 atm) were green, but those recovered from preparations at higher oxygen pressures were red-brown. The titration method of Nakamura et al. (5) was used to determine the oxidation state of vanadium in the crystals. The result was 4.00 for the green crystals and 4.01 for the red-brown crystals. Diffraction patterns taken on a precession camera indicated that the green crystals were of higher quality from a diffraction point of view. Therefore, only those crystals were used in further structural studies.

TABLE 1
Crystal Data and Intensity Collection for $(VO)_2P_2O_7$

| | |
|--|--|
| Empirical formula | $V_2P_2O_8$ |
| Formula weight (g/mol) | 307.82 |
| Space Group | $P2_1$ (no. 4) |
| a (Å) | 7.7276(3) |
| b (Å) | 16.5885(4) |
| c (Å) | 9.5796(5) |
| β | 89.975(3) |
| V (Å ³) | 1228(1) |
| Z | 4 |
| Diffractometer | Rigaku AFC6R |
| Radiation | MoK_{α} ($\lambda = 0.71069$ Å) graphite-monochromated |
| Temperature | 23°C |
| Maximum 2θ | 70° |
| Data collected | $-9 < h < 9, -21 < k < 21, -12 < l < 12$ |
| Scan speed (degrees/min) | 16.0 in ω and 32.0 in 2θ |
| No. of unique data with $F_{obs} > 3\sigma(F_{obs})$ | 4254 |
| R_{merge} | 3.6% |
| Data/parameter ratio | 14.72 |
| R | 0.031 |
| R_w | 0.044 |

STRUCTURE DETERMINATION

Single crystal diffraction data were collected on a Rigaku AFC6R diffractometer. Despite the apparent orthorhombic symmetry, a full sphere of reflections was collected (Table 1). The intensities of three standard reflections measured every 300 reflections throughout the data collection exhibited no significant excursions. The structure was solved and refined with programs from the TEXAN software packages (6). The absorption correction was performed with the DIFABS software (7).

TABLE 2
Atomic and Thermal Parameters for $(VO)_2P_2O_7$

| Atom | x | y | z | Beq (\AA^2)* | Atom | x | y | z | Beq (\AA^2) |
|------|------------|-------------|------------|-------------------------|------|-----------|------------|------------|------------------------|
| V1 | 0.2973(2) | 0.0958 | 0.4912(1) | 0.47(5) | O11 | 0.4960(7) | -0.2162(3) | 0.2871(4) | 0.59(7) |
| V2 | 0.2039(2) | 0.5960(1) | 0.4909(1) | 0.36(5) | O12 | 0.0066(8) | 0.2846(3) | 0.2862(5) | 0.79(8) |
| V3 | 0.2087(2) | -0.09442(9) | 0.5023(1) | 0.33(5) | O13 | 0.3007(9) | 0.1734(3) | 0.6421(5) | 1.0(1) |
| V4 | 0.2923(2) | 0.4057(1) | 0.5017(1) | 0.25(5) | O14 | 0.2017(8) | 0.6734(3) | 0.6410(5) | 0.71(9) |
| V5 | 0.2055(2) | -0.1548(1) | -0.0005(2) | 0.45(6) | O15 | 0.2133(8) | 0.1761(3) | 0.3613(5) | 0.44(8) |
| V6 | 0.2935(2) | 0.3453(1) | -0.0002(1) | 0.43(6) | O16 | 0.2943(9) | 0.6756(3) | 0.3600(5) | 0.9(1) |
| V7 | 0.2081(2) | 0.1561(1) | -0.0014(1) | 0.34(5) | O17 | 0.281(1) | 0.0010(4) | 0.6307(6) | 0.9(1) |
| V8 | 0.2914(2) | 0.65607(7) | -0.0021(1) | 0.33(5) | O18 | 0.2200(8) | 0.5007(3) | 0.6300(5) | 0.3(1) |
| P1 | 0.2952(3) | -0.0003(1) | 0.7924(2) | 0.3(1) | O19 | 0.2221(9) | 0.0017(4) | 0.3639(5) | 0.4(1) |
| P2 | 0.2026(4) | 0.4997(2) | 0.7925(2) | 0.39(8) | O30 | 0.276(1) | 0.5012(4) | 0.3619(6) | 0.7(1) |
| P3 | 0.1948(3) | 0.0005(2) | 0.2012(2) | 0.3(1) | O31 | 0.2860(9) | -0.1795(3) | 0.6294(6) | 0.7(1) |
| P4 | 0.3031(4) | 0.5008(1) | 0.2012(2) | 0.24(8) | O32 | 0.2117(9) | 0.3207(3) | 0.6287(5) | 0.6(1) |
| P5 | 0.1920(4) | -0.2447(2) | 0.7064(2) | 0.4(1) | O33 | 0.199(1) | -0.1772(3) | 0.3578(5) | 0.8(1) |
| P6 | 0.3106(4) | 0.2556(1) | 0.7061(2) | 0.37(9) | O34 | 0.298(1) | 0.3230(3) | 0.3572(5) | 0.7(1) |
| P7 | 0.3020(3) | -0.2436(1) | 0.2907(2) | 0.4(1) | O35 | 0.2659(9) | 0.0775(3) | 0.1395(5) | 1.0(1) |
| P8 | 0.2008(4) | 0.2562(1) | 0.2906(2) | 0.27(8) | O36 | 0.2355(8) | 0.5768(3) | 0.1390(5) | 0.67(9) |
| O1 | 0.5014(8) | 0.5914(3) | 0.5399(4) | 0.59(8) | O37 | 0.2274(8) | 0.0749(3) | -0.1455(5) | 0.79(9) |
| O2 | -0.0024(8) | 0.0933(3) | 0.5397(4) | 0.76(8) | O38 | 0.2809(9) | 0.5743(3) | -0.1465(5) | 1.05(9) |
| O3 | 0.005(1) | -0.0904(3) | 0.5313(5) | 0.8(1) | O39 | 0.2727(8) | -0.0754(3) | 0.1391(6) | 0.6(1) |
| O4 | 0.496(1) | 0.4121(3) | 0.5320(4) | 0.41(9) | O50 | 0.2329(9) | 0.4247(3) | 0.1390(6) | 0.7(1) |
| O5 | -0.001(1) | -0.1597(3) | 0.0131(6) | 0.59(8) | O51 | 0.2234(9) | -0.0768(3) | -0.1513(5) | 0.7(1) |
| O6 | 0.500(1) | 0.3435(4) | 0.0115(7) | 0.9(1) | O52 | 0.2823(9) | 0.4233(3) | -0.1514(6) | 0.8(1) |
| O7 | -0.0002(9) | 0.1599(3) | 0.0125(6) | 0.67(8) | O53 | 0.2510(8) | -0.2504(3) | 0.1328(6) | 0.3(1) |
| O8 | 0.500(1) | 0.6595(3) | 0.0108(6) | 0.88(8) | O54 | 0.255(1) | 0.2497(4) | 0.1352(6) | 0.8(1) |
| O9 | 0.4975(8) | -0.0014(3) | 0.8272(6) | 1.55(9) | O55 | 0.2496(9) | -0.2482(4) | -0.1394(6) | 0.6(1) |
| O10 | 0.0044(8) | 0.4983(3) | 0.8252(6) | 1.41(9) | O56 | 0.2535(9) | 0.2517(4) | -0.1382(6) | 0.5(1) |

*Beq = $(8\pi^2/3)\sum_i U_{ij}a_i^*a_j^*$

Refinements were initially carried out in space groups $Pcam$ and $Pca2_1$ because these were the space groups that earlier investigators (2-4) had considered. However, significant peaks near vanadium atoms were always found in the final difference electron density maps. Furthermore, three different crystals clearly showed reflections which violate these space groups both in precession photographs and in data collected with the diffractometer. Thirty-seven of these "forbidden" reflections had an observed intensity higher than six times sigma. The only space group consistent with the observed systematic absences is $P2_1$. Refinement in this space group was successful (Table 1), and the final difference Fourier map was featureless. The final positional parameters are given in Table 2, and the final thermal parameters are given in Table 3. Bond distances are given in Table 4, and selected bond angles are given in Table 5.

TABLE 3
Anisotropic Thermal Parameters ($\times 10^{-4}\text{\AA}$) for $(VO)_2P_2O_7$

| Atom | U11 | U22 | U33 | U12 | U13 | U23 |
|------|------|-------|-------|--------|--------|--------|
| V1 | 7(1) | 59(6) | 53(5) | -15(5) | 16(7) | 21(6) |
| V2 | 4(1) | 56(6) | 45(5) | 10(5) | 1(7) | -26(5) |
| V3 | 3(1) | 60(6) | 36(5) | 29(5) | 1(7) | -17(5) |
| V4 | 6(9) | 43(6) | 45(5) | -2(4) | 6(7) | 27(5) |
| V5 | 8(1) | 40(6) | 47(5) | 21(5) | 4(7) | -16(6) |
| V6 | 7(1) | 52(6) | 46(5) | -4(5) | -18(7) | 32(6) |
| V7 | 6(1) | 38(5) | 33(5) | 15(5) | 12(7) | -20(5) |
| V8 | 5(1) | 40(5) | 37(5) | -20(5) | -6(6) | 16(5) |
| P1 | 1(1) | 8(1) | 17(8) | 8(8) | 1(1) | 6(6) |
| P2 | 6(1) | 5(1) | 33(8) | 11(9) | -20(1) | -6(6) |
| P3 | 2(1) | 7(1) | 37(9) | -19(8) | -1(1) | -1(6) |
| P4 | 1(1) | 6(1) | 17(8) | 14(8) | -10(9) | -4(6) |
| P5 | 9(2) | 6(1) | 2(1) | -26(9) | 1(1) | 7(6) |
| P6 | 5(1) | 4(1) | 48(9) | 6(9) | -1(1) | 5(6) |
| P7 | 6(2) | 5(1) | 4(1) | -4(9) | 10(1) | 6(6) |
| P8 | 3(1) | 5(1) | 23(8) | 10(8) | 1(1) | -11(6) |

TABLE 4
Bond Distances (Å) for (VO)₂P₂O₇

| | | | | | |
|----|-----|----------|----|-----|----------|
| V1 | O1 | 1.585(6) | V8 | O7 | 2.253(7) |
| | O2 | 2.337(6) | | O8 | 1.619(7) |
| | O13 | 1.936(5) | | O36 | 1.934(5) |
| | O15 | 1.935(5) | | O38 | 1.939(5) |
| | O17 | 2.067(6) | | O53 | 2.043(6) |
| | O19 | 2.065(6) | | O55 | 2.088(6) |
| V2 | O1 | 2.307(6) | P1 | O9 | 1.599(7) |
| | O2 | 1.585(6) | | O17 | 1.554(6) |
| | O14 | 1.929(5) | | O37 | 1.478(5) |
| | O16 | 1.951(6) | P2 | O51 | 1.486(6) |
| | O18 | 2.071(6) | | O10 | 1.564(7) |
| | O30 | 2.075(7) | | O18 | 1.563(5) |
| V3 | O3 | 1.597(8) | P3 | O38 | 1.497(6) |
| | O4 | 2.307(7) | | O52 | 1.509(6) |
| | O17 | 2.081(7) | | O10 | 1.560(7) |
| | O19 | 2.076(6) | P4 | O19 | 1.573(5) |
| | O31 | 1.957(6) | | O35 | 1.510(6) |
| | O33 | 1.951(5) | | O39 | 1.518(6) |
| V4 | O3 | 2.323(8) | P5 | O9 | 1.565(7) |
| | O4 | 1.606(7) | | O30 | 1.554(6) |
| | O18 | 2.075(6) | | O36 | 1.489(6) |
| | O30 | 2.079(6) | P6 | O50 | 1.497(6) |
| | O32 | 1.964(6) | | O12 | 1.612(6) |
| | O34 | 1.949(5) | | O14 | 1.498(5) |
| V5 | O5 | 1.602(8) | P7 | O31 | 1.498(6) |
| | O6 | 2.277(9) | | O55 | 1.544(6) |
| | O39 | 1.948(6) | | O11 | 1.568(6) |
| | O51 | 1.945(5) | P8 | O13 | 1.497(6) |
| | O53 | 2.066(6) | | O32 | 1.516(6) |
| | O55 | 2.070(6) | | O56 | 1.557(6) |
| V6 | O5 | 2.266(8) | P7 | O11 | 1.567(6) |
| | O6 | 1.601(9) | | O16 | 1.496(5) |
| | O50 | 1.932(6) | | O33 | 1.506(6) |
| | O52 | 1.943(5) | P8 | O53 | 1.568(6) |
| | O54 | 2.071(6) | | O12 | 1.574(6) |
| | O56 | 2.062(6) | | O15 | 1.495(5) |
| V7 | O7 | 1.617(7) | P8 | O34 | 1.484(6) |
| | O8 | 2.257(7) | | O54 | 1.549(6) |
| | O35 | 1.930(5) | | | |
| | O37 | 1.935(5) | | | |
| | O54 | 2.062(6) | | | |
| | O56 | 2.087(6) | | | |

TABLE 5
Bond Angles (°) for $(VO)_2P_2O_7$

| Bond | Angle (°) | Bond | Angle (°) | Bond | Angle (°) | Bond | Angle (°) |
|------------|-----------|------------|-----------|------------|-----------|------------|-----------|
| O1-V1-O13 | 99.0(3) | O38-V8-O53 | 167.4(3) | O19-V3-O31 | 159.4(3) | O31-P5-O55 | 111.0(3) |
| O1-V1-O15 | 103.9(2) | O38-V8-O55 | 94.3(2) | O19-V3-O33 | 95.1(2) | O11-P6-O13 | 109.8(4) |
| O1-V1-O17 | 98.3(3) | O53-V8-O55 | 78.3(2) | O31-V3-O33 | 86.9(2) | O11-P6-O32 | 106.7(3) |
| O1-V1-O19 | 97.5(3) | O9-P1-O17 | 106.0(4) | O3-V4-O4 | 174.0(2) | O11-P6-O56 | 104.0(3) |
| O13-V1-O15 | 91.6(2) | O9-P1-O37 | 105.8(3) | O3-V4-O18 | 78.0(2) | O13-P6-O32 | 115.0(3) |
| O13-V1-O17 | 91.4(2) | O9-P1-O51 | 106.2(3) | O3-V4-O30 | 80.3(2) | O13-P6-O56 | 109.9(3) |
| O13-V1-O19 | 161.4(2) | O17-P1-O37 | 111.4(4) | O3-V4-O32 | 7.9(2) | O32-P6-O56 | 110.8(4) |
| O15-V1-O17 | 156.8(3) | O17-P1-O51 | 110.3(3) | O3-V4-O34 | 86.9(2) | O11-P7-O16 | 108.0(3) |
| O15-V1-O19 | 92.6(2) | O37-P1-O51 | 116.3(3) | O4-V4-O18 | 96.1(2) | O11-P7-O33 | 107.7(3) |
| O17-V1-O19 | 77.9(2) | O10-P2-O18 | 106.5(4) | O4-V4-O30 | 97.2(3) | O11-P7-O53 | 103.9(3) |
| O2-V2-O14 | 98.5(3) | O10-P2-O38 | 109.3(4) | O4-V4-O32 | 104.3(3) | O16-P7-O33 | 116.4(3) |
| O2-V2-O16 | 104.6(3) | O10-P2-O52 | 108.4(3) | O4-V4-O34 | 98.8(3) | O16-P7-O53 | 110.7(3) |
| O2-V2-O18 | 99.0(3) | O18-P2-O38 | 110.2(3) | O18-V4-O30 | 77.7(2) | O33-P7-O53 | 109.3(3) |
| O2-V2-O30 | 97.6(3) | O18-P2-O52 | 109.2(3) | O18-V4-O32 | 95.3(2) | O12-P8-O15 | 109.9(3) |
| O14-V2-O16 | 91.8(2) | O38-P2-O52 | 113.0(4) | O18-V4-O34 | 163.9(3) | O12-P8-O34 | 105.7(3) |
| O14-V2-O18 | 91.6(2) | O10-P3-O19 | 107.1(3) | O30-V4-O32 | 158.1(3) | O12-P8-O54 | 104.5(3) |
| O14-V2-O30 | 161.9(2) | O10-P3-O35 | 108.4(3) | O30-V4-O34 | 94.6(2) | O15-P8-O34 | 115.9(3) |
| O16-V2-O18 | 155.3(3) | O10-P3-O39 | 107.9(3) | O32-V4-O34 | 86.7(2) | V2-O30-P4 | 129.0(4) |
| O16-V2-O30 | 92.0(2) | O19-P3-O35 | 109.2(3) | O5-V5-O6 | 175.9(2) | V4-O30-P4 | 128.8(4) |
| O18-V2-O30 | 77.8(2) | O19-P3-O39 | 110.2(3) | O5-V5-O39 | 104.2(3) | V3-O31-P5 | 132.8(4) |
| O3-V3-O4 | 174.6(2) | O35-P3-O39 | 113.8(3) | O52-V6-O56 | 91.0(2) | V4-O32-P6 | 130.9(4) |
| O3-V3-O17 | 97.5(3) | O9-P4-O30 | 107.6(4) | O54-V6-O56 | 78.6(2) | V3-O33-P7 | 142.8(4) |
| O3-V3-O19 | 97.3(3) | O9-P4-O36 | 107.2(3) | O7-V7-O8 | 175.6(2) | V4-O34-P8 | 145.1(4) |
| O3-V3-O31 | 102.9(3) | O15-P8-O54 | 110.9(3) | O7-V7-O35 | 101.5(3) | V7-O35-P3 | 139.5(4) |
| O3-V3-O33 | 96.5(3) | O34-P8-O54 | 109.2(3) | O7-V7-O37 | 99.3(3) | V8-O36-P4 | 140.9(4) |
| O4-V3-O17 | 77.4(2) | V3-O3-V4 | 161.7(3) | O7-V7-O54 | 95.3(3) | V7-O37-P1 | 154.4(4) |
| O5-V5-O51 | 99.5(3) | V3-O4-V4 | 160.2(3) | O7-V7-O56 | 100.9(3) | V8-O38-P2 | 150.9(4) |
| O5-V5-O53 | 94.6(3) | V5-O5-V6 | 170.8(3) | O8-V7-O35 | 79.3(2) | V5-O39-P3 | 136.4(4) |
| O5-V5-O55 | 100.3(3) | V5-O6-V6 | 173.1(4) | O8-V7-O37 | 85.0(2) | V6-O50-P4 | 139.6(4) |
| O6-V5-O39 | 76.9(2) | V7-O7-V8 | 171.8(3) | O8-V7-O54 | 80.3(2) | V5-O51-P1 | 149.8(4) |
| O6-V5-O51 | 84.4(2) | V7-O8-V8 | 172.5(3) | O8-V7-O56 | 77.8(2) | V6-O52-P2 | 147.5(4) |
| O6-V5-O53 | 81.3(2) | P1-O9-P4 | 157.9(4) | O35-V7-O37 | 90.6(2) | V5-O53-V8 | 102.6(2) |
| O6-V5-O55 | 8.2(2) | P2-O10-P3 | 159.0(4) | O35-V7-O54 | 91.4(2) | V5-O53-P7 | 125.8(3) |
| O39-V5-O51 | 92.4(2) | P6-O11-P7 | 145.5(3) | O35-V7-O56 | 156.1(3) | V8-O53-P7 | 128.8(3) |
| O39-V5-O53 | 92.8(2) | P5-O12-P8 | 144.7(3) | O37-V7-O54 | 164.5(3) | V6-O54-V7 | 101.8(3) |
| O39-V5-O55 | 154.6(3) | V1-O13-P6 | 155.8(3) | O37-V7-O56 | 93.9(2) | V6-O54-P8 | 126.0(4) |
| O51-V5-O53 | 163.3(2) | V2-O14-P5 | 156.4(3) | O54-V7-O56 | 78.3(2) | V7-O54-P8 | 128.0(4) |
| O51-V5-O55 | 90.5(2) | V1-O15-P8 | 157.7(4) | O7-V8-O8 | 175.9(2) | V5-O55-V8 | 100.9(3) |
| O53-V5-O55 | 78.2(2) | V2-O16-P7 | 154.9(4) | O7-V8-O36 | 80.1(2) | V5-O55-P5 | 122.7(4) |
| O5-V6-O6 | 176.7(2) | V1-O17-V3 | 102.3(3) | O7-V8-O38 | 86.9(2) | V8-O55-P5 | 132.5(4) |
| O5-V6-O50 | 79.7(2) | V1-O17-P1 | 130.6(4) | O7-V8-O53 | 81.6(2) | V6-O56-V7 | 101.2(2) |
| O5-V6-O52 | 86.5(3) | V3-O17-P1 | 126.7(4) | O7-V8-O55 | 78.2(2) | V6-O56-P6 | 122.7(3) |
| O5-V6-O54 | 82.0(2) | V2-O18-V4 | 102.4(2) | O8-V8-O36 | 101.1(3) | V7-O56-P6 | 132.9(4) |
| O5-V6-O56 | 77.7(2) | V2-O18-P2 | 130.1(3) | O8-V8-O38 | 97.0(3) | | |
| O6-V6-O50 | 101.9(3) | V4-O18-P2 | 127.3(3) | O8-V8-O53 | 94.4(3) | | |
| O6-V6-O52 | 96.3(3) | V1-O19-V3 | 102.6(2) | O8-V8-O55 | 100.1(3) | | |
| O6-V6-O54 | 94.9(3) | V1-O19-P3 | 129.2(4) | O9-P4-O50 | 105.5(3) | | |
| O6-V6-O56 | 100.4(3) | V3-O19-P3 | 128.1(3) | O30-P4-O36 | 110.2(4) | | |
| O50-V6-O52 | 92.9(2) | V2-O30-V4 | 102.1(3) | O30-P4-O50 | 110.4(3) | | |
| O50-V6-O54 | 93.1(2) | O4-V3-O19 | 79.9(2) | O36-P4-O50 | 115.3(3) | | |
| O50-V6-O56 | 156.8(3) | O4-V3-O31 | 79.7(2) | O12-P5-O14 | 109.9(3) | | |
| O52-V6-O54 | 165.9(3) | O4-V3-O33 | 88.4(2) | O12-P5-O31 | 105.4(3) | | |
| O36-V8-O38 | 90.8(2) | O17-V3-O19 | 77.3(2) | O12-P5-O55 | 104.1(3) | | |
| O36-V8-O53 | 92.3(2) | O17-V3-O31 | 95.7(2) | O14-P5-O31 | 115.1(3) | | |
| O36-V8-O55 | 157.3(3) | O17-V3-O33 | 164.8(3) | O14-P5-O55 | 110.6(3) | | |

DISCUSSION

The vanadium atoms in $(VO_2)_2P_2O_7$ show a typical vanadyl arrangement (Table 4): four nearly coplanar V-O bonds of about 2 Å, one short V-O bond of about 1.6 Å opposite a long V-O bond of about 2.3 Å. Likewise, the P_2O_7 groups show normal bond lengths and angles (Tables 4 and 5). The P-O-P angles lie within the normal range of 130° to 160°. The structure we find (Fig. 1) for $(VO)_2P_2O_7$ is basically the same as that found in previous studies (2-4). There are no differences in the way in which the vanadyl and pyrophosphate groups are connected.

We conclude that much of the problem with earlier single crystal studies (2-4) of $(VO_2)_2P_2O_7$ was due to the use of an incorrect space group. The deviation from orthorhombic symmetry is small but significant. Another difference between our study and the previous studies is the quality of our crystals. None of the streaking effects reported in the previous studies (2-4) were evident in the diffraction pattern of our green crystals. It is likely that the better quality of our crystals can be attributed to our control oxygen fugacity during crystal growth. Ebner and Thompson (4) also reported both green and red-brown crystals, and they also indicated that the red-brown crystals were of poorer quality. They did not, however, show that the red-brown crystals have an average vanadium oxidation state higher than four.

Bond valence calculations (8) based on our structure for $(VO)_2P_2O_7$ show a range of 4.02 to 4.18 for the eight vanadium atoms and a range of 4.76 to 4.97 for the eight phosphorous atoms. These are well within the range expected for V^{4+} and P^{5+} , and this indicates that all vanadium atoms and all phosphorous atoms are essentially equivalent chemically. The bond valences calculated (9) based on the best previously available structural data give a range of 3.69 to 4.44 for vanadium. These values are smaller and larger, respectively, than expected for V^{4+} , and this confirms the fact that our structure is considerably more accurate.

The $(VO)_2P_2O_7$ structure presented here should not be regarded as the structure of the real vanadyl pyrophosphate catalyst. The real catalyst gives an X-ray diffraction pattern similar to that generated from parameters determined in this study. However, there are peak shifts and broadening effects due to extended defects (10). These defects can account for the observation that the real catalysts always have an average vanadium oxidation state significantly higher than four (e.g., ~4.1). These same defects can also account for the streaking effects observed by Ebner and Thompson (4) in single crystal X-ray studies and by Bordes and Courtine (11) in electron diffraction studies. One of the most significant conclusions of our work is that it is in fact possible to obtain $(VO)_2P_2O_7$ crystals which are relatively free of both extended and point defects.

ACKNOWLEDGMENT

This work was supported in part by DuPont de Nemours and Company.

REFERENCES

1. R.M. Contractor, H.E. Bergna, H.S. Horowitz, C.M. Blackstone, U. Chowdhry, and A.W. Sleight, *Catalysis* **198**, Elsevier, 645 (1988).
2. N.E. Middlemiss, doctoral dissertation, Dept. of Chemistry, McMaster Univer., Hamilton, Ontario, Canada (1978)
3. Yu.E. Gorbunova and S.A. Linde, *Sov. Phys. Dokl.* **24**, 138 (1979)
4. J.R. Ebner and M.R. Thompson, *Studies in Surface Science and Catalysis*, ed. R.K. Grasselli and A.W. Sleight, Elsevier (1992)
5. M. Nakamura, K. Kawai, and Y. Fujiwara, *J. Catal.* **34**, 345 (1974).
6. TEXSAN, Molecular Structure Corporation, The Woodlands, TX, USA, (1988)
7. N. Walker and A. Stuart, *Acta Cryst.* **A39**, 158-166, (1983)
8. I.D. Brown and K.K. Wu, *Acta Cryst.* **B32**, 1957 (1976).
9. E. Bordes, *Catalysis Today* **1**, 499 (1987).
10. P.T. Nguyen, N. Roberts, W.W. Warren, and A.W. Sleight, *14th North American Meeting of the Catalysis Society*, T-258 (1995)
11. E. Bordes and P. Courtine, *J. Catal.* **57**, 236-252 (1979).