

Extended Results and Diagnostics for the Paper Moving Beyond Crude Trial–Real-World Comparisons Through Population Alignment and Decomposition

Juho Lähteenmaa, Pauliina Ilmonen, Elli-Noora Hännikäinen,
Joonas Laaksolilja, Anders Ståhls, Kirsi Santti, Peeter Karihtala, Johanna Mattson

This document accompanies the paper *Moving Beyond Crude Trial–Real-World Comparisons Through Population Alignment and Decomposition* and provides extended results and diagnostic information that complement the main article and supplementary information.

1 Results

This section presents extended results from the trial–real-world comparisons in stages corresponding to the analytical framework described in the article. We first summarize results from the multilevel outcome models fitted to the real-world cohort, which characterize associations between baseline variables and pCR. We then compare observed pCR rates between the real-world cohort and the randomized trial arms and present population-aligned estimates obtained using MAIC and Bayesian MRP with raking. Finally, we decompose the adjustment differences into variable-level contributions.

All results are descriptive. Baseline alignment is restricted to variables that are observed and harmonizable across the real-world and trial populations. Differences arising from unmeasured baseline characteristics, treatment implementation, and other contextual factors may therefore contribute to the residual differences, and limited covariate overlap can increase uncertainty in the population-aligned estimates.

1.1 Results of the Outcome Models

Each real-world–trial comparison uses a distinct set of baseline variables, resulting in six multilevel logistic regression models: three models based solely on baseline characteristics and three that additionally include an indicator of treatment discontinuation. This structure allows assessment of the robustness of estimated associations to model specification.

Table 1: Baseline characteristics of the HUS cohort.

Characteristic	N	Summary
Median age (n = 268)	—	52 (42–63)
Tumor size, median (mm)	221	26.0 (20–37)
Tumor size ≤ 20 mm	58	22% [17–27%]
Tumor size 20–50 mm	143	53% [47–60%]
Tumor size > 50 mm	19	7% [4–11%]
Tumor size: unknown	47	18%
Nodal status: positive	155	58% [52–64%]
Nodal status: negative	93	35% [29–41%]
Nodal status: unknown	20	8% [5–11%]
Bilateral breast cancer	8	3% [1–6%]
Unilateral breast cancer	260	97% [94–99%]
Bilaterality: unknown	0	—
Histology: ductal	211	79% [73–84%]
Histology: lobular	13	5% [3–8%]
Histology: other	44	16% [12–21%]
Histology: unknown	0	—
Tumor grade 1	< 3	$< 1\%$ [0–2%]
Tumor grade 2	64	24% [19–29%]
Tumor grade 3	152	57% [51–63%]
Tumor grade: unknown	51	19% [15–24%]
ER and/or PR positive	88	33% [27–39%]
ER and PR negative	180	67% [61–73%]
ER or PR unknown	0	—
HER2 IHC/ISH: IHC 2+ and ISH+	71	27% [21–32%]
HER2 IHC/ISH: IHC 3+	183	68% [62–74%]
HER2 IHC/ISH: other	14	5% [3–9%]
HER2 IHC/ISH: unknown	0	—
Ki-67, median (n = 263)	—	40 (40–60)
Ki-67 $\leq 30\%$	91	34% [28–40%]
Ki-67 $> 30\%$	172	64% [58–70%]
Ki-67: unknown	5	2% [0–4%]

Notes: Continuous variables are reported as median (interquartile range). Categorical variables are reported as n (%) with 95% binomial confidence intervals. Counts between 1 and 2 are masked as “ < 3 ” to preserve patient confidentiality.

Abbreviations: ER, estrogen receptor; PR, progesterone receptor; HER2, human epidermal growth factor receptor 2; IHC, immunohistochemistry; ISH, in situ hybridization; Ki-67, proliferation index.

These outcome models are not intended to estimate causal effects or to

provide optimal predictive performance. Instead, covariate selection and discretization are guided by the objective of harmonizing marginal distributions between the real-world and trial populations. Posterior summaries of coefficients for all six models are reported in Supplementary Tables 5–10. All the coefficients are identified via a sum-to-zero constraint and should be interpreted as deviations from the overall mean of the linear predictor. Pairwise differences between categories are obtained by subtracting the corresponding coefficients.

Despite their descriptive role, several baseline variables show consistent associations with pCR across model specifications. Pre-treatment hormone receptor status is among the most strongly associated variables. As an example in the HELEN-006 outcome model (Supplementary Table 9), hormone receptor positivity has a mean posterior coefficient of -0.64 with a 94% highest density interval (HDI) from -0.93 to -0.35 , which is consistent over all the models. Using the “divide-by-4” rule as a rough approximation, this coefficient of -0.64 corresponds to an upper bound of approximately 16 percentage points difference in pCR rates between hormone-positive and hormone-negative patients, holding other variables constant. The magnitude of this association is broadly consistent with published results in similar patient populations, while lying near the upper end of reported estimates Canino et al. [2024], Zhou et al. [2022], Kim et al. [2022], Falcón González et al. [2024], Hall et al. [2022].

Hormone receptor positivity was associated with lower modeled pCR rates, while higher HER2 expression was associated with higher modeled pCR rates in the HUS sample. For example, comparing IHC3+ to IHC2+ corresponds to an approximate difference of 35 percentage points under the same approximation. Some other coefficients show less intuitive patterns, such as T3 tumors having comparable or higher modeled pCR rates than T2 tumors, which may reflect model misspecification, omitted interactions, or other unmeasured differences.

Individual coefficients should be interpreted cautiously. Some variables in the HELEN-006 model are structurally related (e.g., stage, tumor size, and nodal status), which may introduce collinearity and reduce interpretability of individual parameters. This issue affects interpretation of individual coefficients but does not materially influence the post-stratified predictions that form the basis of the population-alignment analysis. In addition, many posterior intervals include zero, reflecting substantial uncertainty about the sign of the coefficient, particularly for imbalanced variables such as bilaterality.

Coefficients for NAT discontinuation are modest in magnitude in KRISTINE and HELEN-006 variable outcome models (Supplementary Table 8 and Supplementary Table 10), suggesting limited association with modeled pCR rates. However, in the NeoSphere variable outcome model, discontinuation is defined using a lower treatment threshold than in the KRISTINE and HELEN-006 comparisons, which may partly explain the larger magnitude (more negative) but more uncertain coefficient estimates observed in that setting.

1.2 Comparison of pCR Rates Across RCT, Real-World, and Population-Aligned Estimates

We next compare pCR rates between the HUS cohort and each RCT arm, considering both observed and population-aligned estimates. For each trial, we present (i) the unadjusted comparison, (ii) the adjusted HUS estimate obtained by aligning baseline distributions, (iii) the resulting adjustment and residual differences, and (iv) Shapley decomposition of the adjustment difference $\Delta\hat{\theta}_{\text{adj}}$. The same analytical framework is applied separately to the NeoSphere, KRISTINE, and HELEN-006 comparisons.

1.2.1 NeoSphere comparison

We first estimate unadjusted posterior distributions for pCR rates in the HUS cohort and in each RCT arm using binomial models. Posteriors are obtained from the observed sample sizes and numbers of patients achieving pCR in each group. A uniform Beta(1, 1) prior is used, as the likelihood dominates given the sample sizes.

For the HUS cohort, the posterior mean pCR rate $\hat{\theta}_{\text{RWD}}$ is 0.54 with a 95% HDI from 0.49 to 0.60. For the NeoSphere arm B group, the posterior mean is 0.46 with a 95% HDI from 0.37 to 0.55. The posterior for the observed difference $\Delta\hat{\theta}_{\text{obs}} = \hat{\theta}_{\text{RCT}} - \hat{\theta}_{\text{RWD}}$ has mean -0.09 , with a posterior probability of 94% that the NeoSphere pCR rate is lower than the HUS rate.

We then adjust for differences in baseline marginals using Bayesian MRP with raking and MAIC. Under MRP with raking, the adjusted HUS pCR rate has posterior mean 0.51 with a 95% HDI from 0.42 to 0.59. The corresponding difference $\hat{\theta}_{\text{RWD} \rightarrow \text{RCT}} - \hat{\theta}_{\text{RCT}}$ is positive in 79% of posterior draws. The estimated posterior distributions of the pCR rates for HUS ($\hat{\theta}_{\text{RWD}}$), NeoSphere ($\hat{\theta}_{\text{RCT}}$), and the adjusted HUS estimate ($\hat{\theta}_{\text{RWD} \rightarrow \text{RCT}}$) are shown in Figure 2.

The MAIC-adjusted pCR estimate had median 0.58 with a bootstrapped 95% confidence interval of 0.46–0.68. Unlike the MRP with raking point estimate, the MAIC point estimate shifted upward relative to the crude HUS rate, although the uncertainty intervals were wide. This result should be interpreted cautiously because the MAIC effective sample size was low: the median ESS was 31 with a 95% bootstrapped confidence interval of 23–38 (Supplementary Figure 1).

The posterior distributions of the adjustment difference $\Delta\hat{\theta}_{\text{adj}}$ and the residual difference $\Delta\hat{\theta}_{\text{res}}$ are shown in Figure 1. The posterior mean of the adjustment difference is -0.03 , with a 95% HDI from -0.13 to 0.08 . The posterior probability that the adjustment difference is negative is approximately 75%, while the probability that its magnitude exceeds 5 percentage points is 39%.

Among the three comparisons, the NeoSphere analysis is the only one in which the residual difference $\Delta\hat{\theta}_{\text{res}}$ is predominantly negative, with 79% of posterior mass below zero. The posterior for $\Delta\hat{\theta}_{\text{res}}$ is displayed in the lower panel of Figure 1. By construction, this residual component represents differences not accounted for by the baseline adjustment and may include contributions




Figure 1: Posterior distributions of the adjustment difference $\Delta\hat{\theta}_{\text{adj}}$ (top) and residual difference $\Delta\hat{\theta}_{\text{res}}$ (bottom) for the NeoSphere comparison.

from treatment differences, unmeasured baseline factors, and other contextual differences. In the NeoSphere case, plausible contributors include differences in disease staging (see Supplementary Table 2) and the shorter NeoSphere protocol of four THP cycles compared with the six cycles typically used in HUS. The magnitude of the residual difference is modest, which should be taken into account when interpreting this posterior distribution: the posterior probability that the adjusted HUS pCR rate exceeds the NeoSphere rate by more than 5 percentage points is 52%.

The Shapley decomposition of $\Delta\hat{\theta}_{\text{adj}}$ reflects the overall uncertainty in the adjustment difference (Figure 3). Most 95% HDIs for variable-level contributions are wide relative to their means. The only contribution showing a relatively consistent negative direction arises from the higher proportion of hormone receptor-positive patients at baseline in NeoSphere (47%) compared with HUS (33%), with mean contribution -0.03 and 95% HDI from -0.06 to 0.00 .




Figure 2: Estimated pCR rates for the NeoSphere B arm, the crude HUS cohort, and the adjusted HUS cohort after baseline covariate alignment.

1.2.2 KRISTINE comparison

The KRISTINE comparison is analyzed using the same framework as for NeoSphere. The posterior mean pCR rate in KRISTINE, $\hat{\theta}_{\text{RCT}}$, is 0.56 with a 95% HDI from 0.50 to 0.63. This does not indicate a clear difference from the HUS pCR rate $\hat{\theta}_{\text{RWD}}$; the posterior probability that $\hat{\theta}_{\text{RCT}} > \hat{\theta}_{\text{RWD}}$ is 0.62.

The baseline-adjustment shift estimated using Bayesian MRP with raking, $\Delta\hat{\theta}_{\text{adj}}$, has mean -0.12 with a 95% HDI from -0.23 to -0.01 (Figure 4). This corresponds to an adjusted HUS pCR rate $\hat{\theta}_{\text{RWD} \rightarrow \text{RCT}}$ with posterior mean 0.42 (95% HDI 0.33–0.51). The MAIC analysis yields a similar adjusted estimate of 0.41 with a bootstrapped 95% confidence interval of 0.30–0.53, indicating




Figure 3: Shapley decomposition of the adjustment difference for the NeoSphere comparison. Bars represent mean Shapley contributions and error bars show 95% HDIs. Mean contributions sum to the total adjustment difference.

broad agreement between the two adjustment approaches despite their different modeling assumptions. Posterior distributions for $\hat{\theta}_{\text{RCT}}$, $\hat{\theta}_{\text{RWD}}$, and $\hat{\theta}_{\text{RWD} \rightarrow \text{RCT}}$ are shown in Figure 5.

Limited overlap between cohorts contributes to uncertainty in both adjustment approaches. The exclusion of nodal-positive patients reduces the pool of comparable cases, requiring extrapolation in the raking model and producing highly variable weights in MAIC. In the MAIC diagnostics, the median effective sample size was 34 with a 95% bootstrapped confidence interval of 27–40 (Supplementary Figure 2), indicating substantial loss of effective information after reweighting.

The residual difference $\Delta\hat{\theta}_{\text{res}}$ (Figure 4) has mean 0.13 with a 95% HDI from 0.02 to 0.24. Despite substantial uncertainty, 94% of posterior mass supports a difference exceeding 5 percentage points between $\hat{\theta}_{\text{RCT}}$ and $\hat{\theta}_{\text{RWD} \rightarrow \text{RCT}}$. This suggests that factors not captured by baseline adjustment account for a sizable residual difference after baseline alignment.

The Shapley decomposition of $\Delta\hat{\theta}_{\text{adj}}$ indicates that the higher prevalence of hormone receptor-positive patients in KRISTINE (62%; Supplementary Table 3) relative to HUS (33%) corresponds to a mean -0.06 contribution to the adjustment difference (95% HDI -0.09 to -0.03). This estimate likely understates the contribution of hormone receptor status, because patients excluded due to nodal-positive biopsy have a lower hormone receptor-positive proportion than those retained (29% compared to 37%). As a result, part of the




Figure 4: Top: Adjustment difference $\Delta\hat{\theta}_{\text{adj}}$ and Bottom: residual difference $\Delta\hat{\theta}_{\text{res}}$ for KRISTINE.

hormone-receptor contribution may be redistributed to the nodal-biopsy exclusion variable. The Shapley value for nodal-biopsy exclusion has mean -0.04 (95% HDI -0.09 to 0.01), a direction that is difficult to interpret clinically and likely reflects this redistribution of attribution.




Figure 5: Estimated pCR rates for the KRISTINE B arm, the crude HUS cohort, and the adjusted HUS cohort after baseline covariate alignment.

1.2.3 HELEN-006 comparison

The unadjusted pCR rate $\hat{\theta}_{\text{RCT}}$ for the HELEN-006 arm B cohort is 0.57 with a 95% HDI from 0.52 to 0.63, comparable to the TCHP arm in the KRISTINE trial. The posterior probability that $\hat{\theta}_{\text{RCT}} > \hat{\theta}_{\text{RWD}}$ is 0.77, indicating moderate evidence for a higher pCR rate in HELEN-006 relative to HUS. However, the probability that this difference exceeds 5 percentage points is 0.31, suggesting limited support for a clinically large difference.

HELEN-006 provides the most detailed reporting of baseline characteristics (Supplementary Table 4), enabling adjustment for a broader set of variables than in the other comparisons. Posterior distributions for $\hat{\theta}_{\text{RCT}}$, $\hat{\theta}_{\text{RWD}}$, and $\hat{\theta}_{\text{RWD} \rightarrow \text{RCT}}$ are shown in Figure 8, and the corresponding adjustment and residual differences are presented in Figure 7.

The adjustment difference $\Delta\hat{\theta}_{\text{adj}}$ has mean -0.07 with a 95% HDI from -0.17 to 0.03 . This corresponds to an adjusted HUS pCR rate $\hat{\theta}_{\text{RWD} \rightarrow \text{RCT}}$ with mean 0.48 (95% HDI 0.40–0.56). The MAIC estimate is similar at 0.49 (95% CI 0.40–0.58), providing convergent evidence across methods. The posterior probability that $\Delta\hat{\theta}_{\text{adj}}$ is positive is 0.10, indicating that a positive adjustment difference is unlikely. The residual difference $\Delta\hat{\theta}_{\text{res}}$ has mean 0.10 with a 95% HDI from 0.03 to 0.19.




Figure 6: Shapley decomposition of the adjustment difference for the KRISTINE comparison. Bars represent mean Shapley contributions and error bars show 95% HDIs. Mean contributions sum to the total adjustment difference.

The Shapley decomposition (Figure 9) shows that the higher prevalence of hormone receptor-positive patients in HELEN-006 (57% vs. 33% in HUS) is the dominant contributor to the adjustment difference, with mean contribution -0.09 (95% HDI -0.13 to -0.05), accounting for a substantial portion of the total absolute Shapley contributions. Pre-treatment histology contributes positively (mean 0.04 , 95% HDI 0.02 – 0.06), largely reflecting the higher proportion of non-ductal tumors in HUS. Differences in HER2 IHC/ISH distribution also contribute modestly; the higher proportion of IHC2+/ISH+ cases in HELEN-006 (31% vs. 27% in HUS) corresponds to a Shapley contribution of -0.03 (95% HDI -0.05 to -0.01).

1.3 Sensitivity of results to NAT discontinuation

One potential contributor to the residual differences $\Delta\hat{\theta}_{\text{res}}$ in the KRISTINE and HELEN-006 comparisons is a higher proportion of patients in the HUS cohort who did not complete the planned NAT. To assess this, a binary treatment-discontinuation variable was added to the Bayesian MRP with raking model. The marginal distribution of this variable was aligned to the discontinuation proportions reported in the RCTs (NeoSphere B: 5/107 Gianni et al. [2012]; KRISTINE B: 18/221 Hurvitz et al. [2018]; HELEN-006 B: 23/337) and its influence on the estimates was evaluated.

Including treatment discontinuation in the adjustment set had minimal im-




Figure 7: Top: Adjustment difference $\Delta\hat{\theta}_{\text{adj}}$ and Bottom: residual difference $\Delta\hat{\theta}_{\text{res}}$ for HELEN-006.




Figure 8: Estimated pCR rates for the HELEN-006 B arm, the crude HUS cohort, and the adjusted HUS cohort after baseline covariate alignment.




Figure 9: Shapley decomposition of the adjustment difference for the HELEN-006 comparison. Bars represent mean Shapley contributions and error bars show 95% HDIs. Mean contributions sum to the total adjustment difference.

Characteristic	N	Summary
Median age	337	50 (IQR 44-56)
T stage: T1 to T2	281	83%
T stage: T3 to T4	56	17%
Nodal status: Positive	247	73%
Nodal status: Negative	90	27%
Bilateral: False	337	100%
Tumor grade: 1	1	< 1%
Tumor grade: 2	142	42%
Tumor grade: 3	194	58%
Histological type: Ductal	317	94%
Histological type: Lobular	4	1%
Histological type: Other	16	5%
ER or PR positive	193	57%
ER and PR negative	144	43%
IHC: 2+ and ISH	104	31%
IHC: 3+	233	69%
Ki-67 index ≤ 30 %	75	22%
Ki-67 index > 30 %	262	78%
Disease stage: II	212	63%
Disease stage: III	125	37%

Table 2: Baseline characteristics for the HELEN-006 arm B (TCHP) population, restricted to variables used for adjustment and staging.

impact on the estimated residual differences. Across NeoSphere, KRISTINE, and HELEN-006, posterior means of $\Delta\hat{\theta}_{\text{res}}$ changed by less than 1 percentage point. Consistent with this, the outcome-model coefficients for treatment discontinuation in the KRISTINE and HELEN-006 analyses indicated only weak associations with pCR (Subsection 1.1). Although the discontinuation threshold differed in NeoSphere and was associated with a more negative coefficient, incorporating this variable did not materially alter the posterior distributions of $\Delta\hat{\theta}_{\text{adj}}$ or $\Delta\hat{\theta}_{\text{res}}$. These tests suggest that treatment completion is unlikely to be a major explanation for the observed residual differences.

References

- Fabio Canino, Monica Barbolini, Ugo De Giorgi, Tommaso Fontana, Valeria Gaspari, Caterina Gianni, Lorenzo Gianni, Antonio Maestri, Santino Minichillo, Luca Moschetti, Antonella Mura, Stefania Vittoria Luisa Nicoletti, Claudia Omarini, Rachele Pagani, Samanta Sarti, Angela Toss, Claudio Zambagni, Riccardo Cuoghi Costantini, Federica Caggia, Giuseppina Antonelli, Federica Baglio, Lorenzo Belluzzi, Giulio Martinelli, Salvatore Natalizio, Ornella Ponzoni, Massimo Dominici, and Federico Piacentini. Safety and efficacy analysis of neoadjuvant pertuzumab, trastuzumab and standard chemotherapy for HER2-positive early breast cancer: real-world data from NeoPower study. *BMC Cancer*, 24(1):735, 2024. ISSN 1471-2407. doi: 10.1186/s12885-024-12506-0.
- Meirong Zhou, Shouman Wang, Nengbin Wan, Songlin Yuan, Xiongqiang Hu, Wei Zhou, Bohua Qing, Mingwen Liu, Weihua Sun, Peizhi Fan, Jianguo Wang, Hong Cao, Haifan Xu, Bin Dai, Peizhi Tang, Liyuan Qian, Xi Zhao, Jun Xiao, Huaiying Zhou, Jinhui Hu, Li Ding, Domenico Tripodi, Nicholas Zdenkowski, Thomas J. O’Keefe, Alejandro Martin Sanchez, Li Chen, Ping Zhang, and Feng Xu. Efficacy and safety of neoadjuvant pertuzumab plus trastuzumab in combination with chemotherapy regimen in Chinese patients with HER2-positive early breast cancer: a real-world retrospective multicenter cohort study. *Annals of Translational Medicine*, 10:1387, December 2022.
- Ji-Yeon Kim, Seok Jin Nam, Jeong Eon Lee, Jonghan Yu, Byung Joo Chae, Se Kyung Lee, Jai Min Ryu, Jin Seok Ahn, Young-Hyuck Im, Seok Won Kim, and Yeon Hee Park. Real world evidence of neoadjuvant docetaxel/carboplatin/trastuzumab/pertuzumab (TCHP) in patients with HER2-positive early or locally advanced breast cancer: A single-institutional clinical experience. *Cancer Research and Treatment*, 54(4):1091–1098, January 2022. ISSN 1598-2998. doi: 10.4143/crt.2021.901.
- Alejandro Falcón González, Josefina Cruz Jurado, Elisenda Llabrés Valenti, Rocío Urbano Cubero, Maria Carmen Álamo de la Gala, María Antonia Martínez Guisado, Rocío Álvarez Ambite, Carlos José Rodríguez González,

- Marta Américo Góngora, Lourdes Rodríguez Pérez, Pilar López Álvarez, Pedro Sánchez Rovira, Encarnación González Flores, Fernando Henao Carrasco, Juan Bayo Calero, María Valero Arbizu, Alicia Quílez Cutillas, Javier Salvador Boffil, Eloísa Rubio Pérez, and Manuel Ruiz-Borrego. Real-world experience with pertuzumab and trastuzumab combined with chemotherapy in neoadjuvant treatment for patients with early-stage HER2-positive breast cancer: the NEOPERSUR study. *Clinical and Translational Oncology*, 26(9): 2217–2226, 2024. ISSN 1699-3055. doi: 10.1007/s12094-024-03440-5.
- Benjamin James Hall, Ajay Ashok Bhojwani, Helen Wong, Andrea Law, Helen Flint, Eliyaz Ahmed, Helen Innes, Joanne Cliff, Zaf Malik, Julie Elizabeth O’Hagan, Allison Hall, Rajaram Sripadam, Shaun Tolan, Zulfiqar Ali, Clare Hart, Douglas Errington, Farida Alam, Rosa Giuliani, Shaveta Mehta, Sheena Khanduri, Nicky Thorp, Richard Jackson, Silvia Cicconi, and Carlo Palmieri. Neoadjuvant trastuzumab and pertuzumab for early HER2-positive breast cancer: A real-world experience. *The Breast Journal*, 2022(1):7146172, January 2022. doi: 10.1155/2022/7146172.
- Luca Gianni, Tadeusz Pienkowski, Young-Hyuck Im, Laslo Roman, Ling-Ming Tseng, Mei-Ching Liu, Ana Lluch, Elżbieta Staroslawska, Juan de la Haba-Rodriguez, Seock-Ah Im, Jose Luiz Pedrini, Brigitte Poirier, Paolo Morandi, Vladimir Semiglazov, Vichien Srimuninnimit, Giulia Bianchi, Tania Szado, Jayantha Ratnayake, Graham Ross, and Pinuccia Valagussa. Efficacy and safety of neoadjuvant pertuzumab and trastuzumab in women with locally advanced, inflammatory, or early HER2-positive breast cancer (NeoSphere): a randomised multicentre, open-label, phase 2 trial. *The Lancet Oncology*, 13(1):25–32, 2012. ISSN 1470-2045. doi: 10.1016/S1470-2045(11)70336-9.
- Sara A Hurvitz, Miguel Martin, W Fraser Symmans, Kyung Hae Jung, Chiun-Sheng Huang, Alastair M Thompson, Nadia Harbeck, Vicente Valero, Daniil Stroyakovskiy, Hans Wildiers, Mario Campone, Jean-François Boileau, Matthias W Beckmann, Karen Afenjar, Rodrigo Fresco, Hans-Joachim Helms, Jin Xu, Yvonne G Lin, Joseph Sparano, and Dennis Slamon. Neoadjuvant trastuzumab, pertuzumab, and chemotherapy versus trastuzumab emtansine plus pertuzumab in patients with HER2-positive breast cancer (KRISTINE): a randomised, open-label, multicentre, phase 3 trial. *The Lancet Oncology*, 19(1):115–126, 2018. ISSN 1470-2045. doi: 10.1016/S1470-2045(17)30716-7.