

# A MAGIC DETERMINANT FORMULA FOR SYMMETRIC POLYNOMIALS OF EIGENVALUES

Jules Jacobs

$$\sum_i p_i \lambda_1^{i_1} \lambda_2^{i_2} \cdots \lambda_n^{i_n} = \sum_i p_i \det(A_1^{i_1} | A_2^{i_2} | \cdots | A_n^{i_n})$$

## TRACE AND DETERMINANT

$$A = \begin{pmatrix} A_{11} & A_{12} & A_{13} \\ A_{21} & A_{22} & A_{23} \\ A_{31} & A_{32} & A_{33} \end{pmatrix} \quad (\text{e.g. integer matrix})$$

## TRACE AND DETERMINANT

$$A = \begin{pmatrix} A_{11} & A_{12} & A_{13} \\ A_{21} & A_{22} & A_{23} \\ A_{31} & A_{32} & A_{33} \end{pmatrix} \quad (\text{e.g. integer matrix})$$

$$\text{tr}(A) = \lambda_1 + \lambda_2 + \lambda_3 = A_{11} + A_{22} + A_{33}$$

$$\begin{aligned} \det(A) = \lambda_1 \lambda_2 \lambda_3 = & A_{11}A_{22}A_{33} - A_{11}A_{32}A_{23} - A_{12}A_{21}A_{33} + \\ & A_{12}A_{23}A_{31} + A_{13}A_{21}A_{32} - A_{13}A_{22}A_{31} \end{aligned}$$

## TRACE AND DETERMINANT

$$A = \begin{pmatrix} A_{11} & A_{12} & A_{13} \\ A_{21} & A_{22} & A_{23} \\ A_{31} & A_{32} & A_{33} \end{pmatrix} \quad (\text{e.g. integer matrix})$$

$$\text{tr}(A) = \lambda_1 + \lambda_2 + \lambda_3 = A_{11} + A_{22} + A_{33}$$

$$\begin{aligned} \det(A) = \lambda_1 \lambda_2 \lambda_3 = & A_{11}A_{22}A_{33} - A_{11}A_{32}A_{23} - A_{12}A_{21}A_{33} + \\ & A_{12}A_{23}A_{31} + A_{13}A_{21}A_{32} - A_{13}A_{22}A_{31} \end{aligned}$$

Polynomial of irrational complex numbers = polynomial of integer entries!

## TRACE AND DETERMINANT

$$A = \begin{pmatrix} A_{11} & A_{12} & A_{13} \\ A_{21} & A_{22} & A_{23} \\ A_{31} & A_{32} & A_{33} \end{pmatrix} \quad (\text{e.g. integer matrix})$$

$$\text{tr}(A) = \lambda_1 + \lambda_2 + \lambda_3 = A_{11} + A_{22} + A_{33}$$

$$\begin{aligned} \det(A) = \lambda_1 \lambda_2 \lambda_3 = & A_{11}A_{22}A_{33} - A_{11}A_{32}A_{23} - A_{12}A_{21}A_{33} + \\ & A_{12}A_{23}A_{31} + A_{13}A_{21}A_{32} - A_{13}A_{22}A_{31} \end{aligned}$$

Polynomial of irrational complex numbers = polynomial of integer entries!

Q: Which other polynomials  $p(\lambda_1, \lambda_2, \lambda_3)$  can we compute exactly?

## TRACE AND DETERMINANT

$$A = \begin{pmatrix} A_{11} & A_{12} & A_{13} \\ A_{21} & A_{22} & A_{23} \\ A_{31} & A_{32} & A_{33} \end{pmatrix} \quad (\text{e.g. integer matrix})$$

$$\text{tr}(A) = \lambda_1 + \lambda_2 + \lambda_3 = A_{11} + A_{22} + A_{33}$$

$$\begin{aligned} \det(A) = \lambda_1 \lambda_2 \lambda_3 = & A_{11}A_{22}A_{33} - A_{11}A_{32}A_{23} - A_{12}A_{21}A_{33} + \\ & A_{12}A_{23}A_{31} + A_{13}A_{21}A_{32} - A_{13}A_{22}A_{31} \end{aligned}$$

Polynomial of irrational complex numbers = polynomial of integer entries!

Q: Which other polynomials  $p(\lambda_1, \lambda_2, \lambda_3)$  can we compute exactly?

A: All symmetric ones:

$$p(\lambda_1, \lambda_2, \lambda_3) = p(\lambda_2, \lambda_1, \lambda_3) = \cdots = p(\lambda_3, \lambda_2, \lambda_1)$$

## THE MAGIC METHOD

$$p(\lambda_1, \lambda_2, \lambda_3) = \lambda_1 \lambda_2^4 \lambda_3^4 + \lambda_1^4 \lambda_2 \lambda_3^4 + \lambda_1^4 \lambda_2^4 \lambda_3$$

## THE MAGIC METHOD

$$\begin{aligned} p(\lambda_1, \lambda_2, \lambda_3) &= \lambda_1 \lambda_2^4 \lambda_3^4 + \lambda_1^4 \lambda_2 \lambda_3^4 + \lambda_1^4 \lambda_2^4 \lambda_3 \\ &= \det(A_1 | A_2^4 | A_3^4) + \det(A_1^4 | A_2 | A_3^4) + \det(A_1^4 | A_2^4 | A_3) \end{aligned}$$



## THE MAGIC METHOD

$$\begin{aligned} p(\lambda_1, \lambda_2, \lambda_3) &= \lambda_1 \lambda_2^4 \lambda_3^4 + \lambda_1^4 \lambda_2 \lambda_3^4 + \lambda_1^4 \lambda_2^4 \lambda_3 \\ &= \det(A_1 | A_2^4 | A_3^4) + \det(A_1^4 | A_2 | A_3^4) + \det(A_1^4 | A_2^4 | A_3) \end{aligned}$$

$A_2^4$  = the second column of  $A^4$

## THE MAGIC METHOD

$$\begin{aligned} p(\lambda_1, \lambda_2, \lambda_3) &= \lambda_1 \lambda_2^4 \lambda_3^4 + \lambda_1^4 \lambda_2 \lambda_3^4 + \lambda_1^4 \lambda_2^4 \lambda_3 \\ &= \det(A_1 | A_2^4 | A_3^4) + \det(A_1^4 | A_2 | A_3^4) + \det(A_1^4 | A_2^4 | A_3) \end{aligned}$$

$A_2^4$  = the second column of  $A^4$

$$\cdots + a \lambda_1^{i_1} \cdot \lambda_2^{i_2} \cdots \lambda_n^{i_n} + \cdots$$

$\Downarrow$

$$\cdots + a \det(A_1^{i_1} | A_2^{i_2} | \cdots | A_n^{i_n}) + \cdots$$

# THE TRACE

$$\lambda_1 + \lambda_2 + \lambda_3$$

## THE TRACE

$$\lambda_1 + \lambda_2 + \lambda_3 = \lambda_1 \lambda_2^0 \lambda_3^0 + \lambda_1^0 \lambda_2 \lambda_3^0 + \lambda_1^0 \lambda_2^0 \lambda_3$$

## THE TRACE

$$\begin{aligned}\lambda_1 + \lambda_2 + \lambda_3 &= \lambda_1 \lambda_2^0 \lambda_3^0 + \lambda_1^0 \lambda_2 \lambda_3^0 + \lambda_1^0 \lambda_2^0 \lambda_3 \\ &= \det(A_1|A_2^0|A_3^0) + \det(A_1^0|A_2|A_3^0) + \det(A_1^0|A_2^0|A_3)\end{aligned}$$

## THE TRACE

$$\begin{aligned}\lambda_1 + \lambda_2 + \lambda_3 &= \lambda_1 \lambda_2^0 \lambda_3^0 + \lambda_1^0 \lambda_2 \lambda_3^0 + \lambda_1^0 \lambda_2^0 \lambda_3 \\ &= \det(A_1 | A_2^0 | A_3^0) + \det(A_1^0 | A_2 | A_3^0) + \det(A_1^0 | A_2^0 | A_3) \\ &= \det \begin{pmatrix} A_{11} & 0 & 0 \\ A_{12} & 1 & 0 \\ A_{13} & 0 & 1 \end{pmatrix} + \det \begin{pmatrix} 1 & A_{21} & 0 \\ 0 & A_{22} & 0 \\ 0 & A_{23} & 1 \end{pmatrix} + \det \begin{pmatrix} 1 & 0 & A_{31} \\ 0 & 1 & A_{32} \\ 0 & 0 & A_{33} \end{pmatrix}\end{aligned}$$

## THE TRACE

$$\begin{aligned}\lambda_1 + \lambda_2 + \lambda_3 &= \lambda_1 \lambda_2^0 \lambda_3^0 + \lambda_1^0 \lambda_2 \lambda_3^0 + \lambda_1^0 \lambda_2^0 \lambda_3 \\&= \det(A_1 | A_2^0 | A_3^0) + \det(A_1^0 | A_2 | A_3^0) + \det(A_1^0 | A_2^0 | A_3) \\&= \det \begin{pmatrix} A_{11} & 0 & 0 \\ A_{12} & 1 & 0 \\ A_{13} & 0 & 1 \end{pmatrix} + \det \begin{pmatrix} 1 & A_{21} & 0 \\ 0 & A_{22} & 0 \\ 0 & A_{23} & 1 \end{pmatrix} + \det \begin{pmatrix} 1 & 0 & A_{31} \\ 0 & 1 & A_{32} \\ 0 & 0 & A_{33} \end{pmatrix} \\&= A_{11} + A_{22} + A_{33}\end{aligned}$$