

HW_2: Ch 6: Ex 32, 34, Ch 7: Ex 2, 6, 16, and 20

Julia Andrus

Chp 6: 32, 34

32a) Let X_1, \dots, X_n be a random sample from a uniform distribution on $[0, \theta]$. Then the mle of θ is $\hat{\theta} = Y = \max(X_i)$. Use the fact that $Y \leq y$ iff each $X_i \leq y$ to the derivative the cdf of Y . Then show that the pdf of $Y = \max(X_i)$ is:

$$f_Y(y) = \begin{cases} \frac{ny^{n-1}}{\theta^n} & 0 \leq y \leq \theta \\ 0 & \text{otherwise} \end{cases}$$

cdf:

$$F_Y(y) = P(Y \leq y) = P(\max(X_i) \leq y)$$

all the X_i are small, if max is small:

$$= P(X_1 \leq y * \dots * X_n \leq y)$$

they are independent:

$$= P(X_1 \leq y) * \dots * (X_n \leq y)$$

cdf of uniform distribution on $[0, \theta]$:

$$= \left(\frac{y}{\theta}\right)^n, 0 \leq y \leq \theta$$

Now taking the derivative:

$$\frac{d}{df_Y} = \begin{cases} \frac{ny^{n-1}}{\theta^n} & 0 \leq y \leq \theta \\ 0 & \text{otherwise} \end{cases}$$

32b) Use the result of part (a) to show that mle is biased but that $(n + 1)\max(X_i)/n$ is unbiased.

For the estimator $(E(y) = \theta)$ to be unbiased, then:

$$\begin{aligned} E(Y) &= \int_0^\theta y * \frac{ny^n - 1}{\theta^n} dy \\ &= \frac{n}{\theta^n} \frac{y^n + 1}{n + 1} \Big|_0^\theta \\ &= \frac{n}{n + 1} \theta \end{aligned}$$

As you can see that $E(Y) \neq \theta$, and \therefore the estimator is biased.

To show that $(n + 1)\max(X_i)/n$ is unbiased:

Let $\tilde{Y} = \frac{n+1}{n}$, then

$$E(\tilde{Y}) = E \frac{(n + 1)}{n}$$

Using result from (a):

$$\begin{aligned} &= E(Y) \frac{n + 1}{n} \\ &= \theta \frac{n}{n + 1} \frac{n + 1}{n} \end{aligned}$$

Cancel:

$$= \theta$$

This shows that \tilde{Y} is **unbiased**.

34) The mean squared error of an estimator $\hat{\theta}$ is $MSE(\hat{\theta}) = E(\hat{\theta} - \theta)^2$. Consider $\hat{\sigma}^2 = KS^2$, where S^2 = sample variance. If $\hat{\theta}$ is unbiased, then $MSE(\hat{\theta}) = V(\hat{\theta}) + (bias)^2$. What value of K minimizes the mean squared error of this estimator when the population distribution is normal?

First we have to find the mean squared error of $\hat{\sigma}^2 = KS^2$. Since S^2 is our estimator, then:

$$MSE(\hat{\sigma}^2) = V(\hat{\sigma}^2) + (bias)(\sigma^2)$$

$$Bias(\sigma^2) = E(\hat{\sigma}^2 - \sigma^2) = E(KS^2) - \sigma^2 = K\sigma^2 - \sigma^2 = \sigma^2(k - 1).$$

Let's find the variance given that S^2 is unbiased of σ^2 :

$$\begin{aligned} V(\hat{\sigma}^2) &= V(KS^2) \\ &= K^2V(S^2) = K^2(E((S^2)^2) - (E(S^2))^2) \\ &= K^2\left(\frac{n+1}{n-1}\sigma^4 - (\sigma^2)^2\right) \\ &= K^2\sigma^4\left(\frac{n+1}{n-1} - 1\right) \end{aligned}$$

Now to find the value of K that minimize the MSE, we need to take the derivative of the MSE:

$$\begin{aligned} MSE(\sigma^2) &= K^2\sigma^4\left(\frac{n+1}{n-1} - 1\right) - (\sigma^2(k - 1))^2 \\ &= K^2\sigma^4\left(\frac{n+1}{n-1} - 1\right) - \sigma^4(k - 1)^2 \end{aligned}$$

Now doing the derivative:

$$\begin{aligned} \frac{d}{dK} MSE(\sigma^2) &= \frac{d}{dK} (K^2\sigma^4\left(\frac{n+1}{n-1} - 1\right) - \sigma^4(k - 1)^2) \\ &= 2K\sigma^4\left(\frac{n+1}{n-1} - 1\right) - 2\sigma^4(k - 1) \end{aligned}$$

Now equal it to 0 so we can find the minimizer of the MSE:

$$2K\sigma^4\left(\frac{n+1}{n-1} - 1\right) - 2\sigma^4(k - 1) = 0$$

$$2K\sigma^4\left(\frac{n+1}{n-1} - 1\right) = 2\sigma^4(k - 1)$$

$$K\left(\frac{n+1}{n-1} - 1\right) = (k - 1)$$

.

$$K = \frac{n-1}{n-3}$$

Answer: $K = \frac{n-1}{n-3}$

Ch 7: Ex 2, 6, 16, and 20

2a) Each of the following is a confidence interval for μ =true average (i.e., population mean) resonance frequency (Hz) for all tennis rackets of a certain type. (114.4 , 115.6) and (114.1 , 115.9). What is the value of the sample mean resonance frequency?

To find the sample mean, we just

$$\bar{X} = \frac{x_1 + x_2 \dots x_n}{n}$$

So for (114.4 , 115.6):

$$\bar{X} = \frac{114.4 + 115.6}{2} = 115$$

For (114.1 , 115.9):

$$\bar{X} = \frac{114.1 + 115.9}{2} = 115$$

Answer: Mean for both intervals is 115

2b) Both intervals were calculated from the same sample data. The Confidence level for one of these intervals is 90% and the other has 99%. Which has 90%?

In class we talked about how the length of confidence interval depends on couple of points. One of the points mentioned about the sample size (n): the larger the value, the shorter the length of confidence interval. So for 90% C.I, the interval would have to be narrower. If we have a wider interval, then our C.I. would be higher. With that being said, I believe that (114.4 , 115.6) has 90% C.I.

6a) Mild-reinforcing bar is known to be normally distributed with $\sigma = 100$. The composition of bars has been slightly modified, but does not affect the normality or value of σ . If $n=25$, $\bar{X} = 8439\text{lb}$, compute a 90% CI for the true average yield point of the modified bar.

First we have to find what the area for a 90% interval:

$$A = \frac{1 + .90}{2} = .95$$

Now using the appendix in the back of the book to find z-score:

$$.95 = 1.64$$

$$z = 1.64$$

Now we have to use this formula to compute the intervals: $\bar{X} \pm z(\frac{\sigma}{n})$

$$n=25$$

$$\bar{X} = 8439\text{lb}$$

$$\sigma = 100$$

$$z=1.64$$

So,

$$\begin{aligned} & \bar{X} \pm z\left(\frac{\sigma}{n}\right) \\ & 8439 \pm 1.64\left(\frac{100}{25}\right) \\ & \left(8439 - 1.64\left(\frac{100}{25}\right), 8439 + 1.64\left(\frac{100}{25}\right)\right) \\ & (8432.44, 8445.56) \end{aligned}$$

Answer: (8432.44, 8445.56)

6b) How would you modify the interval in part (a) to obtain a confidence level of 92%?

The interval would have to be a bit wider to get a higher CI. So, same process as we did in part (a):

First we have to find what the area for a 90% interval:

$$A = \frac{1 + .92}{2} = .96$$

Now using the appendix in the back of the book to find z-score:


$$.96 = 1.75$$

$$z = 1.75$$

Once changing z that correspond with the percentage CI, then I can find out what my interval is by going through the same process I did in part (a). Note that the interval will be a little wider because our higher CI.

16a) Construct a boxplot of the data and comment on interesting features.

I have constructed a boxplot in R:



This boxplot is not symmetric so that means that is it not normally distributed.

16b) Calculate and interpret a 95% CI for true average breakdown voltage μ . Does it appear that μ has been precisely estimated? Explain.

We find some information to use the margin of error formula: margin of error = $E = z_{\frac{\alpha}{2}} \times \frac{s}{\sqrt{(n)}}$:

$$\bar{X} = \frac{x_1 + x_2 + \dots + x_n}{n} = 54.70833$$

$$s^2 = \sigma^2 = \sum(x^2) - (\bar{X})^2 = 27.35993$$

$$s = \sigma = \sqrt{(\sigma^2)} = 5.230672$$

$$n=48$$

$$z=1.96$$

So now that we have all of our info, we can solve:

$$\begin{aligned} & z_{\frac{\alpha}{2}} \times \frac{s}{\sqrt{(n)}} \\ & 1.96 \times \frac{5.230672}{\sqrt{(48)}} \\ & E = 1.4798 \end{aligned}$$

Now we can use this formula to get our intervals: $\bar{X} \pm E$

$$\bar{X} \pm E$$

$$(\bar{X} - E, \bar{X} + E)$$

$$(54.70833 - 1.4798, 54.70833 + 1.4798)$$

$$(53.2285, 56.1881)$$

Answer: (53.2285, 56.1881)

16c) all values of breakdown voltage are between 40 and 70. What sample size would be appropriate for the 95% CI to have a width of 2 kV (so that μ is estimated to within 1 kV with 95% confidence)?

For this problem, we have to use this formula: $n = (\frac{z_{\frac{\alpha}{2}} \sigma}{E})^2$

We first need to find the population deviation of 40 to 70, which can be estimated by 1/4: $\sigma = \frac{70-40}{4} = 7.5$

Now we can use the formula to find our n :

$$n = (\frac{z_{\frac{\alpha}{2}} \sigma}{E})^2$$

$$n = (\frac{1.96 \times 7.5}{1.4798})^2$$

$$n = 217$$

Answer: $n=217$

20a) Calculate and interpret a confidence interval at the 99% confidence level

We are given:

$$n = 2343$$

$$\hat{p} = 53\% = .53$$

$$z = 2.575$$

The margin error formula is: $E = z_{\frac{\alpha}{2}} \left(\sqrt{\frac{\hat{p}(1-\hat{p})}{n}} \right)$

$$E = 2.575 \left(\sqrt{\frac{.53(1-.53)}{2343}} \right)$$

$$E = 0.0266$$

Now to get the intervals: $\hat{p} \pm E$

$$.53 - .0266 = .5034$$

$$.53 + .0266 = 0.5566$$

Answer: (0.5034 , 0.5566)

20b) What sample size would be required for the width of a 99% CI to be at most .05 irrespective of the value of \hat{p} ?

Since we are interested in finding the sample size of 99% CI to be at most .05 irrespective of the value of \hat{p} , then we use this formula: $n = \frac{(z_{\frac{\alpha}{2}})^2 \times \hat{p}(1-\hat{p})}{E^2}$

$$z=2.575$$

$$E=0.05$$

$$\hat{p} = .53$$

So,

$$n = \frac{2.575^2 \times .53(1-.53)}{.053^2}$$

$$n = 661$$

Answer: 661 (rounded to the nearest whole number)