

# EMIC-Induced Precipitation as Measured by ELFIN: Source Regions, Impacts, and Energy Dependence

Julia Claxton (she/her), Robert Marshall – University of Colorado Boulder; julia.claxton@colorado.edu

## Motivation

Electromagnetic Ion-Cyclotron (EMIC) waves are thought to be a primary driver of relativistic electron loss from the outer radiation belt (Ni 2015, Thorne 1971). The atmospheric impact of this high-energy precipitation is poorly-characterized due to a lack of measurements resolving the loss cone. Further, the minimum electron energy required for scattering by EMIC waves remains unknown, as the quasi-linear theory traditionally used to analyze wave-particle interactions underestimates scattering of subrelativistic electrons (Cappanolo 2023). This poster aims to move toward an understanding of global EIP fluxes and the energy-domain response of electrons to EMIC waves by using recent data that can resolve the interior of the loss cone at a variety of energies.

## Data Source: ELFIN

Electron Fields and Losses Investigation (Angelopoulos 2020)

- Twin CubeSats in polar LEO
- 3 years of data (2019-2022) over 2 satellites
- $\sim 22.5^\circ$  angular resolution
- LEO  $\Rightarrow$  Resolvable loss cone ( $\alpha_{LC} \approx 67^\circ$ )




Fig. 1: ELFIN data showing EIP at ~07:05:00 UTC

## Methods

Candidate EIP events detected using a scoring algorithm. Process: Look at  $J_{prec}/J_{trap}$  heatmap (see Fig. 1). For high- ( $> 500$  keV) and low- ( $< 500$  keV) energy channels:

1. Remove pixels with low precipitation ( $J_{prec}/J_{trap} < 10^{-75}$ )
2. Remove pixels with  $< 3$  nonzero neighbors
3. Vertically sum remaining  $J_{prec}/J_{trap}$  values over given energy range

This produces a precipitation score for low and high energy channels. The low energy score is then set to its highest value in the surrounding 4 timesteps on either side to reject detections due to the electron isotropy boundary. A candidate EIP event is returned when:

1. **high-energy score**  $> .75$   
Loss cone must be filled at high energies.
2. **high-energy score**  $> \text{low-e score} - 1.5$   
High energy precipitation must be comparable or larger than low energy precipitation.
3. **low-energy score**  $< 4$   
Reject events with very strong low-energy precipitation, e.g. the isotropized plasma sheet.




Fig. 2: Example EIP detections.

## Key Points:

- An in-progress EIP detection algorithm is used to create a set of 237 likely EIP events
- These events primarily originate from dusk to midnight at  $L \approx 5-10$ , in line with previous results
- EIP precipitation is strongest at  $\sim$ MeV energies, extending down to 100s of keV in some cases

## EIP Distributions

Origin of EMIC-Driven Precipitation (N = 237)




Fig. 3: EIP detection rate shown in MLT/L and a polar projection of same distribution. This distribution generally agrees with the distribution of H-band EMIC waves found in Min et al. 2012 and Usanova et al. 2012.

EMIC-Driven Atmospheric Energy Input




Fig. 4: Global atmospheric energy input due to EIP events analyzed, normalized to the time ELFIN spent in each MLT/latitude bin.

## Future Work

- Improve algorithm to accept EIP with low-energy precipitation
- Reduce need for human verification of EIP detection
- Expand dataset to capture majority of EIP measured by ELFIN
- Quantify atmospheric backscatter of EIP

## Open Science

All code used in this analysis is available at [github.com/julia-claxton/2024\\_GEM\\_workshop](https://github.com/julia-claxton/2024_GEM_workshop). ELFIN data are freely available at [data.elfin.ucla.edu](https://data.elfin.ucla.edu).



## EIP Energy Profile

EIP Pitch Angle Distributions




Fig. 4: Average flux of EIP events, normalized to perpendicular flux, in pitch angle-energy space. This demonstrates a filling of the loss cone at high energies, with filling intensity decreasing with energy down to  $\sim 200$  keV. Atmospheric backscatter can be seen at  $\sim 400$  keV in the anti loss cone.

## Precipitating Flux Distribution by Energy Channel




Fig. 6: Quantiles of  $J_{prec}/J_{trap}$  distribution for each of ELFIN's energy channels. Precipitating flux increases with increasing energy, beginning in the 100s of keV range.

## Findings

- EIP events can be detected in particle data by looking for patterns of increasing  $J_{prec}/J_{perp}$  with increasing energy
- EIP identification does not require conjunctions with other instruments
- EMIC-induced precipitation originates from dusk to nightside at  $L \approx 5-10$
- In this set of events, EMIC scattering begins at  $\sim 400$  keV

## Limitations

- EIP selection punishes low-energy precipitation, potentially excluding EIP events with sub-MeV effects
- Varying loss cone coverage requires extrapolation, leading to some inaccuracy in precipitating flux estimates

## References

Cappanolo+ 2023, <https://doi.org/10.1029/2023GL103519>  
Ni+ 2015, <https://doi.org/10.1002/2015JA021466>  
Thorne+ 1971, <https://doi.org/10.1029/JA076i019p04446>  
Angelopoulos+ 2020, <https://doi.org/10.1007/s11214-020-00721-7>  
Min+ 2012, <https://doi.org/10.1029/2012JA017515>  
Usanova+ 2012, <https://doi.org/10.1029/2012JA018049>

Software:  
G. Van Rossum+ 1995, Python reference manual.  
Bezanson+ 2017, <https://doi.org/10.1137/141000671>  
Morley+, <https://doi.org/10.5281/zenodo.3252523>