

# Math 302: Series methods

Kameryn J Williams

University of Hawai'i at Mānoa

Spring 2021

# Undetermined coefficients

To solve an equation like

$$y'' + y' + y = x^3$$

we guessed that a particular solution looks like

$$A + Bx + Cx^2 + Dx^3$$

Then we solve for the values of the coefficients.

# Undetermined coefficients

To solve an equation like

$$y'' + y' + y = x^3$$

we guessed that a particular solution looks like

$$A + Bx + Cx^2 + Dx^3$$

Then we solve for the values of the coefficients.

What if, instead, to solve an equation like

$$ay'' + by' + cy = f(x)$$

we guessed that a particular solution looks like

$$A + Bx + Cx^2 + Dx^3 + Ex^4 + Fx^5 + \dots?$$

# Undetermined coefficients

To solve an equation like

$$y'' + y' + y = x^3$$

we guessed that a particular solution looks like

$$A + Bx + Cx^2 + Dx^3$$

Then we solve for the values of the coefficients.

What if, instead, to solve an equation like

$$ay'' + by' + cy = f(x)$$

we guessed that a particular solution looks like

$$A + Bx + Cx^2 + Dx^3 + Ex^4 + Fx^5 + \dots?$$

That is, we want to see what we can figure out if we represent the solution as a power series.

# Power series

A **power series** is an infinite series of the form

$$\sum_{n=0}^{\infty} a_n x^n = a_0 + a_1 x + a_2 x^2 + a_3 x^3 + \cdots$$

# Power series

A **power series** is an infinite series of the form

$$\sum_{n=0}^{\infty} a_n x^n = a_0 + a_1 x + a_2 x^2 + a_3 x^3 + \cdots$$

We can also talk about power series centered at an arbitrary point  $p$  rather than centered at 0:

$$\sum_{n=0}^{\infty} a_n (x-p)^n = a_0 + a_1 (x-p) + a_2 (x-p)^2 + \cdots$$

# Power series

A **power series** is an infinite series of the form

$$\sum_{n=0}^{\infty} a_n x^n = a_0 + a_1 x + a_2 x^2 + a_3 x^3 + \cdots$$

We can also talk about power series centered at an arbitrary point  $p$  rather than centered at 0:

$$\sum_{n=0}^{\infty} a_n (x-p)^n = a_0 + a_1 (x-p) + a_2 (x-p)^2 + \cdots$$

A power series centered at  $p$  has a **radius of convergence**  $R$ :

- If  $R = 0$  the series converges iff  $x = p$ .
- If  $0 < R < \infty$  the series converges if  $|x - p| < R$ . At the end points  $x = p \pm R$  it may either converge or diverge
- If  $R = \infty$  the series converges for any value of  $x$ .

# Power series

A **power series** is an infinite series of the form

$$\sum_{n=0}^{\infty} a_n x^n = a_0 + a_1 x + a_2 x^2 + a_3 x^3 + \cdots$$

We can also talk about power series centered at an arbitrary point  $p$  rather than centered at 0:

$$\sum_{n=0}^{\infty} a_n (x-p)^n = a_0 + a_1 (x-p) + a_2 (x-p)^2 + \cdots$$

A power series centered at  $p$  has a **radius of convergence**  $R$ :

- If  $R = 0$  the series converges iff  $x = p$ .
- If  $0 < R < \infty$  the series converges if  $|x - p| < R$ . At the end points  $x = p \pm R$  it may either converge or diverge
- If  $R = \infty$  the series converges for any value of  $x$ .

You can use convergence tests like you learned in Calc II to figure out the radius of convergence of a given power series.



# From power series to functions

$$\sum_{n=0}^{\infty} a_n x^n$$

If the **interval of convergence** of this power series is nontrivial (i.e.  $R > 0$ ), then the power series defines a continuous function on the interval  $p - R < x < p + R$ :

$$f(x) = \sum_{n=0}^{\infty} a_n x^n.$$

# From power series to functions

$$\sum_{n=0}^{\infty} a_n x^n$$

If the **interval of convergence** of this power series is nontrivial (i.e.  $R > 0$ ), then the power series defines a continuous function on the interval  $p - R < x < p + R$ :

$$f(x) = \sum_{n=0}^{\infty} a_n x^n.$$

Indeed, we can say more than just that  $f(x)$  is continuous. It is differentiable, and we know its derivative.

# From power series to functions

$$\sum_{n=0}^{\infty} a_n x^n$$

If the **interval of convergence** of this power series is nontrivial (i.e.  $R > 0$ ), then the power series defines a continuous function on the interval  $p - R < x < p + R$ :

$$f(x) = \sum_{n=0}^{\infty} a_n x^n.$$

Indeed, we can say more than just that  $f(x)$  is continuous. It is differentiable, and we know its derivative.

$$f'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{k=0}^{\infty} (k+1) a_{k+1} x^k$$

That is, you get the derivative by differentiating the power series term by term.

# From power series to functions

$$\sum_{n=0}^{\infty} a_n x^n$$

If the **interval of convergence** of this power series is nontrivial (i.e.  $R > 0$ ), then the power series defines a continuous function on the interval  $p - R < x < p + R$ :

$$f(x) = \sum_{n=0}^{\infty} a_n x^n.$$

Indeed, we can say more than just that  $f(x)$  is continuous. It is differentiable, and we know its derivative.

$$f'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{k=0}^{\infty} (k+1) a_{k+1} x^k$$

That is, you get the derivative by differentiating the power series term by term. We can repeat this to find  $f''(x)$ ,  $f'''(x)$ , and so on. In particular,  $f(x)$  is **infinitely differentiable**.

# From power series to functions

$$\sum_{n=0}^{\infty} a_n x^n$$

If the **interval of convergence** of this power series is nontrivial (i.e.  $R > 0$ ), then the power series defines a continuous function on the interval  $p - R < x < p + R$ :

$$f(x) = \sum_{n=0}^{\infty} a_n x^n.$$

Indeed, we can say more than just that  $f(x)$  is continuous. It is differentiable, and we know its derivative.

$$f'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{k=0}^{\infty} (k+1) a_{k+1} x^k$$

That is, you get the derivative by differentiating the power series term by term. We can repeat this to find  $f''(x)$ ,  $f'''(x)$ , and so on. In particular,  $f(x)$  is **infinitely differentiable**.

This works centered at  $p$ , not just centered at 0.

# From functions to power series

Consider a function given by a power series

$$f(x) = \sum_{n=0}^{\infty} a_n x^n.$$

# From functions to power series

Consider a function given by a power series

$$f(x) = \sum_{n=0}^{\infty} a_n x^n.$$

Using the facts about its derivatives we can determine the coefficients  $a_0$ , working backward from the formula for the power series of its derivative.

- $f(0) = a_0$

# From functions to power series

Consider a function given by a power series

$$f(x) = \sum_{n=0}^{\infty} a_n x^n.$$

Using the facts about its derivatives we can determine the coefficients  $a_0$ , working backward from the formula for the power series of its derivative.

- $f(0) = a_0$
- $f'(0) = a_1$



# From functions to power series

Consider a function given by a power series

$$f(x) = \sum_{n=0}^{\infty} a_n x^n.$$

Using the facts about its derivatives we can determine the coefficients  $a_0$ , working backward from the formula for the power series of its derivative.

- $f(0) = a_0$
- $f'(0) = a_1$
- $f''(0) = 2a_2$

# From functions to power series

Consider a function given by a power series

$$f(x) = \sum_{n=0}^{\infty} a_n x^n.$$

Using the facts about its derivatives we can determine the coefficients  $a_0$ , working backward from the formula for the power series of its derivative.

- $f(0) = a_0$
- $f'(0) = a_1$
- $f''(0) = 2a_2$
- $f'''(0) = 6a_3$

# From functions to power series

Consider a function given by a power series

$$f(x) = \sum_{n=0}^{\infty} a_n x^n.$$

Using the facts about its derivatives we can determine the coefficients  $a_0$ , working backward from the formula for the power series of its derivative.

- $f(0) = a_0$
- $f'(0) = a_1$
- $f''(0) = 2a_2$
- $f'''(0) = 6a_3$
- $f^{(n)}(0) = n!a_n$

# From functions to power series

Consider a function given by a power series

$$f(x) = \sum_{n=0}^{\infty} a_n x^n.$$

$$\text{So } a_n = \frac{f^{(n)}(0)}{n!}.$$

Using the facts about its derivatives we can determine the coefficients  $a_0$ , working backward from the formula for the power series of its derivative.

- $f(0) = a_0$
- $f'(0) = a_1$
- $f''(0) = 2a_2$
- $f'''(0) = 6a_3$
- $f^{(n)}(0) = n!a_n$

# From functions to power series

Consider a function given by a power series

$$f(x) = \sum_{n=0}^{\infty} a_n x^n.$$

So  $a_n = \frac{f^{(n)}(0)}{n!}$ .

If we instead centered at  $p$ :

$$a_n = \frac{f^{(n)}(p)}{n!}.$$

Using the facts about its derivatives we can determine the coefficients  $a_0$ , working backward from the formula for the power series of its derivative.

- $f(0) = a_0$
- $f'(0) = a_1$
- $f''(0) = 2a_2$
- $f'''(0) = 6a_3$
- $f^{(n)}(0) = n!a_n$

# From functions to power series

Consider a function given by a power series

$$f(x) = \sum_{n=0}^{\infty} a_n x^n.$$

Using the facts about its derivatives we can determine the coefficients  $a_0$ , working backward from the formula for the power series of its derivative.

- $f(0) = a_0$
- $f'(0) = a_1$
- $f''(0) = 2a_2$
- $f'''(0) = 6a_3$
- $f^{(n)}(0) = n!a_n$

$$\text{So } a_n = \frac{f^{(n)}(0)}{n!}.$$

If we instead centered at  $p$ :

$$a_n = \frac{f^{(n)}(p)}{n!}.$$

So we can write:

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(0)}{n!} x^n$$

for the power series for  $f(x)$ . We call this its **Taylor series**.

# Analytic functions

An **analytic** function is one which has a Taylor series expansion centered at  $p$  for every  $p$  in its domain.

# Analytic functions

An **analytic** function is one which has a Taylor series expansion centered at  $p$  for every  $p$  in its domain.

- Every analytic function is infinitely differentiable.
- But not every infinitely differentiable function is analytic.

$$f(x) = \begin{cases} e^{-1/x^2} & \text{if } x \neq 0 \\ 0 & \text{if } x = 0 \end{cases}$$

has all its derivatives  $= 0$  at  $x = 0$ , but that would give a Taylor series of  $0 + 0x + \cdots = 0$ .



# Analytic functions

An **analytic** function is one which has a Taylor series expansion centered at  $p$  for every  $p$  in its domain.

- Every analytic function is infinitely differentiable.
- But not every infinitely differentiable function is analytic.

$$f(x) = \begin{cases} e^{-1/x^2} & \text{if } x \neq 0 \\ 0 & \text{if } x = 0 \end{cases}$$

has all its derivatives  $= 0$  at  $x = 0$ , but that would give a Taylor series of  $0 + 0x + \dots = 0$ .

Let's remember the Taylor series centered at 0 for some important functions:

$$e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!} + \dots$$

$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} \pm \dots$$

$$\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} \pm \dots$$

# Solving differential equations using series

Let's see a basic example:  $xy' - y = 0$ .

# Solving differential equations using series

Let's see a basic example:  $xy' - y = 0$ .

Guess that the solution is given by a power series:

$$y = a_0 + a_1x + a_2x^2 + \cdots$$

# Solving differential equations using series

Let's see a basic example:  $xy' - y = 0$ .

Guess that the solution is given by a power series:

$$y = a_0 + a_1x + a_2x^2 + \cdots$$

Then,

$$y' = a_1 + 2a_2x + 3a_3x^2 + \cdots$$

# Solving differential equations using series

Let's see a basic example:  $xy' - y = 0$ .

Guess that the solution is given by a power series:

$$y = a_0 + a_1x + a_2x^2 + \cdots$$

Then,

$$y' = a_1 + 2a_2x + 3a_3x^2 + \cdots$$

Let's plug these into the equation:

$$\begin{aligned} & x(a_1 + 2a_2x + 3a_3x^2 + \cdots) \\ & - (a_0 + a_1x + a_2x^2 + \cdots) = 0 + 0x + 0x^2 + \cdots \end{aligned}$$

# Solving differential equations using series

Let's see a basic example:  $xy' - y = 0$ .

Guess that the solution is given by a power series:

$$y = a_0 + a_1x + a_2x^2 + \cdots$$

Then,

$$y' = a_1 + 2a_2x + 3a_3x^2 + \cdots$$

Let's plug these into the equation:

$$\begin{aligned} & x(a_1 + 2a_2x + 3a_3x^2 + \cdots) \\ & - (a_0 + a_1x + a_2x^2 + \cdots) = 0 + 0x + 0x^2 + \cdots \end{aligned}$$

Combine like terms:

$$-a_0 + 0x + a_2x^2 + \cdots = 0 + 0x + 0x^2 + \cdots$$

# Solving differential equations using series

Let's see a basic example:  $xy' - y = 0$ .

Guess that the solution is given by a power series:

$$y = a_0 + a_1x + a_2x^2 + \dots$$

Then,

$$y' = a_1 + 2a_2x + 3a_3x^2 + \dots$$

Let's plug these into the equation:

$$\begin{aligned} & x(a_1 + 2a_2x + 3a_3x^2 + \dots) \\ & - (a_0 + a_1x + a_2x^2 + \dots) = 0 + 0x + 0x^2 + \dots \end{aligned}$$

Combine like terms:

$$-a_0 + 0x + a_2x^2 + \dots = 0 + 0x + 0x^2 + \dots$$

Finally, solve for what values of the coefficients  $a_n$  makes this true. In this case, we quickly see that  $a_1$  can be anything and  $a_n = 0$  for  $n \neq 1$ .

# Solving differential equations using series

Let's see a basic example:  $xy' - y = 0$ .

Guess that the solution is given by a power series:

$$y = a_0 + a_1x + a_2x^2 + \dots$$

Then,

$$y' = a_1 + 2a_2x + 3a_3x^2 + \dots$$

Let's plug these into the equation:

$$\begin{aligned} & x(a_1 + 2a_2x + 3a_3x^2 + \dots) \\ & - (a_0 + a_1x + a_2x^2 + \dots) = 0 + 0x + 0x^2 + \dots \end{aligned}$$

Combine like terms:

$$-a_0 + 0x + a_2x^2 + \dots = 0 + 0x + 0x^2 + \dots$$

Finally, solve for what values of the coefficients  $a_n$  makes this true. In this case, we quickly see that  $a_1$  can be anything and  $a_n = 0$  for  $n \neq 1$ .

So the solution is  $y = a_1x$ , where  $a_1$  is an arbitrary constant.



# An existence theorem

$$y^{(n)} + \cdots + a_1(x)y' + a_0(x) = b(x)$$

If the coefficient functions  $a_i(x)$  and the function  $b(x)$  are all analytic on the same interval centered on  $p$ , then there is a unique solution satisfying the initial conditions

$$y(p) = v_0, \quad y'(p) = v_1, \quad \cdots, \quad y^{(n-1)}(p) = v_{n-1}$$

which is also analytic in the interval.

# An existence theorem

$$y^{(n)} + \cdots + a_1(x)y' + a_0(x) = b(x)$$

If the coefficient functions  $a_i(x)$  and the function  $b(x)$  are all analytic on the same interval centered on  $p$ , then there is a unique solution satisfying the initial conditions

$$y(p) = v_0, \quad y'(p) = v_1, \quad \cdots, \quad y^{(n-1)}(p) = v_{n-1}$$

which is also analytic in the interval.

In short, this result tells us that if all the parts of the equation are analytic functions, then it's valid to guess that the solution is given by a power series and use that to determine the solution.

# An existence theorem

$$y^{(n)} + \cdots + a_1(x)y' + a_0(x) = b(x)$$

If the coefficient functions  $a_i(x)$  and the function  $b(x)$  are all analytic on the same interval centered on  $p$ , then there is a unique solution satisfying the initial conditions

$$y(p) = v_0, \quad y'(p) = v_1, \quad \cdots, \quad y^{(n-1)}(p) = v_{n-1}$$

which is also analytic in the interval.

In short, this result tells us that if all the parts of the equation are analytic functions, then it's valid to guess that the solution is given by a power series and use that to determine the solution.

In particular, if the functions are all polynomials, exponential functions, sine/cosine, or combinations thereof, then we get a solution which is valid for all of  $\mathbb{R}$ .

# Approximating solutions

$$y'' - xy' + y = x, \text{ where } y(0) = y'(0) = 1$$

# Approximating solutions

$$y'' - xy' + y = x, \text{ where } y(0) = y'(0) = 1$$

Guess that the solution is given by a power series

$$y = a_0 + a_1x + a_2x^2 + \cdots$$

# Approximating solutions

$$y'' - xy' + y = x, \text{ where } y(0) = y'(0) = 1$$

Guess that the solution is given by a power series

$$y = a_0 + a_1x + a_2x^2 + \cdots$$

We know  $a_0 = y(0)$  and  $a_1 = y'(0)$ , so those are free. The rest we have to figure out by plugging things in.

# Approximating solutions

$$y'' - xy' + y = x, \text{ where } y(0) = y'(0) = 1$$

Guess that the solution is given by a power series

$$y = a_0 + a_1x + a_2x^2 + \cdots$$

We know  $a_0 = y(0)$  and  $a_1 = y'(0)$ , so those are free. The rest we have to figure out by plugging things in.

Solving for the first few terms:

$$y = 1 + x - \frac{x^2}{2} + \frac{x^3}{6} - \frac{x^4}{24} + \cdots$$

## Another example

$$(1+x)y'' + y = e^x$$

Let's try to work out what  $a_n$  looks like in terms of previous coefficients.



## Another example

$$(1+x)y'' + y = e^x$$

Let's try to work out what  $a_n$  looks like in terms of previous coefficients.

Get:

$$a_0 = \text{arbitrary}$$

$$a_1 = \text{arbitrary}$$

$$a_{n+2} = \underbrace{\frac{1}{n!} - \frac{na_{n+1}}{n+2} - \frac{a_n}{(n+2)(n+1)}}_{=b_{n+2}}$$

## Another example

$$(1+x)y'' + y = e^x$$

Let's try to work out what  $a_n$  looks like in terms of previous coefficients.

So we can write the solution as

Get:

$$y = a_0 + a_1x + \sum_{n=2}^{\infty} b_n x^n,$$

$a_0 = \text{arbitrary}$

$a_1 = \text{arbitrary}$

where  $a_0$  and  $a_1$  are arbitrary constants.

$$a_{n+2} = \underbrace{\frac{1}{n!} - \frac{na_{n+1}}{n+2} - \frac{a_n}{(n+2)(n+1)}}_{=b_{n+2}}$$

## Another example

$$(1+x)y'' + y = e^x$$

Let's try to work out what  $a_n$  looks like in terms of previous coefficients.

So we can write the solution as

Get:

$$y = a_0 + a_1x + \sum_{n=2}^{\infty} b_n x^n,$$

$a_0 = \text{arbitrary}$

$a_1 = \text{arbitrary}$

$$a_{n+2} = \underbrace{\frac{1}{n!} - \frac{na_{n+1}}{n+2} - \frac{a_n}{(n+2)(n+1)}}_{=b_{n+2}}$$

where  $a_0$  and  $a_1$  are arbitrary constants. This doesn't give us a nice way to write the solution in terms of elementary functions, but why should we expect to always be able to do so?

Nonetheless, we could use this to compute an approximation, using more terms for more precision.