

Math 302: Operators

Kameryn J Williams

University of Hawai'i at Mānoa

Spring 2021

Previously

We've been learning various methods for solving linear differential equations.

The next big topic is the [Laplace transform](#) which gives, among other things, yet another method for solving these.

But before we talk about the Laplace transform, let's step back a bit to take a more general view on [operators](#).

This isn't new content per se, but rather is placing content we've already learned into a more general framework.

Operators

- Probably, when you've seen functions in math classes, the inputs/outputs have been real numbers (or maybe complex numbers).
- But it's a more general concept.
- You can have a functions that take any type of mathematical object as input or output.
- You can even have a function whose inputs and outputs are functions.

Operators

- Probably, when you've seen functions in math classes, the inputs/outputs have been real numbers (or maybe complex numbers).
- But it's a more general concept.
- You can have a functions that take any type of mathematical object as input or output.
- You can even have a function whose inputs and outputs are functions.

To avoid confusion, let's use a different name for these.

- An **operator** is a function whose inputs/outputs are functions $\mathbb{R} \rightarrow \mathbb{R}$.

Operators

- Probably, when you've seen functions in math classes, the inputs/outputs have been real numbers (or maybe complex numbers).
- But it's a more general concept.
- You can have a functions that take any type of mathematical object as input or output.
- You can even have a function whose inputs and outputs are functions.

To avoid confusion, let's use a different name for these.

- An **operator** is a function whose inputs/outputs are functions $\mathbb{R} \rightarrow \mathbb{R}$.

The most important example is the **differential operator**, let's call it D :

$$Dy = y'$$

Operators

- Probably, when you've seen functions in math classes, the inputs/outputs have been real numbers (or maybe complex numbers).
- But it's a more general concept.
- You can have a functions that take any type of mathematical object as input or output.
- You can even have a function whose inputs and outputs are functions.

To avoid confusion, let's use a different name for these.

- An **operator** is a function whose inputs/outputs are functions $\mathbb{R} \rightarrow \mathbb{R}$.

The most important example is the **differential operator**, let's call it D :

$$Dy = y'$$

Another important example is integration:

$$\int_{x_0}^x f(t) dt$$

converts a function f into a new function.

Linear operators

We can write a basic property of differentiation in terms of operators:

$$D(af + bg) = aDf + bDg,$$

where a, b are constant real numbers and f, g are functions.

Linear operators

We can write a basic property of differentiation in terms of operators:

$$D(af + bg) = aDf + bDg,$$

where a, b are constant real numbers and f, g are functions.

If an operator satisfies this property we call it a **linear operator**.

- If you've taken linear algebra: linear operators are linear transformations acting on vector spaces of real-valued functions.

Linear operators

We can write a basic property of differentiation in terms of operators:

$$D(af + bg) = aDf + bDg,$$

where a, b are constant real numbers and f, g are functions.

If an operator satisfies this property we call it a **linear operator**.

- If you've taken linear algebra: linear operators are linear transformations acting on vector spaces of real-valued functions.

Another linear operator is integration.

Linear operators

We can write a basic property of differentiation in terms of operators:

$$D(af + bg) = aDf + bDg,$$

Can you come up with more linear operators?

where a, b are constant real numbers and f, g are functions.

If an operator satisfies this property we call it a **linear operator**.

- If you've taken linear algebra: linear operators are linear transformations acting on vector spaces of real-valued functions.

Another linear operator is integration.

Linear operators

We can write a basic property of differentiation in terms of operators:

$$D(af + bg) = aDf + bDg,$$

where a, b are constant real numbers and f, g are functions.

If an operator satisfies this property we call it a **linear operator**.

- If you've taken linear algebra: linear operators are linear transformations acting on vector spaces of real-valued functions.

Another linear operator is integration.

Can you come up with more linear operators?

- The **identity operator** $\text{id } f = f$

Linear operators

We can write a basic property of differentiation in terms of operators:

$$D(af + bg) = aDf + bDg,$$

where a, b are constant real numbers and f, g are functions.

If an operator satisfies this property we call it a **linear operator**.

- If you've taken linear algebra: linear operators are linear transformations acting on vector spaces of real-valued functions.

Another linear operator is integration.

Can you come up with more linear operators?

- The **identity operator** $\text{id } f = f$
- D^2 , i.e. $D^2f = f''$

Linear operators

We can write a basic property of differentiation in terms of operators:

$$D(af + bg) = aDf + bDg,$$

where a, b are constant real numbers and f, g are functions.

If an operator satisfies this property we call it a **linear operator**.

- If you've taken linear algebra: linear operators are linear transformations acting on vector spaces of real-valued functions.

Another linear operator is integration.

Can you come up with more linear operators?

- The **identity operator** $\text{id } f = f$
- D^2 , i.e. $D^2f = f''$
- D^n for any natural number n

Linear operators

We can write a basic property of differentiation in terms of operators:

$$D(af + bg) = aDf + bDg,$$

where a, b are constant real numbers and f, g are functions.

If an operator satisfies this property we call it a **linear operator**.

- If you've taken linear algebra: linear operators are linear transformations acting on vector spaces of real-valued functions.

Another linear operator is integration.

Can you come up with more linear operators?

- The **identity operator** $\text{id } f = f$
- D^2 , i.e. $D^2f = f''$
- D^n for any natural number n
- $S + T$, where S and T are linear

Linear operators

We can write a basic property of differentiation in terms of operators:

$$D(af + bg) = aDf + bDg,$$

where a, b are constant real numbers and f, g are functions.

If an operator satisfies this property we call it a **linear operator**.

- If you've taken linear algebra: linear operators are linear transformations acting on vector spaces of real-valued functions.

Another linear operator is integration.

Can you come up with more linear operators?

- The **identity operator** $\text{id } f = f$
- D^2 , i.e. $D^2f = f''$
- D^n for any natural number n
- $S + T$, where S and T are linear
- aT , where T is linear and a is a constant

Polynomial operators

Consider a polynomial

$$P(r) = a_n r^n + \cdots + a_1 r + a_0.$$

Then if T is a linear operator, so is

$$P(D) = a_n D^n + \cdots a_1 D^1 + a_0 D^0,$$

where $D^0 = \text{id}$ is the identity operator. We call $P(D)$ a **polynomial operator**.

Polynomial operators

Consider a polynomial

$$P(r) = a_n r^n + \cdots + a_1 r + a_0.$$

Then if T is a linear operator, so is

$$P(D) = a_n D^n + \cdots a_1 D^1 + a_0 D^0,$$

where $D^0 = \text{id}$ is the identity operator. We call $P(D)$ a **polynomial operator**.

Notation: we write e.g. 3 to mean the operator 3id .

Polynomial operators

Consider a polynomial

$$P(r) = a_n r^n + \cdots + a_1 r + a_0.$$

Then if T is a linear operator, so is

$$P(D) = a_n D^n + \cdots a_1 D^1 + a_0 D^0,$$

where $D^0 = \text{id}$ is the identity operator. We call $P(D)$ a **polynomial operator**.

Notation: we write e.g. 3 to mean the operator 3id .

For example, consider $P(r) = r^2 + 2r + 3$.

Then, $P(D)y = 0$ is another way to write the equation

$$y'' + 2y' + y = 0.$$

Polynomial operators

Consider a polynomial

$$P(r) = a_n r^n + \cdots + a_1 r + a_0.$$

Then if T is a linear operator, so is

$$P(D) = a_n D^n + \cdots a_1 D^1 + a_0 D^0,$$

where $D^0 = \text{id}$ is the identity operator. We call $P(D)$ a **polynomial operator**.

Notation: we write e.g. 3 to mean the operator 3id .

For example, consider $P(r) = r^2 + 2r + 3$.

Then, $P(D)y = 0$ is another way to write the equation

$$y'' + 2y' + y = 0.$$

Any linear homogeneous differential equation with constant coefficients is of the form $P(D)y = 0$ for some polynomial operator $P(D)$. Solving this equation is computing the **kernel** of the linear operator $P(D)$.

Polynomial operators

Consider a polynomial

$$P(r) = a_n r^n + \cdots + a_1 r + a_0.$$

Then if T is a linear operator, so is

$$P(D) = a_n D^n + \cdots a_1 D^1 + a_0 D^0,$$

where $D^0 = \text{id}$ is the identity operator. We call $P(D)$ a **polynomial operator**.

Notation: we write e.g. 3 to mean the operator 3id .

For example, consider $P(r) = r^2 + 2r + 3$.

Then, $P(D)y = 0$ is another way to write the equation

$$y'' + 2y' + y = 0.$$

Any linear homogeneous differential equation with constant coefficients is of the form

$P(D)y = 0$ for some polynomial operator $P(D)$. Solving this equation is computing the **kernel** of the linear operator $P(D)$.

Jargon: we say that $P(D)$ **annihilates** y if $P(D)y = 0$.

Polynomial operators

Consider a polynomial

$$P(r) = a_n r^n + \cdots + a_1 r + a_0.$$

Then if T is a linear operator, so is

$$P(D) = a_n D^n + \cdots + a_1 D^1 + a_0 D^0,$$

where $D^0 = \text{id}$ is the identity operator. We call $P(D)$ a **polynomial operator**.

Notation: we write e.g. 3 id .

For example, consider $P(r) = r^2 + 2r + 3$. Then, $P(D)y = 0$ is another way to write the equation

$$y'' + 2y' + y = 0.$$

Any linear homogeneous differential equation with constant coefficients is of the form $P(D)y = 0$ for some polynomial operator $P(D)$. Solving this equation is computing the **kernel** of the linear operator $P(D)$.

Jargon: we say that $P(D)$ **annihilates** y if $P(D)y = 0$.

If you've taken linear algebra: this is like when you find kernels of matrices, but in an infinite dimensional vector space.

The principle of superposition

Recall that the general solution to

$$a_n y^{(n)} + \cdots + a_1 y' + a_0 y = 0$$

is a linear combination of n many linearly independent individual solutions.

The principle of superposition

Recall that the general solution to

$$a_n y^{(n)} + \cdots + a_1 y' + a_0 y = 0$$

is a linear combination of n many linearly independent individual solutions.

This fact generalizes.

The principle of superposition

Recall that the general solution to

$$a_n y^{(n)} + \cdots + a_1 y' + a_0 y = 0$$

is a linear combination of n many linearly independent individual solutions.

This fact generalizes.

Suppose T is a linear operator and consider the equation

$$Ty = a_1 b_1 + \cdots + a_n b_n,$$

where a_i are constants and b_i are functions. Suppose you have solutions y_i to the equations

$$Ty = b_i.$$

Then, $a_1 y_1 + \cdots + a_n y_n$ is a solution to the original equation.

In other words: you can get the solution by breaking up the sum into pieces and solving each piece individually. For example, this is how you differentiate.

The algebra of linear operators

Linear operators satisfy most of the algebraic properties of the integers, where addition and multiplication of operators are defined as:

$$(S + T)y = Sy + Ty, \quad (ST)y = S(Ty).$$

The algebra of linear operators

Linear operators satisfy most of the algebraic properties of the integers, where addition and multiplication of operators are defined as:

$$(S + T)y = Sy + Ty, \quad (ST)y = S(Ty).$$

The exception is commutativity of multiplication: in general, ST and TS won't be the same operator. However, if they are polynomial operators $P(D)$ and $Q(D)$, then they will be the same:

$$P(D)Q(D) = Q(D)P(D).$$

The algebra of linear operators

Linear operators satisfy most of the algebraic properties of the integers, where addition and multiplication of operators are defined as:

$$(S + T)y = Sy + Ty, \quad (ST)y = S(Ty).$$

The exception is commutativity of multiplication: in general, ST and TS won't be the same operator. However, if they are polynomial operators $P(D)$ and $Q(D)$, then they will be the same:

$$P(D)Q(D) = Q(D)P(D).$$

The distributive law, associativity of addition and multiplication, and commutativity of addition all hold:

$$S(T + U) = ST + SU$$

$$(S + T)U = SU + TU$$

$$(S + T) + U = S + (T + U)$$

$$(ST)U = S(TU)$$

$$S + T = T + S$$

The algebra of linear operators

Linear operators satisfy most of the algebraic properties of the integers, where addition and multiplication of operators are defined as:

$$(S + T)y = Sy + Ty, \quad (ST)y = S(Ty).$$

The exception is commutativity of multiplication: in general, ST and TS won't be the same operator. However, if they are polynomial operators $P(D)$ and $Q(D)$, then they will be the same:

$$P(D)Q(D) = Q(D)P(D).$$

The distributive law, associativity of addition and multiplication, and commutativity of addition all hold:

$$S(T + U) = ST + SU$$

$$(S + T)U = SU + TU$$

$$(S + T) + U = S + (T + U)$$

$$(ST)U = S(TU)$$

$$S + T = T + S$$

(In math jargon: linear operators form a **noncommutative ring**, and restricting to the polynomial operators gives a **commutative ring**.)

Solving linear ODEs with operators

Due to time constraints, we won't cover it in this class, but I want to briefly mention that you can use operators to solve ODEs.

Solving linear ODEs with operators

Due to time constraints, we won't cover it in this class, but I want to briefly mention that you can use operators to solve ODEs.

- Consider the ODE
$$a_n y^{(n)} + \cdots + a_1 y' + a_0 y = b(x).$$
- This is in the form $P(D)y = b(x)$, where $P(r) = a_n r^n + \cdots + a_1 r + a_0$.
- You can solve $P(D)y = b(x)$ by determining an **inverse operator** for $P(D)$.
- Compare to how you solve $Dy = b(x)$ by means of the inverse operator for differentiation, namely integration.