

The Σ_1 -definable universal finite sequence

Kameryn J Williams

University of Hawai'i at Mānoa

Cantor Meets Robinson

Set Theory, Model Theory, and their Philosophy

2018 Dec 15

This is joint work with Joel David Hamkins and Philip Welch.

Multiversism versus universism in set theory

- **The universalist:** *The* universe of sets is uniquely determined.
- **The multiversalist:** There are many universes of sets, and every universe is contained inside a bigger, better universe.

Set-theoretic multiverses

- Zermelo: The set-theoretic universe is always dynamically growing upward; V is V_κ for κ inaccessible in a larger universe.

Set-theoretic multiverses

- Zermelo: The set-theoretic universe is always dynamically growing upward; V is V_κ for κ inaccessible in a larger universe.
- The generic multiverse: move to new universes by forcing or by going to grounds.

Set-theoretic multiverses

- Zermelo: The set-theoretic universe is always dynamically growing upward; V is V_κ for κ inaccessible in a larger universe.
- The generic multiverse: move to new universes by forcing or by going to grounds.
- The toy multiverse of countable transitive models.

Set-theoretic multiverses

- Zermelo: The set-theoretic universe is always dynamically growing upward; V is V_κ for κ inaccessible in a larger universe.
- The generic multiverse: move to new universes by forcing or by going to grounds.
- The toy multiverse of countable transitive models.
- A more radical multiversism: Hamkins's multiverse axioms, including the [Well-foundedness Mirage axiom](#)—every universe is seen to be ill-founded from some larger universe.

Set-theoretic multiverses

- Zermelo: The set-theoretic universe is always dynamically growing upward; V is V_κ for κ inaccessible in a larger universe.
- The generic multiverse: move to new universes by forcing or by going to grounds.
- The toy multiverse of countable transitive models.
- A more radical multiversism: Hamkins's multiverse axioms, including the [Well-foundedness Mirage axiom](#)—every universe is seen to be ill-founded from some larger universe.
 - The multiverse consisting of the countable, [recursively saturated](#) models of set theory satisfy Hamkins's axioms. (Gitman and Hamkins)

Potentialism as a general framework

- A **potentialist system** is a collection of structures of the same type, ordered by a relation \subseteq which refines the substructure relation.
- Used to formalize a domain we think of as dynamically growing.

Potentialism as a general framework

- A **potentialist system** is a collection of structures of the same type, ordered by a relation \subseteq which refines the substructure relation.
- Used to formalize a domain we think of as dynamically growing.
- $\Box\varphi$ holds in a world M if φ holds in every N extending M .
- $\Diamond\varphi$ holds in a world M if φ holds in some N extending M .

Potentialism as a general framework

- A **potentialist system** is a collection of structures of the same type, ordered by a relation \subseteq which refines the substructure relation.
- Used to formalize a domain we think of as dynamically growing.
- $\Box\varphi$ holds in a world M if φ holds in every N extending M .
- $\Diamond\varphi$ holds in a world M if φ holds in some N extending M .
- **Warning!** There is no guarantee that a potentialist system be **linearly ordered** or even **directed**. There are **branching** potentialist systems which have incompatible extensions.

The modal logic of potentialism

- A modal assertion is **valid** if it is true at any world under any substitution of propositional variables.
- $\Box p \Rightarrow p$ is valid, because if φ holds in every structure extending M , it in particular holds at M .

The modal logic of potentialism

- A modal assertion is **valid** if it is true at any world under any substitution of propositional variables.
- $\Box p \Rightarrow p$ is valid, because if φ holds in every structure extending M , it in particular holds at M .
- Key Question: given a potentialist system, what are its modal validities?

The modal logic of potentialism

- A modal assertion is **valid** if it is true at any world under any substitution of propositional variables.
- $\Box p \Rightarrow p$ is valid, because if φ holds in every structure extending M , it in particular holds at M .
- Key Question: given a potentialist system, what are its modal validities?
- The modal theory **S4** is valid in every potentialist system, where S4 has axioms

$$\Box(p \Rightarrow q) \Rightarrow (\Box p \Rightarrow \Box q)$$

$$\neg \Diamond p \Leftrightarrow \Box \neg p$$

$$\Box p \Rightarrow p$$

$$\Box p \Rightarrow \Box \Box p$$

- Because the potentialist system is partially ordered by \subseteq .

The modal logic of potentialism

- A modal assertion is **valid** if it is true at any world under any substitution of propositional variables.
- $\Box p \Rightarrow p$ is valid, because if φ holds in every structure extending M , it in particular holds at M .
- Key Question: given a potentialist system, what are its modal validities?
- The modal theory **S4** is valid in every potentialist system, where S4 has axioms

$$\Box(p \Rightarrow q) \Rightarrow (\Box p \Rightarrow \Box q)$$

$$\neg \Diamond p \Leftrightarrow \Box \neg p$$

$$\Box p \Rightarrow p$$

$$\Box p \Rightarrow \Box \Box p$$

- Because the potentialist system is partially ordered by \subseteq .
- This gives an easy lower bound. The real work is in getting upper bounds.

Buttons and switches (Hamkins and Löwe)

We can analyze the modal validities of a potentialist system using **control statements**.

- A **button** is a statement β so that $\Diamond\Box\beta$ holds in every world. The button is **pushed** if $\Box\beta$ holds.

Buttons and switches (Hamkins and Löwe)

We can analyze the modal validities of a potentialist system using **control statements**.

- A **button** is a statement β so that $\Diamond\Box\beta$ holds in every world. The button is **pushed** if $\Box\beta$ holds.
- A **switch** is a statement σ so that $\Diamond\sigma$ and $\Diamond\neg\sigma$ hold in every world.

Buttons and switches (Hamkins and Löwe)

We can analyze the modal validities of a potentialist system using **control statements**.

- A **button** is a statement β so that $\Diamond\Box\beta$ holds in every world. The button is **pushed** if $\Box\beta$ holds.
- A **switch** is a statement σ so that $\Diamond\sigma$ and $\Diamond\neg\sigma$ hold in every world.
- For forcing potentialism: CH is a switch, “ ω_1^L is countable” is a button.

Buttons and switches (Hamkins and Löwe)

We can analyze the modal validities of a potentialist system using **control statements**.

- A **button** is a statement β so that $\Diamond\Box\beta$ holds in every world. The button is **pushed** if $\Box\beta$ holds.
- A **switch** is a statement σ so that $\Diamond\sigma$ and $\Diamond\neg\sigma$ hold in every world.
- For forcing potentialism: CH is a switch, “ ω_1^L is countable” is a button.
- If there are arbitrarily large families of independent switches, then the modal validities are contained within S5, which is S4 plus the axiom $\Diamond\Box p \Rightarrow p$.

Buttons and switches (Hamkins and Löwe)

We can analyze the modal validities of a potentialist system using **control statements**.

- A **button** is a statement β so that $\Diamond\Box\beta$ holds in every world. The button is **pushed** if $\Box\beta$ holds.
- A **switch** is a statement σ so that $\Diamond\sigma$ and $\Diamond\neg\sigma$ hold in every world.
- For forcing potentialism: CH is a switch, “ ω_1^L is countable” is a button.
- If there are arbitrarily large families of independent switches, then the modal validities are contained within S5, which is S4 plus the axiom $\Diamond\Box p \Rightarrow p$.
- If there are arbitrarily large families of independent buttons and switches, then the modal validities are contained within S4.2, which is S4 plus the axiom $\Diamond\Box p \Rightarrow \Box\Diamond p$.

Potentialism in set theory

- Zermelo upward potentialism: worlds are V_κ for κ inaccessible.
 - Height potentialist.
- Forcing potentialism: worlds are forcing extensions of M .
 - Width potentialist.
- Countable transitive model potentialism.
 - Height and width potentialist.

Potentialism in set theory

- Zermelo upward potentialism: worlds are V_κ for κ inaccessible.
 - Height potentialist.
 - [Modal validities are S4.3.](#) (Hamkins and Linnebo)
 - S4.3 is S4 plus

$$(\Diamond \varphi \wedge \Diamond \psi) \Rightarrow (\varphi \wedge \Diamond \psi) \vee (\Diamond \varphi \wedge \psi).$$

- Forcing potentialism: worlds are forcing extensions of M .
 - Width potentialist.
 - [Modal validities are S4.2.](#) (Hamkins and Löwe)
- Countable transitive model potentialism.
 - Height and width potentialist.
 - [Modal validities are S4.2.](#) (Hamkins and Linnebo)

Flavors of potentialism

- S5 maximality principles
- S4.3 linear
- S4.2 directed
- S4 incompatible branching

The Σ_2 -definable universal finite set for rank-extensions

$N \supseteq M$ is a **rank-extension** of M if every $b \in N \setminus M$ has rank in $N \setminus M$.

Theorem (Hamkins and Woodin)

There is a Σ_2 definition for a finite set $\{b_0, \dots, b_n\}$ with the following properties.

- 1 ZFC proves that the definition defines a finite set.
- 2 In any transitive model of ZFC the set is empty.
- 3 If $M \models \text{ZFC}$ is countable, has s as its universal finite set, and $t \in M$ is a finite set extending s , then there is $N \models \text{ZFC}$ a rank-extension of M which has t as its universal finite set.

The Σ_2 -definable universal finite set for rank-extensions

$N \supseteq M$ is a **rank-extension** of M if every $b \in N \setminus M$ has rank in $N \setminus M$.

Theorem (Hamkins and Woodin)

There is a Σ_2 definition for a finite set $\{b_0, \dots, b_n\}$ with the following properties.

- 1 ZFC proves that the definition defines a finite set.
- 2 In any transitive model of ZFC the set is empty.
- 3 If $M \models \text{ZFC}$ is countable, has s as its universal finite set, and $t \in M$ is a finite set extending s , then there is $N \models \text{ZFC}$ a rank-extension of M which has t as its universal finite set.

Theorem (Hamkins and Woodin)

The modal validities of rank-extensional set theoretic potentialism are precisely S4.

End-extensions versus rank-extensions

- $N \supseteq M$ is an **end-extension** of M if $a \in b$ in N and b in M implies a is in M .
- Elementary end-extensions are always rank-extensions.
- But not all end-extensions are rank-extensions. For example, if M is an inner model of N then N end-extends M .

End-extensions versus rank-extensions

- $N \supseteq M$ is an **end-extension** of M if $a \in b$ in N and b in M implies a is in M .
- Elementary end-extensions are always rank-extensions.
- But not all end-extensions are rank-extensions. For example, if M is an inner model of N then N end-extends M .
- Key fact: the assertions which are preserved in arbitrary end-extensions are the Σ_1 assertions.

$$\exists y \underbrace{\varphi(x, y)}_{\text{quantifiers bounded}}$$

The Σ_1 -definable universal finite sequence for end-extensions

Let ZF^+ be a computably enumerable extension of ZF .

Theorem (Hamkins, Welch, W.)

There is a Σ_1 definition for a finite sequence b_0, b_1, \dots, b_n with the following properties.

- ❶ ZF^+ proves that the sequence is finite.
- ❷ In any transitive model of ZF^+ the sequence is empty.
- ❸ Let M be a countable model of ZF^+ which defines the sequence as s . Then if t in M is any finite sequence extending s , there is $N \models ZF^+$ end-extending M in which the universal sequence is t .

The Σ_1 -definable universal finite sequence for end-extensions

Let ZF^+ be a computably enumerable extension of ZF .

Theorem (Hamkins, Welch, W.)

There is a Σ_1 definition for a finite sequence b_0, b_1, \dots, b_n with the following properties.

- ❶ ZF^+ proves that the sequence is finite.
- ❷ In any transitive model of ZF^+ the sequence is empty.
- ❸ Let M be a countable model of ZF^+ which defines the sequence as s . Then if t in M is any finite sequence extending s , there is $N \models ZF^+$ end-extending M in which the universal sequence is t .
- ❹ Indeed, it suffices in (3) that M has an inner model W of ZF^+ satisfying such.

The definition of the universal finite sequence—Process A

Intended for ω -nonstandard models. A different process is used for ω -standard models.

- Proceed in stages to produce b_0, b_1, \dots, b_n , using auxiliary information: countable ordinals $\alpha_0 < \alpha_1 < \dots < \alpha_n$ and natural numbers $k_0 > k_1 > \dots > k_n$.

The definition of the universal finite sequence—Process A

Intended for ω -nonstandard models. A different process is used for ω -standard models.

- Proceed in stages to produce b_0, b_1, \dots, b_n , using auxiliary information: countable ordinals $\alpha_0 < \alpha_1 < \dots < \alpha_n$ and natural numbers $k_0 > k_1 > \dots > k_n$.
- At stage n : Are there $\alpha > \alpha_{n-1}$, $k < k_{n-1}$, and $b \in L_\alpha$ so that L_α has **no** end-extension to N satisfying the first k axioms of ZF^+ plus “process A succeeds at stage n and defines b ”? If so, stage n is successful and set (b_n, α_n, k_n) to be the L -least triple of such (b, α, k) .

The definition of the universal finite sequence—Process A

Intended for ω -nonstandard models. A different process is used for ω -standard models.

- Proceed in stages to produce b_0, b_1, \dots, b_n , using auxiliary information: countable ordinals $\alpha_0 < \alpha_1 < \dots < \alpha_n$ and natural numbers $k_0 > k_1 > \dots > k_n$.
- At stage n : Are there $\alpha > \alpha_{n-1}$, $k < k_{n-1}$, and $b \in L_\alpha$ so that L_α has **no** end-extension to N satisfying the first k axioms of ZF^+ plus “process A succeeds at stage n and defines b ”? If so, stage n is successful and set (b_n, α_n, k_n) to be the L -least triple of such (b, α, k) .
- This self-reference is allowed by the **Gödel–Carnap fixed-point lemma**.

The definition of the universal finite sequence—Process A

Intended for ω -nonstandard models. A different process is used for ω -standard models.

- Proceed in stages to produce b_0, b_1, \dots, b_n , using auxiliary information: countable ordinals $\alpha_0 < \alpha_1 < \dots < \alpha_n$ and natural numbers $k_0 > k_1 > \dots > k_n$.
- At stage n : Are there $\alpha > \alpha_{n-1}$, $k < k_{n-1}$, and $b \in L_\alpha$ so that L_α has **no** end-extension to N satisfying the first k axioms of ZF^+ plus “process A succeeds at stage n and defines b ”? If so, stage n is successful and set (b_n, α_n, k_n) to be the L -least triple of such (b, α, k) .
- This self-reference is allowed by the **Gödel–Carnap fixed-point lemma**.
- Claim: The map $n \mapsto (b_n, \alpha_n, k_n)$ is Σ_1 -definable.
- Claim: Each k_i must be nonstandard.
- Claim: There are only finitely many successful stages.

Seeing that Process A has the plus one extension property

Consider countable M in which the universal sequence is b_0, \dots, b_{n-1} and take any $b \in M$ and nonstandard $k < k_{n-1}$.

- Because stage n is unsuccessful in M this means L^M thinks every countable set can be end-extended to a model of the first k axioms of ZF^+ in which the universal sequence is b_0, \dots, b_{n-1}, b .

Seeing that Process A has the plus one extension property

Consider countable M in which the universal sequence is b_0, \dots, b_{n-1} and take any $b \in M$ and nonstandard $k < k_{n-1}$.

- Because stage n is unsuccessful in M this means L^M thinks every countable set can be end-extended to a model of the first k axioms of ZF^+ in which the universal sequence is b_0, \dots, b_{n-1}, b .
- This is a Π_2^1 assertion, so by the [Shoenfield absoluteness theorem](#) is true in M . It is also true in any elementary end-extension of M , as well as their forcing extensions.

Seeing that Process A has the plus one extension property

Consider countable M in which the universal sequence is b_0, \dots, b_{n-1} and take any $b \in M$ and nonstandard $k < k_{n-1}$.

- Because stage n is unsuccessful in M this means L^M thinks every countable set can be end-extended to a model of the first k axioms of ZF^+ in which the universal sequence is b_0, \dots, b_{n-1}, b .
- This is a Π_2^1 assertion, so by the [Shoenfield absoluteness theorem](#) is true in M . It is also true in any elementary end-extension of M , as well as their forcing extensions.
- By the [Keisler–Morley theorem](#), let M^+ be an elementary end extension of M and fix an ordinal θ in $M^+ \setminus M$. Consider $M^+[G]$ the forcing extension in which V_θ is collapsed to be countable.

Seeing that Process A has the plus one extension property

Consider countable M in which the universal sequence is b_0, \dots, b_{n-1} and take any $b \in M$ and nonstandard $k < k_{n-1}$.

- Because stage n is unsuccessful in M this means L^M thinks every countable set can be end-extended to a model of the first k axioms of ZF^+ in which the universal sequence is b_0, \dots, b_{n-1}, b .
- This is a Π_2^1 assertion, so by the [Shoenfield absoluteness theorem](#) is true in M . It is also true in any elementary end-extension of M , as well as their forcing extensions.
- By the [Keisler–Morley theorem](#), let M^+ be an elementary end extension of M and fix an ordinal θ in $M^+ \setminus M$. Consider $M^+[G]$ the forcing extension in which V_θ is collapsed to be countable.
- So V_θ^M has in $M^+[G]$ an end-extension N in which the universal sequence is b_0, \dots, b_{n-1}, b . But N is also an end-extension of M .

Process B —for ω -standard models

- Again go in stages: produce b_0, b_1, \dots, b_n using auxiliary information countable ordinals $\alpha_0 < \alpha_1 < \dots < \alpha_n$ and countable ordinals $\lambda_0 > \lambda_1 > \dots > \lambda_n$.

Process B —for ω -standard models

- Again go in stages: produce b_0, b_1, \dots, b_n using auxiliary information countable ordinals $\alpha_0 < \alpha_1 < \dots < \alpha_n$ and countable ordinals $\lambda_0 > \lambda_1 > \dots > \lambda_n$.
- At stage n : are there $\alpha > \alpha_{n-1}$, $\lambda < \lambda_n$, and $b \in L_\alpha$ so that L_α has **no** end-extension to N satisfying ZF^+ plus “process B succeeds at stage n and defines b ”, with the well-founded tree canonically representing this Π_1^1 assertion having rank λ ? If so, stage n is successful and set $(b_n, \alpha_n, \lambda_n)$ to be the L -least triple of such (b, α, λ) .

Process B —for ω -standard models

- Again go in stages: produce b_0, b_1, \dots, b_n using auxiliary information countable ordinals $\alpha_0 < \alpha_1 < \dots < \alpha_n$ and countable ordinals $\lambda_0 > \lambda_1 > \dots > \lambda_n$.
- At stage n : are there $\alpha > \alpha_{n-1}$, $\lambda < \lambda_n$, and $b \in L_\alpha$ so that L_α has **no** end-extension to N satisfying ZF^+ plus “process B succeeds at stage n and defines b ”, with the well-founded tree canonically representing this Π_1^1 assertion having rank λ ? If so, stage n is successful and set $(b_n, \alpha_n, \lambda_n)$ to be the L -least triple of such (b, α, λ) .
- Claim: The map $n \mapsto (b_n, \alpha_n, \lambda_n)$ is Σ_1 -definable.
- Claim: Each λ_i must be nonstandard.
- Claim: There are only finitely many successful stages.

Process B —for ω -standard models

- Again go in stages: produce b_0, b_1, \dots, b_n using auxiliary information countable ordinals $\alpha_0 < \alpha_1 < \dots < \alpha_n$ and countable ordinals $\lambda_0 > \lambda_1 > \dots > \lambda_n$.
- At stage n : are there $\alpha > \alpha_{n-1}$, $\lambda < \lambda_n$, and $b \in L_\alpha$ so that L_α has **no** end-extension to N satisfying ZF^+ plus “process B succeeds at stage n and defines b ”, with the well-founded tree canonically representing this Π_1^1 assertion having rank λ ? If so, stage n is successful and set $(b_n, \alpha_n, \lambda_n)$ to be the L -least triple of such (b, α, λ) .
- Claim: The map $n \mapsto (b_n, \alpha_n, \lambda_n)$ is Σ_1 -definable.
- Claim: Each λ_i must be nonstandard.
- Claim: There are only finitely many successful stages.
- Claim: Process B has the plus one extension property.

Process B —for ω -standard models

- Again go in stages: produce b_0, b_1, \dots, b_n using auxiliary information countable ordinals $\alpha_0 < \alpha_1 < \dots < \alpha_n$ and countable ordinals $\lambda_0 > \lambda_1 > \dots > \lambda_n$.
- At stage n : are there $\alpha > \alpha_{n-1}$, $\lambda < \lambda_n$, and $b \in L_\alpha$ so that L_α has **no** end-extension to N satisfying ZF^+ plus “process B succeeds at stage n and defines b ”, with the well-founded tree canonically representing this Π_1^1 assertion having rank λ ? If so, stage n is successful and set $(b_n, \alpha_n, \lambda_n)$ to be the L -least triple of such (b, α, λ) .
- Claim: The map $n \mapsto (b_n, \alpha_n, \lambda_n)$ is Σ_1 -definable.
- Claim: Each λ_i must be nonstandard.
- Claim: There are only finitely many successful stages.
- Claim: Process B has the plus one extension property.

Can merge Processes A and B into a single Process C which works for all models.

The Barwise extension theorem

The Barwise extension theorem can be derived as a corollary of our theorem.

Theorem (Barwise)

Every countable model of ZF end-extends to a model of $ZFC + V = L$.

The universal sequence for L -extensions

Corollary (Hamkins, Welch, W.)

There is a Σ_1 definition for a finite sequence b_0, b_1, \dots, b_n with the following properties.

- ① $ZF + V = L$ proves that the sequence is finite.
- ② In any standard model of $ZF + V = L$ the sequence is empty.
- ③ Let M be a countable model of $ZF + V = L$ which defines the sequence as s . Then if t in M is any finite sequence extending s , there is an L -extension N of M in which the universal sequence is t .


Railyard labelings

- A **tree** is a partial order T so that $\{s \in T : s \leq t\}$ is well-ordered for every $t \in T$. A **pre-tree** is a pre-order which quotients to a tree.
- A **railyard labeling** of a pre-tree T is an assignment ρ_t of statements to nodes $t \in T$ so that each structure satisfies exactly one ρ_t and $\diamond \rho_s$ holds iff $t \leq_T s$.

Railyard labelings


- A **tree** is a partial order T so that $\{s \in T : s \leq t\}$ is well-ordered for every $t \in T$. A **pre-tree** is a pre-order which quotients to a tree.
- A **railyard labeling** of a pre-tree T is an assignment ρ_t of statements to nodes $t \in T$ so that each structure satisfies exactly one ρ_t and $\Diamond \rho_s$ holds iff $t \leq_T s$.
- If there are railyard labelings for every finite pre-tree, then the modal validities for the corresponding potentialist system are contained within S4.

The universal finite sequence and railyard labelings




The universal finite sequence and railyard labelings

- Step 1: the subsequence $\langle n_i \rangle$ of finite ordinals from the universal finite sequence tell you how to descend the tree to determine your cluster. If B is the branching of the current node, then $n_i \bmod B$ tells you where to go.



The universal finite sequence and railyard labelings

- Step 1: the subsequence $\langle n_i \rangle$ of finite ordinals from the universal finite sequence tell you how to descend the tree to determine your cluster. If B is the branching of the current node, then $n_i \bmod B$ tells you where to go.
- Step 2: the final infinite ordinal $\lambda + m$ on the sequence tells you where in your cluster you are. If K is the size of the cluster, then $m \bmod K$ identifies your node in the cluster. (If no infinite ordinals are on the sequence, default to 0.)



The modal validities of end-extensional set theoretic potentialism

Theorem (Hamkins, Welch, W.)

Consider the potentialist system consisting of countable models of ZF^+ ordered by end-extension.

- ① *For any world M , the modal validities, allowing for a single parameter for the length of the universal finite sequence, are precisely S4.*
- ② *For any ω -standard world M , the modal validities, allowing no parameters, are precisely S4.*

Thank you!