

# The $\Sigma_1$ -definable universal finite sequence

Kameryn J Williams

University of Hawai'i at Mānoa

## Cantor Meets Robinson

Set Theory, Model Theory, and their Philosophy

2018 Dec 15

# Multiversism versus universism in set theory

- **The universalist:** *The* universe of sets is uniquely determined.
- **The multiversist:** There are many universes of sets, and every universe is contained inside a bigger, better universe.

# Set-theoretic multiverses

- Zermelo: The set-theoretic universe is always dynamically growing upward;  $V$  is  $V_\kappa$  for  $\kappa$  inaccessible in a larger universe.

# Set-theoretic multiverses

- Zermelo: The set-theoretic universe is always dynamically growing upward;  $V$  is  $V_\kappa$  for  $\kappa$  inaccessible in a larger universe.
- The generic multiverse: move to new universes by forcing or by going to grounds.

# Set-theoretic multiverses

- Zermelo: The set-theoretic universe is always dynamically growing upward;  $V$  is  $V_\kappa$  for  $\kappa$  inaccessible in a larger universe.
- The generic multiverse: move to new universes by forcing or by going to grounds.
- The toy multiverse of countable transitive models.

# Set-theoretic multiverses

- Zermelo: The set-theoretic universe is always dynamically growing upward;  $V$  is  $V_\kappa$  for  $\kappa$  inaccessible in a larger universe.
- The generic multiverse: move to new universes by forcing or by going to grounds.
- The toy multiverse of countable transitive models.
- A more radical multiversism: Hamkins's multiverse axioms, including the [Well-foundedness Mirage axiom](#)—every universe is seen to be ill-founded from some larger universe.

# Potentialism as a general framework

- A **potentialist system** is a collection of structures of the same type, ordered by a relation  $\subseteq$  which refines the substructure relation.
- Used to formalize a domain we think of as dynamically growing.

# Potentialism as a general framework

- A **potentialist system** is a collection of structures of the same type, ordered by a relation  $\subseteq$  which refines the substructure relation.
- Used to formalize a domain we think of as dynamically growing.
- $\Box\varphi$  holds in a world  $M$  if  $\varphi$  holds in every  $N$  extending  $M$ .
- $\Diamond\varphi$  holds in a world  $M$  if  $\varphi$  holds in some  $N$  extending  $M$ .



# Potentialism as a general framework

- A **potentialist system** is a collection of structures of the same type, ordered by a relation  $\subseteq$  which refines the substructure relation.
- Used to formalize a domain we think of as dynamically growing.
- $\Box\varphi$  holds in a world  $M$  if  $\varphi$  holds in every  $N$  extending  $M$ .
- $\Diamond\varphi$  holds in a world  $M$  if  $\varphi$  holds in some  $N$  extending  $M$ .
- **Warning!** There is no guarantee that a potentialist system be **linearly ordered** or even **directed**. There are **branching** potentialist systems which have incompatible extensions.

# The modal logic of potentialism

- A modal assertion is **valid** if it is true at any world under any substitution of propositional variables.
- $\Box p \Rightarrow p$  is valid, because if  $\varphi$  holds in every structure extending  $M$ , it in particular holds at  $M$ .

# The modal logic of potentialism

- A modal assertion is **valid** if it is true at any world under any substitution of propositional variables.
- $\Box p \Rightarrow p$  is valid, because if  $\varphi$  holds in every structure extending  $M$ , it in particular holds at  $M$ .
- Key Question: given a potentialist system, what are its modal validities?

# The modal logic of potentialism

- A modal assertion is **valid** if it is true at any world under any substitution of propositional variables.
- $\Box p \Rightarrow p$  is valid, because if  $\varphi$  holds in every structure extending  $M$ , it in particular holds at  $M$ .
- Key Question: given a potentialist system, what are its modal validities?
- The modal theory **S4** is valid in every potentialist system, where S4 has axioms

$$\Box(p \Rightarrow q) \Rightarrow (\Box p \Rightarrow \Box q)$$

$$\neg \Diamond p \Leftrightarrow \Box \neg p$$

$$\Box p \Rightarrow p$$

$$\Box p \Rightarrow \Box \Box p$$

- Because the potentialist system is partially ordered by  $\subseteq$ .

# The modal logic of potentialism

- A modal assertion is **valid** if it is true at any world under any substitution of propositional variables.
- $\Box p \Rightarrow p$  is valid, because if  $\varphi$  holds in every structure extending  $M$ , it in particular holds at  $M$ .
- Key Question: given a potentialist system, what are its modal validities?
- The modal theory **S4** is valid in every potentialist system, where S4 has axioms

$$\Box(p \Rightarrow q) \Rightarrow (\Box p \Rightarrow \Box q)$$

$$\neg \Diamond p \Leftrightarrow \Box \neg p$$

$$\Box p \Rightarrow p$$

$$\Box p \Rightarrow \Box \Box p$$

- Because the potentialist system is partially ordered by  $\subseteq$ .
- This gives an easy lower bound. The real work is in getting upper bounds.

# Buttons and switches (Hamkins and Löwe)

We can analyze the modal validities of a potentialist system using **control statements**.

- A **button** is a statement  $\beta$  so that  $\Diamond\Box\beta$  holds in every world. The button is **pushed** if  $\Box\beta$  holds.

# Buttons and switches (Hamkins and Löwe)

We can analyze the modal validities of a potentialist system using **control statements**.

- A **button** is a statement  $\beta$  so that  $\Diamond\Box\beta$  holds in every world. The button is **pushed** if  $\Box\beta$  holds.
- A **switch** is a statement  $\sigma$  so that  $\Diamond\sigma$  and  $\Diamond\neg\sigma$  hold in every world.

# Buttons and switches (Hamkins and Löwe)

We can analyze the modal validities of a potentialist system using **control statements**.

- A **button** is a statement  $\beta$  so that  $\Diamond\Box\beta$  holds in every world. The button is **pushed** if  $\Box\beta$  holds.
- A **switch** is a statement  $\sigma$  so that  $\Diamond\sigma$  and  $\Diamond\neg\sigma$  hold in every world.
- For forcing potentialism: CH is a switch, “ $\omega_1^L$  is countable” is a button.



# Buttons and switches (Hamkins and Löwe)

We can analyze the modal validities of a potentialist system using **control statements**.

- A **button** is a statement  $\beta$  so that  $\Diamond\Box\beta$  holds in every world. The button is **pushed** if  $\Box\beta$  holds.
- A **switch** is a statement  $\sigma$  so that  $\Diamond\sigma$  and  $\Diamond\neg\sigma$  hold in every world.
- For forcing potentialism: CH is a switch, “ $\omega_1^L$  is countable” is a button.
- If there are arbitrarily large families of independent switches, then the modal validities are contained within S5, which is S4 plus the axiom  $\Diamond\Box p \Rightarrow p$ .

# Buttons and switches (Hamkins and Löwe)

We can analyze the modal validities of a potentialist system using **control statements**.

- A **button** is a statement  $\beta$  so that  $\Diamond\Box\beta$  holds in every world. The button is **pushed** if  $\Box\beta$  holds.
- A **switch** is a statement  $\sigma$  so that  $\Diamond\sigma$  and  $\Diamond\neg\sigma$  hold in every world.
- For forcing potentialism: CH is a switch, “ $\omega_1^L$  is countable” is a button.
- If there are arbitrarily large families of independent switches, then the modal validities are contained within S5, which is S4 plus the axiom  $\Diamond\Box p \Rightarrow p$ .
- If there are arbitrarily large families of independent buttons and switches, then the modal validities are contained within S4.2, which is S4 plus the axiom  $\Diamond\Box p \Rightarrow \Box\Diamond p$ .

# Potentialism in set theory

- Zermelo upward potentialism: worlds are  $V_\kappa$  for  $\kappa$  inaccessible.
- Forcing potentialism: worlds are forcing extensions of  $M$ .
- Countable transitive model potentialism.

# Potentialism in set theory

- Zermelo upward potentialism: worlds are  $V_\kappa$  for  $\kappa$  inaccessible.
  - Modal validities are S4.3. (Hamkins and Linnebo)
- Forcing potentialism: worlds are forcing extensions of  $M$ .
  - Modal validities are S4.2. (Hamkins and Löwe)
- Countable transitive model potentialism.
  - Modal validities are S4.2. (Hamkins and Linnebo)

# The $\Sigma_2$ -definable universal finite set for rank-extensions

$N \supseteq M$  is a **rank-extension** of  $M$  if every  $b \in N \setminus M$  has rank in  $N \setminus M$ .

## Theorem (Hamkins and Woodin)

*There is a  $\Sigma_2$  definition for a finite set  $\{b_0, \dots, b_n\}$  with the following properties.*

- ① *ZFC proves that the definition defines a finite set.*
- ② *In any transitive model of ZFC the set is empty.*
- ③ *If  $M \models \text{ZFC}$  is countable, has  $s$  as its universal finite set, and  $t \in M$  is a finite set extending  $s$ , then there is  $N \models \text{ZFC}$  a rank-extension of  $M$  which has  $t$  as its universal finite set.*

# The $\Sigma_2$ -definable universal finite set for rank-extensions

$N \supseteq M$  is a **rank-extension** of  $M$  if every  $b \in N \setminus M$  has rank in  $N \setminus M$ .

## Theorem (Hamkins and Woodin)

*There is a  $\Sigma_2$  definition for a finite set  $\{b_0, \dots, b_n\}$  with the following properties.*

- 1 ZFC proves that the definition defines a finite set.
- 2 In any transitive model of ZFC the set is empty.
- 3 If  $M \models \text{ZFC}$  is countable, has  $s$  as its universal finite set, and  $t \in M$  is a finite set extending  $s$ , then there is  $N \models \text{ZFC}$  a rank-extension of  $M$  which has  $t$  as its universal finite set.

## Theorem (Hamkins and Woodin)

*The modal validities of rank-extensional set theoretic potentialism are precisely S4.*

# End-extensions versus rank-extensions

- $N \supseteq M$  is an **end-extension** of  $M$  if  $a \in b$  in  $N$  and  $b$  in  $M$  implies  $a$  is in  $M$ .
- Elementary end-extensions are always rank-extensions.
- But not all end-extensions are rank-extensions. For example, if  $M$  is an inner model of  $N$  then  $N$  end-extends  $M$ .

# End-extensions versus rank-extensions

- $N \supseteq M$  is an **end-extension** of  $M$  if  $a \in b$  in  $N$  and  $b$  in  $M$  implies  $a$  is in  $M$ .
- Elementary end-extensions are always rank-extensions.
- But not all end-extensions are rank-extensions. For example, if  $M$  is an inner model of  $N$  then  $N$  end-extends  $M$ .
- Key fact: the assertions which are preserved in arbitrary end-extensions are the  $\Sigma_1$  assertions.

$$\exists y \underbrace{\varphi(x, y)}_{\text{quantifiers bounded}}$$



# The $\Sigma_1$ -definable universal finite sequence for end-extensions

Let  $ZF^+$  be a computably enumerable extension of Zermelo–Fraenkel set theory  $ZF$ .

## Theorem (Hamkins, Welch, W.)

*There is a  $\Sigma_1$  definition for a finite sequence  $b_0, b_1, \dots, b_n$  with the following properties.*

- 1  $ZF^+$  proves that the sequence is finite.
- 2 In any standard model of  $ZF^+$  the sequence is finite.
- 3 Let  $M$  be a countable model of  $ZF^+$  which defines the sequence as  $s$ . Then if  $t$  in  $M$  is any finite sequence extending  $s$ , there is an end-extension  $N$  of  $M$  in which the universal sequence is  $t$ .

# The $\Sigma_1$ -definable universal finite sequence for end-extensions

Let  $ZF^+$  be a computably enumerable extension of Zermelo–Fraenkel set theory  $ZF$ .

## Theorem (Hamkins, Welch, W.)

*There is a  $\Sigma_1$  definition for a finite sequence  $b_0, b_1, \dots, b_n$  with the following properties.*

- ❶  $ZF^+$  proves that the sequence is finite.
- ❷ In any standard model of  $ZF^+$  the sequence is finite.
- ❸ Let  $M$  be a countable model of  $ZF^+$  which defines the sequence as  $s$ . Then if  $t$  in  $M$  is any finite sequence extending  $s$ , there is an end-extension  $N$  of  $M$  in which the universal sequence is  $t$ .
- ❹ Indeed, it suffices in (3) that  $M$  has an inner model  $W$  of  $ZF^+$  satisfying such.

# The definition of the universal finite sequence—Process A

Intended for  $\omega$ -nonstandard models. A different process is used for  $\omega$ -standard models.

- Proceed in stages to produce  $b_0, b_1, \dots, b_n$ , using auxiliary information: countable ordinals  $\alpha_0 < \alpha_1 < \dots < \alpha_n$  and natural numbers  $k_0 > k_1 > \dots > k_n$ .

# The definition of the universal finite sequence—Process A

Intended for  $\omega$ -nonstandard models. A different process is used for  $\omega$ -standard models.

- Proceed in stages to produce  $b_0, b_1, \dots, b_n$ , using auxiliary information: countable ordinals  $\alpha_0 < \alpha_1 < \dots < \alpha_n$  and natural numbers  $k_0 > k_1 > \dots > k_n$ .
- At stage  $n$ : Are there  $\alpha > \alpha_{n-1}$ ,  $k < k_{n-1}$ , and  $b \in L_{\alpha_n}$  so that  $L_{\alpha_n}$  has **no** end-extension to  $N$  satisfying the first  $k$  axioms of ZF plus “process A succeeds at stage  $n$  and defines  $b$ ”? If so, stage  $n$  is successful and set  $(b_n, \alpha_n, k_n)$  to be the  $L$ -least triple of such  $(b, \alpha, k)$ .

# The definition of the universal finite sequence—Process A

Intended for  $\omega$ -nonstandard models. A different process is used for  $\omega$ -standard models.

- Proceed in stages to produce  $b_0, b_1, \dots, b_n$ , using auxiliary information: countable ordinals  $\alpha_0 < \alpha_1 < \dots < \alpha_n$  and natural numbers  $k_0 > k_1 > \dots > k_n$ .
- At stage  $n$ : Are there  $\alpha > \alpha_{n-1}$ ,  $k < k_{n-1}$ , and  $b \in L_{\alpha_n}$  so that  $L_{\alpha_n}$  has **no** end-extension to  $N$  satisfying the first  $k$  axioms of ZF plus “process A succeeds at stage  $n$  and defines  $b$ ”? If so, stage  $n$  is successful and set  $(b_n, \alpha_n, k_n)$  to be the  $L$ -least triple of such  $(b, \alpha, k)$ .
- This self-reference is allowed by the **Gödel–Carnap fixed-point lemma**.

# The definition of the universal finite sequence—Process A

Intended for  $\omega$ -nonstandard models. A different process is used for  $\omega$ -standard models.

- Proceed in stages to produce  $b_0, b_1, \dots, b_n$ , using auxiliary information: countable ordinals  $\alpha_0 < \alpha_1 < \dots < \alpha_n$  and natural numbers  $k_0 > k_1 > \dots > k_n$ .
- At stage  $n$ : Are there  $\alpha > \alpha_{n-1}$ ,  $k < k_{n-1}$ , and  $b \in L_{\alpha_n}$  so that  $L_{\alpha_n}$  has **no** end-extension to  $N$  satisfying the first  $k$  axioms of ZF plus “process A succeeds at stage  $n$  and defines  $b$ ”? If so, stage  $n$  is successful and set  $(b_n, \alpha_n, k_n)$  to be the  $L$ -least triple of such  $(b, \alpha, k)$ .
- This self-reference is allowed by the **Gödel–Carnap fixed-point lemma**.
- Claim: The map  $n \mapsto (b_n, \alpha_n, k_n)$  is  $\Sigma_1$ -definable.
- Claim: Each  $k_i$  must be nonstandard.

# Seeing that Process $A$ has the plus one extension property

Consider countable  $M$  in which the universal sequence is  $b_0, \dots, b_{n-1}$  and take any  $b \in M$  and nonstandard  $k < k_{n-1}$ .

- Because stage  $n$  is unsuccessful in  $M$  this means  $L^M$  thinks every countable set can be end-extended to a model of the first  $k$  axioms of ZF in which the universal sequence is  $b_0, \dots, b_{n-1}, b$ .

# Seeing that Process $A$ has the plus one extension property

Consider countable  $M$  in which the universal sequence is  $b_0, \dots, b_{n-1}$  and take any  $b \in M$  and nonstandard  $k < k_{n-1}$ .

- Because stage  $n$  is unsuccessful in  $M$  this means  $L^M$  thinks every countable set can be end-extended to a model of the first  $k$  axioms of ZF in which the universal sequence is  $b_0, \dots, b_{n-1}, b$ .
- This is a  $\Pi_2^1$  assertion, so by the [Shoenfield absoluteness theorem](#) is true in  $M$ . It is also true in any elementary end-extensions of  $M$ , as well as their forcing extensions.



# Seeing that Process $A$ has the plus one extension property

Consider countable  $M$  in which the universal sequence is  $b_0, \dots, b_{n-1}$  and take any  $b \in M$  and nonstandard  $k < k_{n-1}$ .

- Because stage  $n$  is unsuccessful in  $M$  this means  $L^M$  thinks every countable set can be end-extended to a model of the first  $k$  axioms of ZF in which the universal sequence is  $b_0, \dots, b_{n-1}, b$ .
- This is a  $\Pi_2^1$  assertion, so by the [Shoenfield absoluteness theorem](#) is true in  $M$ . It is also true in any elementary end-extensions of  $M$ , as well as their forcing extensions.
- By the [Keisler–Morley theorem](#), let  $M^+$  be an elementary end extension of  $M$  and fix an ordinal  $\theta$  in  $M^+ \setminus M$ . Consider  $M^+[G]$  the forcing extension in which  $V_\theta$  is collapsed to be countable.

# Seeing that Process $A$ has the plus one extension property

Consider countable  $M$  in which the universal sequence is  $b_0, \dots, b_{n-1}$  and take any  $b \in M$  and nonstandard  $k < k_{n-1}$ .

- Because stage  $n$  is unsuccessful in  $M$  this means  $L^M$  thinks every countable set can be end-extended to a model of the first  $k$  axioms of ZF in which the universal sequence is  $b_0, \dots, b_{n-1}, b$ .
- This is a  $\Pi_2^1$  assertion, so by the [Shoenfield absoluteness theorem](#) is true in  $M$ . It is also true in any elementary end-extensions of  $M$ , as well as their forcing extensions.
- By the [Keisler–Morley theorem](#), let  $M^+$  be an elementary end extension of  $M$  and fix an ordinal  $\theta$  in  $M^+ \setminus M$ . Consider  $M^+[G]$  the forcing extension in which  $V_\theta$  is collapsed to be countable.
- So  $V_\theta^M$  has in  $M^+[G]$  an end-extension  $N$  in which the universal sequence is  $b_0, \dots, b_{n-1}, b$ . But  $N$  is also an end-extension of  $M$ .

# Process $B$ —for $\omega$ -standard models

- Again go in stages: produce  $b_0, b_1, \dots, b_n$  using auxiliary information countable ordinals  $\alpha_0 < \alpha_1 < \dots < \alpha_n$  and countable ordinals  $\lambda_0 > \lambda_1 > \dots > \lambda_n$ .

# Process $B$ —for $\omega$ -standard models

- Again go in stages: produce  $b_0, b_1, \dots, b_n$  using auxiliary information countable ordinals  $\alpha_0 < \alpha_1 < \dots < \alpha_n$  and countable ordinals  $\lambda_0 > \lambda_1 > \dots > \lambda_n$ .
- At stage  $n$ : are there  $\alpha_n > \alpha_{n-1}$ ,  $\lambda < \lambda_n$ , and  $b \in L_{\alpha_n}$  so that  $L_{\alpha_n}$  has **no** end-extension to  $N$  satisfying  $\text{ZF}^+$  plus “process  $B$  succeeds at stage  $n$  and defines  $b$ ”, with the well-founded tree canonically representing this  $\Pi_1^1$  assertion having rank  $\lambda$ ? If so, stage  $n$  is successful and set  $(b_n, \alpha_n, \lambda_n)$  to be the  $L$ -least triple of such  $(b, \alpha, \lambda)$ .

# Process $B$ —for $\omega$ -standard models

- Again go in stages: produce  $b_0, b_1, \dots, b_n$  using auxiliary information countable ordinals  $\alpha_0 < \alpha_1 < \dots < \alpha_n$  and countable ordinals  $\lambda_0 > \lambda_1 > \dots > \lambda_n$ .
- At stage  $n$ : are there  $\alpha_n > \alpha_{n-1}$ ,  $\lambda < \lambda_n$ , and  $b \in L_{\alpha_n}$  so that  $L_{\alpha_n}$  has **no** end-extension to  $N$  satisfying  $\text{ZF}^+$  plus “process  $B$  succeeds at stage  $n$  and defines  $b$ ”, with the well-founded tree canonically representing this  $\Pi_1^1$  assertion having rank  $\lambda$ ? If so, stage  $n$  is successful and set  $(b_n, \alpha_n, \lambda_n)$  to be the  $L$ -least triple of such  $(b, \alpha, \lambda)$ .
- Claim: The map  $n \mapsto (b_n, \alpha_n, \lambda_n)$  is  $\Sigma_1$ -definable.
- Claim: Each  $\lambda_i$  must be nonstandard.

# Process $B$ —for $\omega$ -standard models

- Again go in stages: produce  $b_0, b_1, \dots, b_n$  using auxiliary information countable ordinals  $\alpha_0 < \alpha_1 < \dots < \alpha_n$  and countable ordinals  $\lambda_0 > \lambda_1 > \dots > \lambda_n$ .
- At stage  $n$ : are there  $\alpha_n > \alpha_{n-1}$ ,  $\lambda < \lambda_n$ , and  $b \in L_{\alpha_n}$  so that  $L_{\alpha_n}$  has **no** end-extension to  $N$  satisfying  $\text{ZF}^+$  plus “process  $B$  succeeds at stage  $n$  and defines  $b$ ”, with the well-founded tree canonically representing this  $\Pi_1^1$  assertion having rank  $\lambda$ ? If so, stage  $n$  is successful and set  $(b_n, \alpha_n, \lambda_n)$  to be the  $L$ -least triple of such  $(b, \alpha, \lambda)$ .
- Claim: The map  $n \mapsto (b_n, \alpha_n, \lambda_n)$  is  $\Sigma_1$ -definable.
- Claim: Each  $\lambda_i$  must be nonstandard.
- Claim: Process  $B$  has the plus one extension property.

# Process $B$ —for $\omega$ -standard models

- Again go in stages: produce  $b_0, b_1, \dots, b_n$  using auxiliary information countable ordinals  $\alpha_0 < \alpha_1 < \dots < \alpha_n$  and countable ordinals  $\lambda_0 > \lambda_1 > \dots > \lambda_n$ .
- At stage  $n$ : are there  $\alpha_n > \alpha_{n-1}$ ,  $\lambda < \lambda_n$ , and  $b \in L_{\alpha_n}$  so that  $L_{\alpha_n}$  has **no** end-extension to  $N$  satisfying  $\text{ZF}^+$  plus “process  $B$  succeeds at stage  $n$  and defines  $b$ ”, with the well-founded tree canonically representing this  $\Pi_1^1$  assertion having rank  $\lambda$ ? If so, stage  $n$  is successful and set  $(b_n, \alpha_n, \lambda_n)$  to be the  $L$ -least triple of such  $(b, \alpha, \lambda)$ .
- Claim: The map  $n \mapsto (b_n, \alpha_n, \lambda_n)$  is  $\Sigma_1$ -definable.
- Claim: Each  $\lambda_i$  must be nonstandard.
- Claim: Process  $B$  has the plus one extension property.

Can merge Processes  $A$  and  $B$  into a single Process  $C$  which works for all models. □

# The Barwise extension theorem

The Barwise extension theorem can be derived as a corollary of our theorem.

## Theorem (Barwise)

*Every countable model of ZF end-extends to a model of  $ZFC + V = L$ .*



# The universal sequence for $L$ -extensions

## Corollary (Hamkins, Welch, W.)

*There is a  $\Sigma_1$  definition for a finite sequence  $b_0, b_1, \dots, b_n$  with the following properties.*

- ①  $ZF + V = L$  proves that the sequence is finite.
- ② In any standard model of  $ZF + V = L$  the sequence is finite.
- ③ Let  $M$  be a countable model of  $ZF + V = L$  which defines the sequence as  $s$ . Then if  $t$  in  $M$  is any finite sequence extending  $s$ , there is an  $L$ -extension  $N$  of  $M$  in which the universal sequence is  $t$ .


# Railyard labelings

- A **tree** is a partial order  $T$  so that  $\{s \in T : s \leq t\}$  is well-ordered for every  $t \in T$ . A **pre-tree** is a pre-order which quotients to a tree.
- A **railyard labeling** of a pre-tree  $T$  is an assignment  $\rho_t$  of statements to nodes  $t \in T$  so that each structure satisfies exactly one  $\rho_t$  and  $\diamond \rho_s$  holds iff  $t \leq_T s$ .

# Railyard labelings


- A **tree** is a partial order  $T$  so that  $\{s \in T : s \leq t\}$  is well-ordered for every  $t \in T$ . A **pre-tree** is a pre-order which quotients to a tree.
- A **railyard labeling** of a pre-tree  $T$  is an assignment  $\rho_t$  of statements to nodes  $t \in T$  so that each structure satisfies exactly one  $\rho_t$  and  $\Diamond \rho_s$  holds iff  $t \leq_T s$ .
- If there are railyard labelings for every finite pre-tree, then the modal validities for the corresponding potentialist system are contained within S4.

# The universal finite sequence and railyard labelings




# The universal finite sequence and railyard labelings

- Step 1: the subsequence  $\langle n_i \rangle$  of finite ordinals from the universal finite sequence tell you how to descend the tree to determine your cluster. If  $B$  is the branching of the current node, then  $n_i \bmod B$  tells you where to go.



# The universal finite sequence and railyard labelings

- Step 1: the subsequence  $\langle n_i \rangle$  of finite ordinals from the universal finite sequence tell you how to descend the tree to determine your cluster. If  $B$  is the branching of the current node, then  $n_i \bmod B$  tells you where to go.
- Step 2: the final infinite ordinal  $\lambda + m$  on the sequence tells you where in your cluster you are. If  $K$  is the size of the cluster, then  $m \bmod K$  identifies your node in the cluster. (If no infinite ordinals are on the sequence, default to 0.)



# The modal validities of end-extensional set theoretic potentialism

## Theorem (Hamkins, Welch, W.)

*Consider the potentialist system consisting of countable models of  $ZF^+$  ordered by end-extension.*

- ① *For any world  $M$ , the modal validities, allowing for a single parameter for the length of the universal finite sequence, are precisely S4.*
- ② *For any  $\omega$ -standard world  $M$ , the modal validities, allowing no parameters, are precisely S4.*

# Thank you!