
Table of Contents

.....	1
QUESTION 1:	1
QUESTION 2:	1
QUESTION 3:	2
QUESTION 4:	5
QUESTION 5:	10
QUESTION 6:	16
QUESTION 7:	20
QUESTION 8:	20
ALL FUNCTIONS SUPPORTING THIS CODE	20

```
clear;
```

QUESTION 1:

```
% Function is at the bottom in the supporting code section
% function out = f(theta)

% Testing theta = pi/4
f(pi/4)

% Testing theta = -pi/4
f(-pi/4)

% Both are close to 0, so we are good
```


QUESTION 2:

```
% Plotting f(theta) on [-pi, pi]
theta_vals = -pi:0.01:pi;

f_vals = f(theta_vals);

figure(1)
plot(theta_vals, f_vals)
xlabel('\theta (radians)')
ylabel('f(\theta)')
title('Plot of f(\theta) on [-\pi, \pi]')
yline(0, '--r');
xline(pi/4, '--g', '\pi/4');
xline(-pi/4, '--g', '-\pi/4');
drawnow;

% Plot clearly shows that there are roots at +/- pi/4
```



QUESTION 3:

```
% Pose from Figure 1.15 (a)

% Connected to (0, 0) aka (x, y)
u1 = 1; v1 = 2;

% Connected to (x1, 0)
u2 = 2; v2 = 1;

% Connected to (x2, y2)
u3 = 2; v3 = 3;

x1 = 4; x2 = 0; y2 = 4;

figure(2)
plot([u1 u2 u3 u1], [v1 v2 v3 v1], 'r'); hold on % Platform triangle
plot([0 x1 x2], [0 0 y2], 'bo') % Base anchors
plot([u1 u2 u3], [v1 v2 v3], 'ro', 'MarkerSize', 8, 'MarkerFaceColor', 'r')
% Platform joints
plot([u1 0], [v1 0], 'k--') % p1
plot([u2 x1], [v2 0], 'k--') % p2
plot([u3 x2], [v3 y2], 'k--') % p3
title('Figure 1.15 (a)')
xlabel('x')
```

```

ylabel('y')
drawnow;

% Pose from Figure 1.15 (b)

% Connected to (0, 0) aka (x, y)
u1 = 2; v1 = 1;

% Connected to (x1, 0)
u2 = 3; v2 = 2;



% Connected to (x2, y2)
u3 = 1; v3 = 2;

x1 = 4; x2 = 0; y2 = 4;

figure(3)
plot([u1 u2 u3 u1], [v1 v2 v3 v1], 'r'); hold on % Platform triangle
plot([0 x1 x2], [0 0 y2], 'bo') % Base anchors
plot([u1 u2 u3], [v1 v2 v3], 'ro', 'MarkerSize', 8, 'MarkerFaceColor', 'r')
% Platform joints
plot([u1 0], [v1 0], 'k--') % p1
plot([u2 x1], [v2 0], 'k--') % p2
plot([u3 x2], [v3 y2], 'k--') % p3
title('Figure 1.15 (b)')
xlabel('x')
ylabel('y')
drawnow;

% Here, we're just reproducing Figure 1.15 (a) and (b)

```



QUESTION 4:

```
% Forward kinematics is when we compute (x, y) and theta for each given p1,
% p2, and p3

% The inverse kinematic problem is when we find p1, p2, p3, given x, y, and
% theta

% The new f(theta) function is in the supporting functions section at the
% bottom (f_4(theta))

% Plotting f_4(theta) on [-pi, pi]
theta_vals = -pi:0.01:pi;

f_vals = f_variable_p2(theta_vals, 5); % p2=5

figure(4)
plot(theta_vals, f_vals)
xlabel('\theta (radians)')
ylabel('f(\theta)')
title('Plot of f(\theta) on [-\pi, \pi]')
yline(0, '--r');
drawnow;

% Finding the four theta values (guesses are from eyeballing the graph)
p2 = 5;
f_p2 = @(theta) f_variable_p2(theta, p2);

theta1 = fzero(f_p2, -0.72);
theta2 = fzero(f_p2, -0.33);
theta3 = fzero(f_p2, 1.14);
theta4 = fzero(f_p2, 2.11);
thetas = [theta1 theta2 theta3 theta4];

% From the above it appears that our roots are at:
% theta = -0.7208, -0.3310, 1.1437, and 2.1159 radians
figure(5)
plot(theta_vals, f_vals)
xlabel('\theta (radians)')
ylabel('f(\theta)')
title('Plot of f(\theta) on [-\pi, \pi] with roots')
yline(0, '--r');
xline(theta1, '--r', '-0.7208');
xline(theta2, '--r', '-0.3310');
xline(theta3, '--r', '1.1437');
xline(theta4, '--r', '2.1159');
drawnow;

% Since we're asked to solve the forward kinematics problem, we need to
% solve for x and y now (we just solved for theta)

% Finding the x and y coordinates for the four poses
% Created a new function at the bottom called
```

```
% forward_kinematics_variable_p2
[x_1 y_1] = forward_kinematics_variable_p2(theta1, p2);
[x_2 y_2] = forward_kinematics_variable_p2(theta2, p2);
[x_3 y_3] = forward_kinematics_variable_p2(theta3, p2);
[x_4 y_4] = forward_kinematics_variable_p2(theta4, p2);
xs = [x_1 x_2 x_3 x_4];
ys = [y_1 y_2 y_3 y_4];



% It was found that
% (x_1, y_1) = (-1.3784, 4.8063)
% (x_2, y_2) = (-0.9147, 4.9156)
% (x_3, y_3) = (4.4818, 2.2167)
% (x_4, y_4) = (4.5718, 2.0244)



% Now we need to plot the four poses
% Helper function is in the supporting functions section



for i = 1:4
    draw_pose(5+i, xs(i), ys(i), thetas(i), 4, i);
    drawnow;
end

% The strut lengths are correct!!!

Pose 1: p1 = 5.0000, p2 = 5.0000, p3 = 3.0000
Pose 2: p1 = 5.0000, p2 = 5.0000, p3 = 3.0000
Pose 3: p1 = 5.0000, p2 = 5.0000, p3 = 3.0000
Pose 4: p1 = 5.0000, p2 = 5.0000, p3 = 3.0000
```







QUESTION 5:

```
% Here we are changing p2 to 7 and resolving problem 4

% Plotting f(theta) on [-pi, pi]
theta_vals = -pi:0.01:pi;

f_vals = f_variable_p2(theta_vals, 7); % p2=7

figure(10)
plot(theta_vals, f_vals)
xlabel('\theta (radians)')
ylabel('f(\theta)')
title('Plot of f(\theta) on [-\pi, \pi] for Question #5')
yline(0, '--r');
drawnow;

% The problem states that there are now six poses

% Finding the six theta values (guesses are from eyeballing the graph)
p2 = 7;
f_p2 = @(theta) f_variable_p2(theta, p2);

theta1 = fzero(f_p2, -0.68);
theta2 = fzero(f_p2, -0.36);
theta3 = fzero(f_p2, 0.03);
theta4 = fzero(f_p2, 0.44);
theta5 = fzero(f_p2, 0.97);
theta6 = fzero(f_p2, 2.5);

thetas = [theta1 theta2 theta3 theta4 theta5 theta6];

% From the above it appears that our roots are at:
% theta = -0.6732, -0.3547, 0.0378, 0.4589, 0.9777, and 2.5139 rad
figure(11)
plot(theta_vals, f_vals)
xlabel('\theta (radians)')
ylabel('f(\theta)')
title('Plot of f(\theta) on [-\pi, \pi] with roots')
yline(0, '--r');
xline(theta1, '--r', '-0.6732');
xline(theta2, '--r', '-0.3547');
xline(theta3, '--r', '0.0378');
xline(theta4, '--r', '0.4589');
xline(theta5, '--r', '0.9777');
xline(theta6, '--r', '2.5139');
drawnow;

% Since we're asked to solve the forward kinematics problem, we need to
% solve for x and y now (we just solved for theta)

% Finding the x and y coordinates for the four poses
[x_1 y_1] = forward_kinematics_variable_p2(theta1, p2);
```

```

[x_2 y_2] = forward_kinematics_variable_p2(theta2, p2);
[x_3 y_3] = forward_kinematics_variable_p2(theta3, p2);
[x_4 y_4] = forward_kinematics_variable_p2(theta4, p2);
[x_5 y_5] = forward_kinematics_variable_p2(theta5, p2);
[x_6 y_6] = forward_kinematics_variable_p2(theta6, p2);

xs = [x_1 x_2 x_3 x_4 x_5 x_6];
ys = [y_1 y_2 y_3 y_4 y_5 y_6];

% It was found that
% (x_1, y_1) = (-4.3148, 2.5264)
% (x_2, y_2) = (-4.8049, 1.3831)
% (x_3, y_3) = (-4.9490, 0.7121)
% (x_4, y_4) = (-0.8198, 4.9323)
% (x_5, y_6) = (2.3036, 4.4378)
% (x_5, y_6) = (3.2157, 3.8287)



% Now we need to plot the four poses
% Helper function is in the supporting functions section



for i = 1:6
    draw_pose(11+i, xs(i), ys(i), thetas(i), 5, i);
    drawnow;
end



% The strut lengths are correct!!!



Pose 1: p1 = 5.0000, p2 = 7.0000, p3 = 3.0000
Pose 2: p1 = 5.0000, p2 = 7.0000, p3 = 3.0000
Pose 3: p1 = 5.0000, p2 = 7.0000, p3 = 3.0000
Pose 4: p1 = 5.0000, p2 = 7.0000, p3 = 3.0000
Pose 5: p1 = 5.0000, p2 = 7.0000, p3 = 3.0000
Pose 6: p1 = 5.0000, p2 = 7.0000, p3 = 3.0000


```











QUESTION 6:

```
% Now we need to find a strut length p2, for which there are only two
% poses. It is found that when p2=4, there are only two poses

theta_vals = -pi:0.01:pi;
p2_range = 1:7; % Test p2 values between 1 and 7
target_num_roots = 2;
found = false;

for p2 = p2_range
    f_vals = f_variable_p2(theta_vals, p2);
    sign_changes = sum(abs(diff(sign(f_vals))) == 2); % # of times f(theta)
crosses 0
    if sign_changes == target_num_roots
        fprintf("Found p2 = %.2f with exactly %d poses\n", p2,
target_num_roots);
        found = true;
        break
    end
end

if ~found
    fprintf("No p2 in range [%0.2f, %0.2f] gives exactly %d poses\n", ...
        p2_range(1), p2_range(end), target_num_roots);
```

```

end

% Here we are changing p2 to 4

% Plotting f(theta) on [-pi, pi]
theta_vals = -pi:0.01:pi;

f_vals = f_variable_p2(theta_vals, 4); % p2=4

figure(18)
plot(theta_vals, f_vals)
xlabel('\theta (radians)')
ylabel('f(\theta)')
title('Plot of f(\theta) on [-\pi, \pi] for Question #6')
yline(0, '--r');
drawnow;

% Finding the six theta values (guesses are from eyeballing the graph)
p2 = 4;
f_p2 = @(theta) f_variable_p2(theta, p2);

theta1 = fzero(f_p2, 1.32);
theta2 = fzero(f_p2, 1.77);

thetas = [theta1 theta2];

% theta vals are 1.3316 and 1.7775 rad

% Since we're asked to solve the forward kinematics problem, we need to
% solve for x and y now (we just solved for theta)

% Finding the x and y coordinates for the four poses
[x_1 y_1] = forward_kinematics_variable_p2(theta1, p2);
[x_2 y_2] = forward_kinematics_variable_p2(theta2, p2);

xs = [x_1 x_2];
ys = [y_1 y_2];



% It was found that
% (x_1, y_1) = (4.8907, 1.0399)
% (x_2, y_2) = (4.8992, 0.9992)



% Now we need to plot the four poses
% Helper function is in the supporting functions section

for i = 1:2
    draw_pose(18+i, xs(i), ys(i), thetas(i), 6, i);
    drawnow;
end

Found p2 = 4.00 with exactly 2 poses
Pose 1: p1 = 5.0000, p2 = 4.0000, p3 = 3.0000
Pose 2: p1 = 5.0000, p2 = 4.0000, p3 = 3.0000

```





QUESTION 7:

QUESTION 8:

ALL FUNCTIONS SUPPORTING THIS CODE

```
% First f(theta) function
function out = f(theta)

    % Platform lengths
    L1 = 2;
    L2 = sqrt(2);
    L3 = sqrt(2);

    % Angle across from L1
    gamma = pi / 2;

    % Strut lengths
    p1 = sqrt(5);
    p2 = sqrt(5);
    p3 = sqrt(5);

    % Strut base positions
    % Got these from Figure 1.15
    x1 = 4;
    x2 = 0;
    y2 = 4;

    A2 = L3 * cos(theta) - x1;
    B2 = L3 * sin(theta);
    A3 = L2 * (cos(theta) * cos(gamma) - sin(theta) * sin(gamma)) - x2;
    B3 = L2 * (cos(theta) * sin(gamma) + sin(theta) * cos(gamma)) - y2;

    N1 = B3 .* (p2^2 - p1^2 - A2.^2 - B2.^2) - B2 .* (p3^2 - p1^2 - A3.^2 - B3.^2);
    N2 = -A3 .* (p2^2 - p1^2 - A2.^2 - B2.^2) + A2 .* (p3^2 - p1^2 - A3.^2 - B3.^2);

    D = 2 * (A2 .* B3 - B2 .* A3);

    out = N1.^2 + N2.^2 - p1.^2 * D.^2;

end

% f(theta) function with ability to change p2
function out = f_variable_p2(theta, p2)
    L1 = 3; L2 = 3 * sqrt(2); L3 = 3;
    gamma = pi / 4;
    p1 = 5; p3 = 3;
    x1 = 5; x2 = 0; y2 = 6;
```

```

    A2 = L3 * cos(theta) - x1;
    B2 = L3 * sin(theta);
    A3 = L2 * (cos(theta) * cos(gamma) - sin(theta) * sin(gamma)) - x2;
    B3 = L2 * (cos(theta) * sin(gamma) + sin(theta) * cos(gamma)) - y2;

    N1 = B3 .* (p2^2 - p1^2 - A2.^2 - B2.^2) - B2 .* (p3^2 - p1^2 - A3.^2 -
B3.^2);
    N2 = -A3 .* (p2^2 - p1^2 - A2.^2 - B2.^2) + A2 .* (p3^2 - p1^2 - A3.^2 -
B3.^2);
    D = 2 * (A2 .* B3 - B2 .* A3);

    out = N1.^2 + N2.^2 - p1.^2 * D.^2;
end

% Forward kinematics problem solver with variable p2
function [x, y] = forward_kinematics_variable_p2(theta, p2)

    % Platform lengths
    L1 = 3;
    L2 = 3 * sqrt(2);
    L3 = 3;

    % Angle across from L1
    gamma = pi / 4;

    % Strut lengths
    p1 = 5;
    p3 = 3;

    % Strut base positions
    x1 = 5;
    x2 = 0;
    y2 = 6;

    % Compute intermediate terms
    A2 = L3 * cos(theta) - x1;
    B2 = L3 * sin(theta);
    A3 = L2 * (cos(theta) * cos(gamma) - sin(theta) * sin(gamma)) - x2;
    B3 = L2 * (cos(theta) * sin(gamma) + sin(theta) * cos(gamma)) - y2;

    % Numerators and denominator
    N1 = B3 .* (p2^2 - p1^2 - A2.^2 - B2.^2) - B2 .* (p3^2 - p1^2 - A3.^2 -
B3.^2);
    N2 = -A3 .* (p2^2 - p1^2 - A2.^2 - B2.^2) + A2 .* (p3^2 - p1^2 - A3.^2 -
B3.^2);
    D = 2 * (A2 .* B3 - B2 .* A3);

    % Solve for x and y
    x = N1 / D;
    y = N2 / D;

end

```

```

function draw_pose(fig_num, x, y, theta, question_number, pose_index)
    % Constants
    L2 = 3 * sqrt(2);
    L3 = 3;
    gamma = pi/4;
    x1 = 5; x2 = 0; y2 = 6;

    % Triangle corner positions
    u1 = x;
    v1 = y;
    u2 = x + L3 * cos(theta);
    v2 = y + L3 * sin(theta);
    u3 = x + L2 * cos(theta + gamma);
    v3 = y + L2 * sin(theta + gamma);

    % Compute strut lengths
    p1 = norm([u1, v1] - [0, 0]);
    p2 = norm([u2, v2] - [x1, 0]);
    p3 = norm([u3, v3] - [x2, y2]);

    % Plot
    figure(fig_num)
    plot([u1 u2 u3 u1], [v1 v2 v3 v1], 'r'); hold on
    plot([0 x1 x2], [0 0 y2], 'bo')
    plot([u1 u2 u3], [v1 v2 v3], 'ro', 'MarkerSize', 8, 'MarkerFaceColor',
'r')
    plot([u1 0], [v1 0], 'k--') % p1
    plot([u2 x1], [v2 0], 'k--') % p2
    plot([u3 x2], [v3 y2], 'k--') % p3

    % Pose label
    title_str = sprintf('Pose %d (\\theta = %.2f)', pose_index, theta);

    title(title_str)
    xlabel('x')
    ylabel('y')
    axis equal
    grid on

    % Print strut lengths
    fprintf("Pose %d: p1 = %.4f, p2 = %.4f, p3 = %.4f\\n", pose_index, p1,
p2, p3);
end

```

ans =

-4.5475e-13

ans =

$-4.5475e-13$

Published with MATLAB® R2024b