
Table of Contents

QUESTION 1 COMMENTING	1
QUESTION 1:	1
QUESTION 2:	1
QUESTION 3:	2
QUESTION 4:	2
QUESTION 5:	2
QUESTION 6:	2
QUESTION 7:	2
QUESTION 8:	2
ALL FUNCTIONS SUPPORTING THIS CODE	2

QUESTION 1 COMMENTING

```
clear; close all; clc;
```

QUESTION 1:

```
% Function is at the bottom!!! function out = f(theta)
```

```
% Testing theta = pi/4  
disp(f(pi/4))
```

```
% Testing theta = -pi/4  
disp(f(-pi/4))
```

```
% Both are close to 0, so we are good
```

QUESTION 2:

```
% Plotting f(theta) on [-pi, pi]  
theta_vals = -pi:0.01:pi;
```

```
f_vals = f(theta_vals);
```

```
figure(1)  
plot(theta_vals, f_vals)  
xlabel('\theta (radians)')  
ylabel('f(\theta)')  
title('Plot of f(\theta) on [-\pi, \pi]')  
yline(0, '--r');  
xline(pi/4, '--g', '\pi/4');  
xline(-pi/4, '--g', '-\pi/4');
```

```
% Plot clearly shows that there are roots at +/- pi/4
```

QUESTION 3:

QUESTION 4:

QUESTION 5:

QUESTION 6:

QUESTION 7:

QUESTION 8:

ALL FUNCTIONS SUPPORTING THIS CODE

```
function out = f(theta)

    % Platform lengths
    L1 = 2;
    L2 = sqrt(2);
    L3 = sqrt(2);

    % Angle across from L1
    gamma = pi / 2;

    % Strut lengths
    p1 = sqrt(5);
    p2 = sqrt(5);
    p3 = sqrt(5);

    % Strut base positions
    % Got these from Figure 1.15
    x1 = 4;
    x2 = 0;
    y2 = 4;

    A2 = L3 * cos(theta) - x1;
    B2 = L3 * sin(theta);
    A3 = L2 * (cos(theta) * cos(gamma) - sin(theta) * sin(gamma)) - x2;
    B3 = L2 * (cos(theta) * sin(gamma) + sin(theta) * cos(gamma)) - y2;

    N1 = B3 .* (p2^2 - p1^2 - A2.^2 - B2.^2) - B2 .* (p3^2 - p1^2 - A3.^2 - B3.^2);
    N2 = -A3 .* (p2^2 - p1^2 - A2.^2 - B2.^2) + A2 .* (p3^2 - p1^2 - A3.^2 - B3.^2);

    D = 2 * (A2 .* B3 - B2 .* A3);

    out = N1.^2 + N2.^2 - p1.^2 * D.^2;
```

end

-4.5475e-13

-4.5475e-13

Published with MATLAB® R2024b