
Table of Contents

QUESTION 1:	1
QUESTION 2:	1
QUESTION 3:	2
QUESTION 4:	4
QUESTION 5:	4
QUESTION 6:	4
QUESTION 7:	4
QUESTION 8:	4
ALL FUNCTIONS SUPPORTING THIS CODE	4

QUESTION 1:

```
% Function is at the bottom!!! function out = f(theta)

% Testing theta = pi/4
disp(f(pi/4))

% Testing theta = -pi/4
disp(f(-pi/4))

% Both are close to 0, so we are good
```


QUESTION 2:

```
% Plotting f(theta) on [-pi, pi]
theta_vals = -pi:0.01:pi;

f_vals = f(theta_vals);

figure(1)
plot(theta_vals, f_vals)
xlabel('\theta (radians)')
ylabel('f(\theta)')
title('Plot of f(\theta) on [-\pi, \pi]')
yline(0, '--r');
xline(pi/4, '--g', '\pi/4');
xline(-pi/4, '--g', '-\pi/4');
drawnow;

% Plot clearly shows that there are roots at +/- pi/4
```



QUESTION 3:

```
% Pose from Figure 1.15 (a)
```

```
u1 = 1; u2 = 2; u3 = 2;
```

```
v1 = 2; v2 = 1; v3 = 3;
```

```
x1 = 4; x2 = 0; y2 = 4;
```

```
figure(2)
```

```
plot([u1 u2 u3 u1], [v1 v2 v3 v1], 'r'); hold on % Platform triangle
```

```
plot([0 x1 x2], [0 0 y2], 'bo') % Base anchors
```

```
plot([u1 u2 u3], [v1 v2 v3], 'ro', 'MarkerSize', 8, 'MarkerFaceColor', 'r')
```

```
% Platform joints
```

```
plot([u1 0], [v1 0], 'k--') % Strut 1
```

```
plot([u2 x1], [v2 0], 'k--') % Strut 2
```

```
plot([u3 x2], [v3 y2], 'k--') % Strut 3
```

```
title('Figure 1.15 (a)')
```

```
xlabel('x')
```

```
ylabel('y')
```

```
drawnow;
```

```
% Pose from Figure 1.15 (b)
```

```
u1 = 2; u2 = 3; u3 = 1;
```

```
v1 = 1; v2 = 2; v3 = 2;
```


```
x1 = 4; x2 = 0; y2 = 4;
```


```

figure(3)
plot([u1 u2 u3 u1], [v1 v2 v3 v1], 'r'); hold on           % Platform triangle
plot([0 x1 x2], [0 0 y2], 'bo')                             % Base anchors
plot([u1 u2 u3], [v1 v2 v3], 'ro', 'MarkerSize', 8, 'MarkerFaceColor', 'r')
% Platform joints
plot([u1 0], [v1 0], 'k--') % Strut 1
plot([u2 x1], [v2 0], 'k--') % Strut 2
plot([u3 x2], [v3 y2], 'k--') % Strut 3
title('Figure 1.15 (b)')
xlabel('x')
ylabel('y')
drawnow;

% Here, we're just reproducing Figure 1.15 (a) and (b)

```





QUESTION 4:

QUESTION 5:

QUESTION 6:

QUESTION 7:

QUESTION 8:

ALL FUNCTIONS SUPPORTING THIS CODE

```
function out = f(theta)

    % Platform lengths
    L1 = 2;
    L2 = sqrt(2);
    L3 = sqrt(2);

    % Angle across from L1
    gamma = pi / 2;
```

```

% Strut lengths
p1 = sqrt(5);
p2 = sqrt(5);
p3 = sqrt(5);

% Strut base positions
% Got these from Figure 1.15
x1 = 4;
x2 = 0;
y2 = 4;

A2 = L3 * cos(theta) - x1;
B2 = L3 * sin(theta);
A3 = L2 * (cos(theta) * cos(gamma) - sin(theta) * sin(gamma)) - x2;
B3 = L2 * (cos(theta) * sin(gamma) + sin(theta) * cos(gamma)) - y2;

N1 = B3 .* (p2^2 - p1^2 - A2.^2 - B2.^2) - B2 .* (p3^2 - p1^2 - A3.^2 -
B3.^2);
N2 = -A3 .* (p2^2 - p1^2 - A2.^2 - B2.^2) + A2 .* (p3^2 - p1^2 - A3.^2 -
B3.^2);

D = 2 * (A2 .* B3 - B2 .* A3);

out = N1.^2 + N2.^2 - p1.^2 * D.^2;

end

-4.5475e-13

-4.5475e-13

```

Published with MATLAB® R2024b