Rational Speech Act models are utterance-independent updates of world priors

Jean-Philippe Bernardy, Julian Grove, and Christine Howes Semdial 2022 - DubDial, August 23, 2022

CLASP, University of Gothenburg

Introduction to RSA



u = 'JP ate five cookies.'

$$[u] = n_{cookies} \ge 5$$



$$u = \text{'JP ate five cookies.'}$$

 $||u|| = n_{cookies} \ge 5$

Three components:



u = 'JP ate five cookies.' $||u|| = n_{cookies} \ge 5$

Three components:

• A literal listener L_0 : $u \mapsto P_{L_0}(w|u)$



$$u = \text{'JP ate five cookies.'}$$

 $||u|| = n_{cookies} \ge 5$

Three components:

- A literal listener L_0 : $u \mapsto P_{L_0}(w|u)$
- A pragmatic speaker S_1 : $w \mapsto P_{S_1}(u|w)$



$$u = \text{'JP ate five cookies.'}$$

 $||u|| = n_{cookies} \ge 5$

Three components:

- A literal listener L_0 : $u \mapsto P_{L_0}(w|u)$
- A pragmatic speaker S_1 : $w \mapsto P_{S_1}(u|w)$
- A pragmatic listener L_1 : $u \mapsto P_{L_1}(w|u)$

The literal listener L_0

$$P_{L_0}(w|u) \propto \mathbb{1}(w \geq n) \times P(w)$$

W	5	6	7
'JP ate 5 cookies'	1/3	1/3	1/3
'JP ate 6 cookies'	0	1/2	1/2
'JP ate 7 cookies'	0	0	1

The pragmatic speaker S_1

$$P_{S_1}(u \mid w) \propto \frac{P_{L_0}(w \mid u)^{\alpha}}{e^{\alpha \times C(u)}}$$

(Assume $\alpha = 1$, and that C(u) is constant.)

W	5	6	7
'JP ate 5 cookies'	1	0.16	0.01
'JP ate 6 cookies'	0	0.84	0.06
'JP ate 7 cookies'	0	0	0.93

The pragmatic listener L_1

$$P_{L_1}(w|u) \propto P_{S_1}(u|w) \times P(w)$$

W	5	6	7
'JP ate 5 cookies'	0.85	0.14	0.01
'JP ate 6 cookies'	0	0.93	0.07
'JP ate 7 cookies'	0	0	1

• Reformulate the components of RSA as *filters* on reconceived prior distributions (using information gain).

- Reformulate the components of RSA as *filters* on reconceived prior distributions (using information gain).
- Discuss some consequences:

- Reformulate the components of RSA as *filters* on reconceived prior distributions (using information gain).
- Discuss some consequences:
 - for computing RSA models;

- Reformulate the components of RSA as *filters* on reconceived prior distributions (using information gain).
- Discuss some consequences:
 - · for computing RSA models;
 - for the algorithmic plausibility of RSA (Marr, 1982).

RSA via information gain

Information gain as K-L divergence

Kullback-Leibler (K-L) divergence (between distributions P and Q over X):

$$D_{\mathsf{KL}}(Q \parallel P) = -\sum_{x \in \mathcal{X}} Q(x) \log \left(\frac{P(x)}{Q(x)} \right)$$

Information gain as K-L divergence

Kullback-Leibler (K-L) divergence (between distributions P and Q over X):

$$D_{KL}(Q \parallel P) = -\sum_{x \in \mathcal{X}} Q(x) \log \left(\frac{P(x)}{Q(x)} \right)$$

Higher values: more information is gained by going from P to Q.

A theorem

Theorem

If $Q(x) \propto f(x) \times P(x)$, and the range of f is $\{0, 1\}$, then:

$$D_{\text{KL}}(Q \parallel P) = -\log \left(\sum_{x} f(x) \times P(x) \right)$$

-

A theorem

Theorem

If $Q(x) \propto f(x) \times P(x)$, and the range of f is $\{0, 1\}$, then:

$$D_{\mathsf{KL}}(Q \parallel P) = -\log \left(\sum_{x} f(x) \times P(x) \right)$$

In words: if f updates P merely by *filtering* it (resulting in Q), then the information gain of Q is the negative log of the expected value of f.

A theorem

Theorem

If $Q(x) \propto f(x) \times P(x)$, and the range of f is $\{0, 1\}$, then:

$$D_{\mathsf{KL}}(Q \parallel P) = -\log \left(\sum_{x} f(x) \times P(x) \right)$$

In words: if f updates P merely by *filtering* it (resulting in Q), then the information gain of Q is the negative log of the expected value of f.

• Important: the expected value of *f* is *also* the normalizing constant for *Q*:

$$Q(x) = \frac{f(x) \times P(x)}{\sum_{x'} f(x') \times P(x')}$$

$$P_{L_0}(w|u) = \frac{l(u,w) \times P(w)}{\sum_{w' \in \mathcal{W}} l(u,w') \times P(w')}$$

where l(u, w) = 1 or 0, depending on whether u is true at w

$$P_{L_0}(w|u) = \frac{l(u,w) \times P(w)}{\sum_{w' \in \mathcal{W}} l(u,w') \times P(w')}$$

where l(u, w) = 1 or 0, depending on whether u is true at w

Let us define the *literal information gain* of *u*:

$$G_{L_0}(u) = D_{\mathsf{KL}}(P_{L_0}(w|u) \parallel P(w))$$

$$P_{L_0}(w|u) = \frac{l(u,w) \times P(w)}{\sum_{w' \in \mathcal{W}} l(u,w') \times P(w')}$$

where l(u, w) = 1 or 0, depending on whether u is true at w

Let us define the *literal information gain* of *u*:

$$G_{L_0}(u) = D_{KL}(P_{L_0}(w|u) \parallel P(w))$$

Using our theorem:

$$G_{L_0}(u) = -\log \sum_{w' \in \mathcal{W}} l(u, w') \times P(w')$$

$$P_{L_0}(w|u) = \frac{l(u,w) \times P(w)}{\sum_{w' \in \mathcal{W}} l(u,w') \times P(w')}$$

where l(u, w) = 1 or 0, depending on whether u is true at w

Let us define the *literal information gain* of *u*:

$$G_{L_0}(u) = D_{KL}(P_{L_0}(w|u) \parallel P(w))$$

Using our theorem:

$$G_{L_0}(u) = -\log \sum_{w' \in \mathcal{W}} l(u, w') \times P(w')$$

$$P_{L_0}(w|u) = l(u, w) \times P(w) \times e^{G_{L_0}(u)}$$

$$P_{S_1}(u \mid w) \propto \frac{P_{L_0}(w \mid u)^{\alpha}}{e^{\alpha \times C(u)}}$$

$$P_{S_1}(u \mid w) \propto \frac{P_{L_0}(w \mid u)^{\alpha}}{e^{\alpha \times C(u)}}$$

$$\propto \frac{\left(l(u, w) \times P(w) \times e^{G_{L_0}(u)}\right)^{\alpha}}{e^{\alpha \times C(u)}}$$

$$P_{S_{1}}(u \mid w) \propto \frac{P_{L_{0}}(w \mid u)^{\alpha}}{e^{\alpha \times C(u)}}$$

$$\propto \frac{\left(l(u, w) \times P(w) \times e^{G_{L_{0}}(u)}\right)^{\alpha}}{e^{\alpha \times C(u)}}$$

$$\propto l(u, w)^{\alpha} \times P(w)^{\alpha} \times e^{\alpha \times (G_{L_{0}}(u) - C(u))}$$

$$P_{S_{1}}(u \mid w) \propto \frac{P_{L_{0}}(w \mid u)^{\alpha}}{e^{\alpha \times C(u)}}$$

$$\propto \frac{\left(l(u, w) \times P(w) \times e^{G_{L_{0}}(u)}\right)^{\alpha}}{e^{\alpha \times C(u)}}$$

$$\propto l(u, w)^{\alpha} \times P(w)^{\alpha} \times e^{\alpha \times (G_{L_{0}}(u) - C(u))}$$

$$\propto l(u, w) \times e^{\alpha \times (G_{L_{0}}(u) - C(u))}$$

$$P_{S_{1}}(u \mid w) \propto \frac{P_{L_{0}}(w \mid u)^{\alpha}}{e^{\alpha \times C(u)}}$$

$$\propto \frac{\left(l(u, w) \times P(w) \times e^{G_{L_{0}}(u)}\right)^{\alpha}}{e^{\alpha \times C(u)}}$$

$$\propto l(u, w)^{\alpha} \times P(w)^{\alpha} \times e^{\alpha \times (G_{L_{0}}(u) - C(u))}$$

$$\propto l(u, w) \times e^{\alpha \times (G_{L_{0}}(u) - C(u))}$$

The pragmatic speaker S_1 :

$$P_{S_{1}}(u \mid w) \propto \frac{P_{L_{0}}(w \mid u)^{\alpha}}{e^{\alpha \times C(u)}}$$

$$\propto \frac{\left(l(u, w) \times P(w) \times e^{G_{L_{0}}(u)}\right)^{\alpha}}{e^{\alpha \times C(u)}}$$

$$\propto l(u, w)^{\alpha} \times P(w)^{\alpha} \times e^{\alpha \times (G_{L_{0}}(u) - C(u))}$$

$$\propto l(u, w) \times e^{\alpha \times (G_{L_{0}}(u) - C(u))}$$

Upshot:

The pragmatic speaker S_1 :

$$P_{S_1}(u \mid w) \propto \frac{P_{L_0}(w \mid u)^{\alpha}}{e^{\alpha \times C(u)}}$$

$$\propto \frac{\left(l(u, w) \times P(w) \times e^{G_{L_0}(u)}\right)^{\alpha}}{e^{\alpha \times C(u)}}$$

$$\propto l(u, w)^{\alpha} \times P(w)^{\alpha} \times e^{\alpha \times (G_{L_0}(u) - C(u))}$$

$$\propto l(u, w) \times e^{\alpha \times (G_{L_0}(u) - C(u))}$$

Upshot:

• l(u, w) is our filter

The pragmatic speaker S_1 :

$$P_{S_1}(u \mid w) \propto \frac{P_{L_0}(w \mid u)^{\alpha}}{e^{\alpha \times C(u)}}$$

$$\propto \frac{\left(l(u, w) \times P(w) \times e^{G_{L_0}(u)}\right)^{\alpha}}{e^{\alpha \times C(u)}}$$

$$\propto l(u, w)^{\alpha} \times P(w)^{\alpha} \times e^{\alpha \times (G_{L_0}(u) - C(u))}$$

$$\propto l(u, w) \times e^{\alpha \times (G_{L_0}(u) - C(u))}$$

Upshot:

- l(u, w) is our filter
- $P_{S_1}(u) \propto e^{\alpha \times (G_{L_0}(u) C(u))}$ is our *new* "prior"!

Reformulating the pragmatic speaker: example

$$[']$$
JP ran u km' $] = w \ge u$

Reformulating the pragmatic speaker: example


$$[']$$
JP ran u km' $] = w \ge u$


$$P_{S_1}(u) \propto e^{\alpha \times (G_{L_0}(u) - C(u))}$$

Reformulating the pragmatic speaker: example

$$[']$$
JP ran u km' $] = w \ge u$

$$P_{S_1}(u) \propto e^{\alpha \times (G_{L_0}(u) - C(u))}$$





Prior over JP's running distance

$$P_{S_1}(u|w=7)$$
 with $\alpha=4$

Making the normalizing constant explicit

As a proportion:

$$P_{S_1}(u|w) \propto l(u,w) \times P_{S_1}(u)$$

As a proportion:

$$P_{S_1}(u|w) \propto l(u,w) \times P_{S_1}(u)$$

Normalizing:

$$P_{S_1}(u|w) = \frac{l(u,w) \times P_{S_1}(u)}{\sum_{u' \in \mathcal{U}} l(u',w) \times P_{S_1}(u')}$$

As a proportion:

$$P_{S_1}(u|w) \propto l(u,w) \times P_{S_1}(u)$$

Normalizing:

$$P_{S_1}(u|w) = \frac{l(u, w) \times P_{S_1}(u)}{\sum_{u' \in \mathcal{U}} l(u', w) \times P_{S_1}(u')}$$
$$= l(u, w) \times P_{S_1}(u) \times e^{G_{S_1}(w)}$$

As a proportion:

$$P_{S_1}(u|w) \propto l(u,w) \times P_{S_1}(u)$$

Normalizing:

$$P_{S_1}(u|w) = \frac{l(u, w) \times P_{S_1}(u)}{\sum_{u' \in \mathcal{U}} l(u', w) \times P_{S_1}(u')}$$
$$= l(u, w) \times P_{S_1}(u) \times e^{G_{S_1}(w)}$$

As a proportion:

$$P_{S_1}(u|w) \propto l(u,w) \times P_{S_1}(u)$$

Normalizing:

$$P_{S_1}(u|w) = \frac{l(u, w) \times P_{S_1}(u)}{\sum_{u' \in \mathcal{U}} l(u', w) \times P_{S_1}(u')}$$
$$= l(u, w) \times P_{S_1}(u) \times e^{G_{S_1}(w)}$$

$$G_{S_1}(w) = -\log \sum_{u \in \mathcal{U}} l(u, w) \times P_{S_1}(u)$$

the specificity of w

$$P_{L_1}(w|u) \propto P_{S_1}(u|w) \times P(w)$$

$$P_{L_1}(w|u) \propto P_{S_1}(u|w) \times P(w)$$

$$\propto l(u, w) \times P_{S_1}(u) \times e^{G_{S_1}(w)} \times P(w)$$

$$P_{L_1}(w|u) \propto P_{S_1}(u|w) \times P(w)$$

$$\propto l(u, w) \times P_{S_1}(u) \times e^{G_{S_1}(w)} \times P(w)$$

$$\propto l(u, w) \times e^{G_{S_1}(w)} \times P(w)$$

$$P_{L_1}(w|u) \propto P_{S_1}(u|w) \times P(w)$$

$$\propto l(u, w) \times P_{S_1}(u) \times e^{G_{S_1}(w)} \times P(w)$$

$$\propto l(u, w) \times e^{G_{S_1}(w)} \times P(w)$$

The pragmatic listener L_1 :

$$P_{L_1}(w|u) \propto P_{S_1}(u|w) \times P(w)$$

$$\propto l(u, w) \times P_{S_1}(u) \times e^{G_{S_1}(w)} \times P(w)$$

$$\propto l(u, w) \times e^{G_{S_1}(w)} \times P(w)$$

Define $P_{L_1}(w) = e^{G_{S_1}(w)} \times P(w)$ as our *new* prior!

The pragmatic listener L_1 :

$$P_{L_1}(w|u) \propto P_{S_1}(u|w) \times P(w)$$

$$\propto l(u, w) \times P_{S_1}(u) \times e^{G_{S_1}(w)} \times P(w)$$


$$\propto l(u, w) \times e^{G_{S_1}(w)} \times P(w)$$

Define $P_{L_1}(w) = e^{G_{S_1}(w)} \times P(w)$ as our *new* prior!

Upshot:

$$P_{L_1}(w|u) \propto l(u,w) \times P_{L_1}(w)$$

Reformulating the pragmatic listener: example



Some consequences

Computing RSA models

Main result:


$$P_{L_1}(w|u) \propto l(u,w) \times P_{L_1}(w)$$

Computing RSA models

Main result:

$$P_{L_1}(w|u) \propto l(u,w) \times P_{L_1}(w)$$

The "pragmatic" interpretation of 'JP ran 2.4 km' is gotten by simply cropping $P_{L_1}(w)$ and normalizing:




Computing RSA models

Main result:

$$P_{L_1}(w|u) \propto l(u,w) \times P_{L_1}(w)$$

The "pragmatic" interpretation of 'JP ran 2.4 km' is gotten by simply cropping $P_{L_1}(w)$ and normalizing:



(Side note: the expected implicature is not always generated!)

From an algorithmic perspective (Marr, 1982), pragmatic interpretations (as according to RSA models) can be obtained the same way as literal interpretations: by filtering a certain prior by the literal meaning of the utterance.

From an algorithmic perspective (Marr, 1982), pragmatic interpretations (as according to RSA models) can be obtained the same way as literal interpretations: by filtering a certain prior by the literal meaning of the utterance.

• For literal interpretations, this prior is P(w).

From an algorithmic perspective (Marr, 1982), pragmatic interpretations (as according to RSA models) can be obtained the same way as literal interpretations: by filtering a certain prior by the literal meaning of the utterance.

- For literal interpretations, this prior is P(w).
- For pragmatic interpretations, this prior is $P_{L_1}(w)$.

From an algorithmic perspective (Marr, 1982), pragmatic interpretations (as according to RSA models) can be obtained the same way as literal interpretations: by filtering a certain prior by the literal meaning of the utterance.

- For literal interpretations, this prior is P(w).
- For pragmatic interpretations, this prior is $P_{L_1}(w)$.

From an algorithmic perspective (Marr, 1982), pragmatic interpretations (as according to RSA models) can be obtained the same way as literal interpretations: by filtering a certain prior by the literal meaning of the utterance.

- For literal interpretations, this prior is P(w).
- For pragmatic interpretations, this prior is $P_{L_1}(w)$.

(RSA-style) pragmatic interpretations can be *learned*, just as probability distributions over events can; pragmatic distributions are additionally sensitive to an event's specificity.

References

- Frank, Michael C., and Noah D. Goodman. 2012. Predicting pragmatic reasoning in language games. *Science* 336:998 998.
- Goodman, Noah D., and Michael C. Frank. 2016. Pragmatic Language Interpretation as Probabilistic Inference. *Trends in Cognitive Sciences* 20:818–829. https://www.sciencedirect.com/science/article/pii/S136466131630122X.
- Marr, David. 1982. Vision: A Computational Investigation into the Human Representation and Processing of Visual Information.

 Cambridge: MIT Press.