

Quick Reference: Context-sensitive Metrics (Adults, 70 kg reference)

Values are model-derived for comparative illustration and weight-normalized.

Drug	ke_0 (min^{-1})	CSHT-50 (10m)	CSHT-50 (1h)	CSHT-50 (4h)	CSHT-50 (8h)	Effect 80% (1h)	Effect 50% (1h)	Effect 25% (1h)
Propofol	0.26	45	83.7	90.2	91	29.8	94.4	190
Dexmedetomidine	0.28	64.2	179	201	204	54.6	198	411
Fentanyl	0.11	1.62	134	191	200	30.8	178	400
Hydromorphone	0.07	2.2	193	293	311	71.5	302	645
Remifentanyl	0.59	7.92	20.7	22.2	22.2	4.9	22.9	50.7
Alfentanil	0.46	2.9	238	318	333	16.1	254	611
Methadone	0.02	3.84	140	1.72e+03	2.09e+03	1.37e+03	3.09e+03	5.62e+03
Rocuronium	0.14	2.27	258	344	359	58.3	318	702
Vecuronium	0.1	2.18	281	418	443	60.2	384	864
Succinylcholine	1.39	27.1	61.3	67.1	67.6	13.6	62.9	136
Midazolam	0.07	7.19	171	207	213	97.4	251	477
Diazepam	0.02	4.72	233	5.7e+03	6.94e+03	2.91e+03	8.57e+03	1.69e+04

Models used: Propofol (Schnider), Remifentanyl (Minto), Fentanyl/Alfentanil (Shafer/Scott), Dexmedetomidine (Ebert/Talke), Hydromorphone (Rose/Lötsch), Methadone (Ferrari/Eap), NMBAs (Miller's + primary).

These values are illustrative unless exact coefficients are extracted from the cited models for the specified covariates.