

## Boring but important disclaimers:

- ▶ If you are not getting this from the GitHub repository or the associated Canvas page (e.g. CourseHero, Chegg etc.), you are probably getting the substandard version of these slides Don't pay money for those, because you can get the most updated version for free at

<https://github.com/julianmak/academic-notes>

The repository principally contains the compiled products rather than the source for size reasons.

- ▶ Associated Python code (as Jupyter notebooks mostly) will be held on the same repository. The source data however might be big, so I am going to be naughty and possibly just refer you to where you might get the data if that is the case (e.g. JRA-55 data). I know I should make properly reproducible binders etc., but I didn't...
- ▶ I do not claim the compiled products and/or code are completely mistake free (e.g. I know I don't write Pythonic code). Use the material however you like, but use it at your own risk.
- ▶ As said on the repository, I have tried to honestly use content that is self made, open source or explicitly open for fair use, and citations should be there. If however you are the copyright holder and you want the material taken down, please flag up the issue accordingly and I will happily try and swap out the relevant material.

# OCES 2003 : Descriptive Physical Oceanography

(a.k.a. physical oceanography by drawing pictures)

## Lecture 15: intro to waves




# Outline

- ▶ Recap: circulation and dependence on small-scale dynamics
- ▶ **waves**: fundamental concepts
  - periodicity, crest/trough/node
  - **wavelength** + **period**
  - **frequency** + **wavenumber**
  - **restoring force** + **dispersion relation**
  - propagation, **phase/group velocity**

**Key terms:** waves, wavenumber, frequency, period, dispersion relation, phase/group velocity

# Reacp: MOC



**Figure:** Schematic of the 3d MOC with watermass distributions. From Talley *et al.* (2011), *Descriptive Physical Oceanography*; see more in their Fig. 14.11. Format after Arnold Gordon (1991).

- **MOC** important for climate, carbon storage, ecology, etc.
  - e.g. warming of Western Europe by **AMOC**
  - e.g. carbon storage by deep water formation
- mostly **along-isopycnal** flow

- isolated places for **watermass transformation** + **deep/abyssal** water formation (**deep convection**)

## Recap: what goes down must come up




Figure: Schematic of the diffusive upwelling.

- diapycnal mixing contribute upwelling, strongest in boundary layers  
→ broad diffusive boundary intensified upwelling

what causes the boundary intensification of  $\kappa_d$ ?

# Recap: what goes down must come up




Figure: Schematic of the diffusive upwelling.

- diapycnal mixing contribute upwelling, strongest in boundary layers  
→ ~~broad~~ diffusive boundary intensified upwelling

what causes the boundary intensification of  $\kappa_d$ ? **dynamics!**

- at the surface, lots of things... (convection, waves, Langmuir turbulence etc.)
- at the bottom, probably tidal conversion (Lec. 18) → internal gravity waves (Lec. 17) → shear instabilities

# Recap: form stress and SO overturning

Role of **baroclinic instability** (Lec. 13), important for

- ▶ vertical **momentum** transfer by **interfacial form stress**
- ▶ scale transfer of **energy**  
→ **mesoscale eddies**, conduit between large-scales and **submesoscales**
- ▶ **along-isopycnal mixing** and also MOC




Figure: Schematic of form stress and eddy induced overturning cell in Southern Ocean (see Lec. 14)

# Outlook of the next few lectures

**Dynamics** important, next four lectures on

- ▶ **waves** (this Lec. + 16, 18) and **instabilities** (Lec. 17)


→ because waves are easier to talk about without maths...

Highlight gross features (i.e. those that can be drawn...)

- ▶ **how to describe waves** (Lec. 15)
- ▶ types of waves (Lec. 16)
  - consequence + leading to instabilities
- ▶ instabilities
- ▶ tides (particularly as **internal gravity waves**) (Lec. 17)



# Examples of waves



**Figure:** Gravity waves with signal at the sea surface (as darker and lighter bands). Taken at HKUST.


## Surface gravity waves<sup>1</sup>

► restoring force is buoyancy

→ treat air-sea as one fluid with a giant jump in density

<sup>1</sup> According to Richard Feynman, while water waves are “...easily seen by everyone... are the worse possible example [of waves], because... they have all the complications that waves can have (From Feynman, Leighton & Sands, 1971, *Lectures of Physics*).

## Examples of waves




**Figure:** Flow over topography (e.g. tidal motion) leading to wave generation.

Tides and/or internal gravity waves forced by tidal motion

- ▶ restoring force is still buoyancy
- wave breaking contributing to mixing

## Examples of waves






**Figure:** Example of jet stream meanders as Rossby waves. Figure from Wikipedia.

**Rossby waves** (cf. same Rossby as Rossby number)

- **restoring force** is Coriolis (or background gradient in vorticity)
  - planetary-scale waves, generic in rotating systems (e.g. rotating tanks, Jupiter, galactic disks)

# Examples of waves



**Figure:** Example of some other waves in systems that support waves: Alfvén waves (magnetic field + Lorentz force), gravitational waves (spacetime + gravity), electromagnetic waves (but also **wave-particle duality**), sound waves (mechanical forcing + any medium). All figures from Wikipedia except the cello one.

# Features of waves

Some observations:

- ▶ waves have some **oscillation/periodicity**  
→ want a measure of **period**
- ▶ waves to **propagate**  
→ **speed/velocity** associated with waves
- ▶ waves need a **medium** to travel through  
→ subtlety with Electro-Magnetic waves (not touched on here)
- ▶ characterised by a **restoring force** (follows from medium)
- ▶ waves can increase in **amplitude** and **steepen**  
→ wave breaking and mixing
- ▶ can **disperse, refract, interfere** etc. (used in Lec. 17, 20)


# Features of waves

Some observations:

- ▶ waves have some **oscillation/periodicity**  
→ want a measure of **period**
- ▶ waves to **propagate**  
→ **speed/velocity** associated with waves
- ▶ waves need a **medium** to travel through  
→ subtlety with Electro-Magnetic waves (not touched on here)
- ▶ characterised by a **restoring force** (follows from medium)
- ▶ waves can increase in **amplitude** and **steepen**  
→ wave breaking and mixing
- ▶ can **disperse, refract, interfere** etc. (used in Lec. 17, 20)

physics  $\Rightarrow$  **dispersion relation**, identifies the type of waves

# Features of waves



**Figure:** Schematic of wave features. Box length  $L = 2\pi$  for simplicity.

- displacement  $\eta$  described by (could also be sine)

$$\eta \sim \cos(x)$$

# Features of waves





Figure: Schematic of wave features. Box length  $L = 2\pi$  for simplicity.

- displacement  $\eta$  described by (could also be sine)

$$\eta \sim \cos(x)$$



# Features of waves



**Figure:** Schematic of wave features. Box length  $L = 2\pi$  for simplicity.

- **displacement**  $\eta$  described by (could also be sine)

$$\eta \sim \cos(x - vt)$$

# Features of waves




Figure: Schematic of wave features. Box length  $L = 2\pi$  for simplicity.

- displacement  $\eta$  described by (could also be sine)

$$\eta \sim A \cos(x - vt), \quad \gamma = v/\lambda$$

# Features of waves




Figure: Schematic of wave features. Box length  $L = 2\pi$  for simplicity.

- displacement  $\eta$  described by (could also be sine)

$$\eta \sim A \cos(x), \quad \eta \sim A \cos(x - \lambda/4)$$

# Features of waves





Figure: Schematic of wave features. Box length  $L = 2\pi$  for simplicity.

- displacement  $\eta$  described by (could also be sine)

$$\eta \sim A \cos(x), \quad \eta \sim A \cos(x - \lambda/2) \sim -A \sin(x)$$

# Features of waves



**Figure:** Schematic of wave features. Box length  $L = 2\pi$  for simplicity.

- displacement  $\eta$  described by (could also be sine)

$$\eta \sim A \cos(x)$$

# Features of waves




Figure: Schematic of wave features. Box length  $L = 2\pi$  for simplicity.

- displacement  $\eta$  described by (could also be sine)

$$\eta \sim A \cos(2x), \quad \eta \sim A \cos(1x), \quad \eta \sim A \cos(3x)$$

# Features of waves

$$\gamma = \frac{v}{\lambda}, \quad k = \frac{2\pi}{\lambda}$$

- ▶  $\gamma$  the **frequency** (units:  $\text{s}^{-1} = \text{Hz}$ )  
→ how quickly the wave oscillates
- ▶  $v = c_p$  the **phase velocity**  
→ how fast the wave itself moves around
- ▶  $\lambda$  the **wavelength**  
→ how long the wave is
- ▶  $k$  the **wavenumber**  
→ intuitively how many waves can you fit in a box (so  $k \sim \lambda^{-1}$ )  
→ does not necessarily have to be an integer

# Features of waves: dispersion relation

Usually describe waves in terms of **wavenumber**  $k$  and the **angular frequency**  $\omega = 2\pi\gamma$ , i.e.

$$\eta = A \cos(kx - \omega t)$$

- generally, for  $\mathbf{x} = (x, y, z)$ , we would have

$$\eta = A \cos(\mathbf{k} \cdot \mathbf{x} - \omega t)$$

→  $k$  is the **wavevector**



# Features of waves: dispersion relation

Usually describe waves in terms of **wavenumber**  $k$  and the **angular frequency**  $\omega = 2\pi\gamma$ , i.e.

$$\eta = A \cos(kx - \omega t)$$

► generally, for  $\mathbf{x} = (x, y, z)$ , we would have

$$\eta = A \cos(\mathbf{k} \cdot \mathbf{x} - \omega t)$$

→  $k$  is the **wavevector**

Aside: If you know your **complex numbers**, the above is neatly encapsulated as

$$\eta = \text{Real} \left[ A e^{i(\mathbf{k} \cdot \mathbf{x} - \omega t - \theta_0)} \right],$$

where  $e$  is Euler's number,  $i = \sqrt{-1}$ , and  $\theta_0$  denotes a phase shift if any (could be sucked into the amplitude). For calculating the **dispersion relation** this form is substantially nicer to deal with (don't have to keep track of sines and cosines when taking derivatives).

# Features of waves: dispersion relation

Note that

$$\omega = 2\pi\gamma = 2\pi\frac{v}{\lambda} = vk$$

- ▶ the physics tells you how  $v = v(k)$
- ▶ the **dispersion relation** is given by

$$\omega = \mathcal{F}(k; \dots)$$


for some function  $\mathcal{F}$

→ the dispersion identifies the types of wave (see Lec. 16), e.g.

$$\omega = \sqrt{gk}, \quad \omega = -\frac{\beta}{k}, \quad \omega = B_0 k, \quad \omega = \frac{\hbar k^2}{2m},$$

# Superposition


(Linear) waves can be superimposed, leading to **interference**



**Figure:** Interference of waves with **Red** = **Orange** + **Green**. For waves in phase (**constructive** interference) and waves in anti-phase (**destructive** interference).

**Q.** but what about waves not quite in phase or anti-phase?

# Superposition



**Figure:** Superposition of two waves slightly out of phase, again with Red = Orange + Green. The crests have a marker marked on to track its progress later.

# Wave propagation

Recall that  $v = v(k)$  is the wave velocity. More precisely,

- ▶ the **phase speed** in a direction ( $= v$ ) is defined as

$$c_{p,x} = \frac{\omega}{k}$$

→ how the wave by itself travels

# Wave propagation

Recall that  $v = v(k)$  is the wave velocity. More precisely,

- ▶ the **phase speed** in a direction ( $= v$ ) is defined as

$$c_{p,x} = \frac{\omega}{k}$$

→ how the wave by itself travels

- ▶ the **group velocity**  $c_g$  is defined as

$$c_{g,x} = \frac{\partial \omega}{\partial k}$$

- ▶ in higher space dimensions,

$$c_{p,x} = \frac{\omega}{k_x}, \quad c_{p,y} = \frac{\omega}{k_y}, \quad c_{p,z} = \frac{\omega}{k_z}, \quad c_g = \nabla_k \omega$$

→ NOTE! Phase propagates in the direction of  $k$

(subtlety:  $c_{p,x}$  is not a component of the velocity that phases propagate at; see Ch.5 App of Vallis (2006))

## Example: 1d Rossby waves (animation)

# Wave propagation

- ▶ group velocity describes
  - how a collection of waves travel as a **group** or **wavepacket**
  - velocity that “stuff” propagates at
- ▶ a type of wave is **non-dispersive** if


$$c_g = c_p$$

- e.g.  $\omega = B_0 k$  and  $\omega = k\sqrt{gH}$  are non-dispersive
- if non-dispersive, wavepacket and phase travel together

- ▶ example just now is **dispersive**
  - 1d Rossby waves,  $\omega = -\beta/k$  (exercise: show  $c_g = -c_p$  for this case)



# Wave propagation




**Figure:** Schematic of **refraction**, **reflection** (and **transmission**), and **diffraction**, nominally using **monochromatic** (i.e. one choice of  $k$ ) surface gravity wave as an example. The orange lines are phase lines (e.g. think wave crests).

**refraction**, **reflection**, and **diffraction** (used in Lec. 17, 20)

- ▶ resulting interference of waves can lead to **wave steepening** and wave breaking

# Wave propagation



**Figure:** Picture of (presumably non-monochromatic) waves over the Arabian sea. Image taken from <https://www.earthglance.com/post/133835790223/wave-diffraction-on-the-arabian-sea>.

# Summary

## smaller-scale dynamics affects large-scale circulation

- ▶ **waves** are ubiquitous physical features
  - depends on physics
- ▶ wave described by the **dispersion relation**  $\omega = \mathcal{F}(k)$ 
  - physics of the system dictates what  $\mathcal{F}(k)$  is
  - usually use **angular frequency**  $\omega$  and **wavenumber**  $k$   
(absorbs factors of  $2\pi$  floating around)
  - $k \sim \lambda^{-1}$  sometimes used to characterise **scale of motion**  
(more on this in Lec. 17)
- ▶ difference in  $c_p$  and  $c_g$ 
  - individual (former) and collective (latter) velocity

**wave breaking contributes to diapycnal mixing** (see Lec. 17)