

University of Manitoba

Math 38200 – Winter 2008

Assignment 1

Due Friday, February 8, 2008

Instructions

This assignment is due in the Mathematics Office (342 Machray Hall) Friday, February 8 before 16:30, or by email (same deadline). I accept email submissions, so if you want to typeset or scan your answers, this is fine (pdf is preferred). Be clear in your answers. If you use a result in the course notes or from another source, please indicate clearly the reference. You can work on this problem with others, but the solution you hand in *must* be yours.

The Ricker model of growth of a single population takes the form

$$N_{t+1} = N_t e^{r(1 - \frac{N_t}{K})}, \quad (1)$$

with $r, K > 0$ and initial condition $N_0 > 0$. The aim of this assignment is to study the behavior of (1).

1. Define $x_t = N_t/K$, and show that the difference equation (1) then takes the form, as a function of the dimensionless variable x_t ,

$$x_{t+1} = x_t e^{r(1-x_t)}, \quad (2)$$

2. Show that if $x_0 > 0$ (where $x_0 = N_0/K$), there holds that $x_t > 0$ for all t in (2).

3. Determine the fixed points of (2), as well as their stability as a function of the parameter r .

4. Show that (2) has no points of period 2 for $0 < r < 2$.

5. Try to find 2-periodic points of (2) analytically (show your work). Then, do so using a numerical software. Under what conditions are these periodic points stable? Evaluate the stability of the points you found for a few sample values of r , using a numerical software.

6. Using numerical software, draw a bifurcation diagram for (2), for r varying in $(0, 5]$. What do you observe?