

# Analysis of simple tumour-growth models

Bob W. Kooi

Dept. Theoretical Biology,  
Vrije Universiteit, de Boelelaan 1087,  
1081 HV Amsterdam,  
The Netherlands

`kooi@bio.vu.nl`

`http://www.bio.vu.nl/thb/`



# Outline

- Introduction
- Simple models
  - exponential growth
  - logistic growth
  - Gompertz growth
  - Minimal mitotic oscillator cascade model
- Attractors
- Bifurcation theory
- Computer exercises (octave)

## Logistic growth population model

Density-dependent growth

$$\frac{dN}{dt} = f(N) = rN\left(1 - \frac{N}{K}\right)$$

$N(t) \geq 0$ : tumour size

$r > 0$ : intrinsic growth rate

$K > 0$ : carrying capacity

Solution of initial value problem:

$$N(t) = \frac{N(0)e^{rt}}{1 + (e^{rt} - 1)N(0)/K}$$

$K = \infty$ : then exponential growth

## Equilibria

$$N = 0 \quad \text{and} \quad N = K$$

Stability analysis of  $N = 0$

consider linearized equation where  $x = N$

$$\frac{dx}{dt} = rx \text{ solution } x(t) = x(0)e^{rt}$$

since  $r > 0$  perturbations lead to divergence

Conclusion: unstable

Stability analysis of  $N = K$

consider linearized equation where  $x = N - K$


$$\frac{dx}{dt} = \left( \frac{df}{dN} \right)_K x = -rx \text{ solution } x(t) = x(0)e^{-rt}$$

since  $r > 0$  perturbations lead to convergence

Conclusion: stable

## Logistic growth and Gompertz:

$$r = 1 \text{ en } K = 1$$



## Gompertz growth population model

Non-autonomous growth equation

$$\frac{dN}{dt} = f(t, N) = r_0 e^{-\alpha t} N$$

$N(t) \geq 0$ : tumour size

$r_0 > 0$ : initial intrinsic growth rate

$\alpha > 0$ : exponential decay rate of growth

Solution of initial value problem:

$$N(t) = K \left( \frac{N_0}{K} \right) e^{-\alpha t}$$

where

$$K = N_0 e^{r_0/\alpha}, \quad \text{that is} \quad \alpha = \frac{r_0}{\ln(K/N_0)}$$

Another formulation

$$\frac{dN}{dt} = -\alpha \ln(N/K) N$$

## Logistic growth with predation

$$\frac{dN}{dt} = rN\left(1 - \frac{N}{K}\right) - p$$

$N(t)$ : tumour size

$r > 0$ : intrinsic growth rate

$K > 0$ : carrying capacity

$p$ : constant loss rate (due to treatment?)

## Equilibria

$$N^* = K \left( \frac{1}{2} \pm \sqrt{\frac{1}{4} - \frac{p}{rK}} \right)$$

$0 \leq p < rK/4$  : two positive equilibria

$p = p_T = rK/4$  : one equilibrium  $N^* = 1/2K$

$p > rK/4$  : no solution!



## Stability

$$0 \leq p \leq rK/4$$

$$\frac{dx}{dt} = \mp r \left( \sqrt{1 - \frac{4p}{rK}} \right) x$$

$p < p_T$  then:

Stable:  $N^* = K(1/2 + \sqrt{(1/4 - p/rK)})$

Unstable:  $N^* = K(1/2 - \sqrt{(1/4 - p/rK)})$

$R(0) > K(1/2 - \sqrt{(1/4 - p/rK)})$ :


convergence to stable equilibrium

$R(0) < K(1/2 - \sqrt{(1/4 - p/rK)})$ :

solution becomes negative

**Logistic growth with predation where:**

$$r = 1 \text{ en } K = 1$$




Bifurcation at  $p = p_T$

solid line: stable equilibrium

dashed line: unstable equilibrium

# Logistic growth with predation

where  $r = 1$  en  $K = 1$



## Stability of Gompertz model

We formulate the model as an equivalent autonomous system:

$$\begin{aligned}\frac{dN}{dt} &= f(N, r) = rN \\ \frac{dr}{dt} &= g(N, r) = -\alpha r\end{aligned}$$

$N(t)$ : tumour size

$r(t)$ : actual growth rate

and initial conditions:

$$N(0) = N_0 \text{ and } r(0) = r_0$$

## Equilibria

Trivial solution

$$N = 0$$

$$r = 0$$

and positive (tumour size) solution

$$N = K$$

$$r = 0$$

Stability analysis is much more difficult!

Now two state-variables

## Linearization: Jacobian matrix $\mathbf{J}$

$$\begin{pmatrix} \frac{dx}{dt} \\ \frac{dy}{dt} \end{pmatrix} = \mathbf{J} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} \mathbf{J}_{11} & \mathbf{J}_{12} \\ \mathbf{J}_{21} & \mathbf{J}_{22} \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}$$

$$\begin{pmatrix} \frac{dx}{dt} \\ \frac{dy}{dt} \end{pmatrix} = \begin{pmatrix} \frac{\partial f}{\partial N} & \frac{\partial f}{\partial r} \\ \frac{\partial g}{\partial N} & \frac{\partial g}{\partial r} \end{pmatrix}_{N^*, r^*} \begin{pmatrix} x \\ y \end{pmatrix}$$

where  $x(t) = N(t) - N^*$  and  $y(t) = r(t) - r^*$

## **characteristic equation**

When characteristic equation is satisfied

$$\lambda^2 + (\mathbf{J}_{11} + \mathbf{J}_{22})\lambda + \mathbf{J}_{11}\mathbf{J}_{22} - \mathbf{J}_{12}\mathbf{J}_{21} = 0$$

There are two solutions:

$$\lambda_{1,2} = 0.5(\mathbf{J}_{11} + \mathbf{J}_{22}) \pm \sqrt{(0.5(\mathbf{J}_{11} + \mathbf{J}_{22}))^2 - (\mathbf{J}_{11}\mathbf{J}_{22} - \mathbf{J}_{12}\mathbf{J}_{21})}$$

$\lambda_{1,2}$  are called *eigenvalues*




General solution

$$\begin{pmatrix} x(t) \\ y(t) \end{pmatrix} = c_1 e^{\lambda_1 t} \mathbf{v}_1 + c_2 e^{\lambda_2 t} \mathbf{v}_2$$

$\lambda_1$  and  $\lambda_2$  are eigenvalues;  $\mathbf{v}_1$  and  $\mathbf{v}_2$  eigenvectors of the Jacobian  $\mathbf{J}$  calculated in equilibrium.

Constants  $c_1$  and  $c_2$  are fixed by the initial conditions  $x(0)$  en  $y(0)$

## Classification of equilibrium points



$$\text{tr } \mathbf{J} = \mathbf{J}_{11} + \mathbf{J}_{22} = \text{Re } \lambda_1 + \text{Re } \lambda_2$$

$$\det \mathbf{J} = \mathbf{J}_{11}\mathbf{J}_{22} - \mathbf{J}_{12}\mathbf{J}_{21} = \lambda_1\lambda_2$$

$$\text{disc } \mathbf{J} = 0.25 (\text{tr } \mathbf{J})^2 - \det \mathbf{J}$$

$$\text{where } \lambda_{1,2} = 0.5(\text{tr } \mathbf{J} \pm \sqrt{\text{disc } \mathbf{J}})$$

Equilibrium is stable when:  $\text{Re } \lambda_1 < 0$  and  $\text{Re } \lambda_2 < 0$

## Stability of Gompertz model

non-zero solution  $N^* = K$

$$\begin{pmatrix} \frac{dx}{dt} \\ \frac{dy}{dt} \end{pmatrix} = \begin{pmatrix} r & N^* \\ 0 & -\alpha \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 & K \\ 0 & -\alpha \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}$$
$$r^* = 0$$

$$\text{tr } \mathbf{J} = -\alpha \text{ and } \det \mathbf{J} = 0$$

eigenvalues  $\lambda_1 = 0$  and  $\lambda_2 = -\alpha$

eigenvectors

$$\mathbf{v}_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \quad \text{and} \quad \mathbf{v}_2 = \begin{pmatrix} -K/\alpha \\ 1 \end{pmatrix}$$

- $\alpha$  primary parameter  
 $K$  depends on initial value  $N_0$  and therefore  
no stability
- $K$  primary parameter  
 $\alpha$  depends on  $N_0$   
Because one zero eigenvalue no stability  
If  $r_0 > 0$  then:  $N_0 < K$  then  $\alpha > 0 \Rightarrow$   
 $\lambda_2 < 0$  and locally convergence to  $K$  while  
 $N_0 > K$  implies  $\alpha < 0$  and convergence to  
 $K$  is impossible


It can be shown that globally with  $r_0 > 0$

- $0 < N(0) \leq K$ : solution converges to  $K$   
for  $t \rightarrow \infty$
- $N(0) > K$ : solution grows unbounded  
for  $t \rightarrow \infty$
- $N(0) = K$ :  $\alpha = 0$  and therefore  $N(t) =$   
 $N_0 = K$

## **Bifurcation theory**

- Find equilibrium
- Stability analysis
- Continuation by varying one free parameter
- Find bifurcation point
- Continuation by varying another free parameter

# minimal cascade model of Goldbeter (1997)



## mitotic oscillator

The Goldbeter model for mitotic oscillator:

$$\begin{aligned}\frac{dC}{dt} &= v_i - v_d X \frac{C}{K_d + C} - k_d C , \\ \frac{dM}{dt} &= V_1 \frac{(1 - M)}{K_1 + (1 - M)} - V_2 \frac{M}{K_2 + M} \\ \frac{dX}{dt} &= V_3 \frac{(1 - X)}{K_3 + (1 - X)} - V_4 \frac{X}{K_4 + X}\end{aligned}$$

where

$$V_1 = V_{M1} \frac{C}{K_c + C} \quad , \quad V_3 = V_{M3} M$$

$C(t)$ : concentration cyclin

$M(t)$ : fraction active cdc2-kinase

$X(t)$ : fraction active cyclin protease

## parameter values

The name cyclin reflects its periodic variation during cell cycle

Only interphase and mitosis in the division cycle are distinguished

Model parameter are realistic for rapidly dividing amphibian embryonic cells

Period of the mitotic oscillator is of the order of 30 min


Parameters:  $K_1 = K_2 = K_3 = K_4 = 0.005$ ,  $v_i = 0.025 \mu\text{M}/\text{min}$ ,  $v_d = 0.25 \mu\text{M}/\text{min}$ ,  $K_d = 0.02 \mu\text{M}$ ,  $k_d = 0.01 \text{ min}^{-1}$ ,  $V_{M1} = 3$ ,  $V_2 = 1.5$ ,  $V_{M3} = 1$  and  $V_4 = 0.5$ ,  $K_c = 0.5 \mu\text{M}$

Initial conditions:


$C(0) = 0.01 \mu\text{M}$ ,  $M(0) = X(0) = 0.01$



## oscillations




## oscillations



# One-parameter diagram


$$K_3 = K_4 = 0.002$$



$H$  Hopf bifurcation point

$T_c$  Tangent bifurcation point for cycle

## One-parameter diagram(details)



# Phase-plane plot

$$K_1 = K_2 = 600, K_3 = K_4 = 0.002$$

