



University
of Manitoba

Math prelims – Linear algebra

MATH 2740 – Mathematics of Data Science – Unit U02

Julien Arino

`julien.arino@umanitoba.ca`

Department of Mathematics @ University of Manitoba

Fall 2026

The University of Manitoba campuses are located on original lands of Anishinaabeg, Ininew, Anisininew, Dakota and Dene peoples, and on the National Homeland of the Red River Métis. We respect the Treaties that were made on these territories, we acknowledge the harms and mistakes of the past, and we dedicate ourselves to move forward in partnership with Indigenous communities in a spirit of Reconciliation and collaboration.

In MATH 2740, we rely on notions you acquired in MATH 1210/1220/1300. We also use some material from first-year calculus

So let us (briefly) go over material in these courses

I also add (for some of you) a few things that will be handy and establish some terminology that we use throughout the course

Outline


Sets and logic

Complex numbers

Vectors and vector spaces

Linear systems and matrices

A crash course in multivariable calculus

A robot with a boxy, metallic body stands in the foreground of a vast, arid desert landscape. The robot is facing away from the viewer, looking towards a large, leafless tree that dominates the mid-ground. The ground is sandy and dotted with small rocks. In the background, there are rolling hills under a hazy, orange-tinted sky, suggesting a sunset or sunrise. The overall mood is desolate and contemplative.

Sets and logic

Complex numbers

Vectors and vector spaces

Linear systems and matrices

A crash course in multivariable calculus

Sets and elements

Definition 1 (Set)

A **set** X is a collection of **elements**

We write $x \in X$ or $x \notin X$ to indicate that the element x belongs to the set X or does not belong to the set X , respectively

Definition 2 (Subset)

Let X be a set. The set S is a **subset** of X , which is denoted $S \subset X$, if all its elements belong to X

Not used here but worth noting: we say S is a **proper subset** of X and write $S \subsetneq X$, if it is a subset of X and not equal to X

Quantifiers

A shorthand notation for “for all elements x belonging to X ” is $\forall x \in X$

For example, if $X = \mathbb{R}$, the *field* of real numbers, then $\forall x \in \mathbb{R}$ means “for all real numbers x ”

A shorthand notation for “there exists an element x in the set X ” is $\exists x \in X$

$\forall$ and $\exists$ are **quantifiers**

Intersection and union of sets

Let X and Y be two sets

Definition 3 (Intersection)

The intersection of X and Y , $X \cap Y$, is the set of elements that belong to X **and** to Y ,

$$X \cap Y = \{x : x \in X \text{ and } x \in Y\}$$

Definition 4 (Union)

The union of X and Y , $X \cup Y$, is the set of elements that belong to X **or** to Y ,

$$X \cup Y = \{x : x \in X \text{ or } x \in Y\}$$

In mathematics, or=and/or in common parlance. We also have an **exclusive or** (xor)

A teeny bit of logic

In a logical sense, a **proposition** is an assertion (or statement) whose truth value (true or false) can be asserted. For example, a theorem is a proposition that has been shown to be true. “The sky is blue” is also a proposition

Let A be a proposition. We generally write

A

to mean that A is true, and

not A

to mean that A is false. **not** A is the **negation** of A (or **not** A is the negative of A)

A teeny bit of logic (cont.)

Let A, B be propositions. Then

- ▶ $A \Rightarrow B$ (read A implies B) means that whenever A is true, then so is B
- ▶ $A \Leftrightarrow B$, also denoted A if and only if B (A iff B for short), means that $A \Rightarrow B$ **and** $B \Rightarrow A$

We also say that A and B are **equivalent**

Let A and B be propositions. Then

$$(A \Rightarrow B) \Leftrightarrow (\text{not } B \Rightarrow \text{not } A)$$

Necessary or sufficient conditions

Suppose we want to establish whether a given statement P is true, depending on the truth value of a statement H . Then we say that

- ▶ H is a **necessary condition** if $P \Rightarrow H$
(It is necessary that H be true for P to be true; so whenever P is true, so is H)
- ▶ H is a **sufficient condition** if $H \Rightarrow P$
(It suffices for H to be true for P to also be true)
- ▶ H is a **necessary and sufficient condition** if $H \Leftrightarrow P$, i.e., H and P are equivalent

Playing with quantifiers

For the quantifiers $\forall$ (for all) and $\exists$ (there exists),

$\exists$ is the negation of $\forall$

Therefore, for example, the negation of

$$\forall x \in X, \exists y \in Y$$

is

$$\exists x \in X, \forall y \in Y$$


Sets and logic

Complex numbers

Vectors and vector spaces

Linear systems and matrices

A crash course in multivariable calculus



Complex numbers

Definition 5 (Complex numbers)

A **complex number** is an ordered pair (a, b) , where $a, b \in \mathbb{R}$. Usually written $a + ib$ or $a + bi$, where $i^2 = -1$ (i.e., $i = \sqrt{-1}$)

The set of all complex numbers is denoted $\mathbb{C}$,

$$\mathbb{C} = \{a + ib : a, b \in \mathbb{R}\}$$

Definition 6 (Addition and multiplication on $\mathbb{C}$)

Letting $a + ib$ and $c + id \in \mathbb{C}$, addition on $\mathbb{C}$ is defined by

$$(a + ib) + (c + id) = (a + c) + i(b + d)$$

and multiplication on $\mathbb{C}$ is defined by

$$(a + ib)(c + id) = (ac - bd) + i(ad + bc)$$

Latter is easy to obtain using regular multiplication and $i^2 = -1$

Properties

$$\forall \alpha, \beta, \gamma \in \mathbb{C},$$

$$\alpha + \beta = \beta + \alpha \text{ and } \alpha\beta = \beta\alpha$$

[commutativity]

$$(\alpha + \beta) + \gamma = \alpha + (\beta + \gamma) \text{ and } (\alpha\beta)\gamma = \alpha(\beta\gamma)$$

[associativity]

$$\gamma + 0 = \gamma \text{ and } \gamma 1 = \gamma$$

[identities]

$$\forall \alpha \in \mathbb{C}, \exists \beta \in \mathbb{C} \text{ unique s.t. } \alpha + \beta = 0$$

[additive inverse]

$$\forall \alpha \neq 0 \in \mathbb{C}, \exists \beta \in \mathbb{C} \text{ unique s.t. } \alpha\beta = 1$$

[multiplicative inverse]

$$\gamma(\alpha + \beta) = \gamma\alpha + \gamma\beta$$

[distributivity]

Additive & multiplicative inverse, subtraction, division

Definition 7

Let $\alpha, \beta \in \mathbb{C}$

- ▶ $-\alpha$ is the **additive inverse** of α , i.e., the unique number in $\mathbb{C}$ s.t. $\alpha + (-\alpha) = 0$

- ▶ **Subtraction** on $\mathbb{C}$:

$$\beta - \alpha = \beta + (-\alpha)$$

- ▶ For $\alpha \neq 0$, $1/\alpha$ is the **multiplicative inverse** of α , i.e., the unique number in $\mathbb{C}$ s.t.

$$\alpha(1/\alpha) = 1$$

- ▶ **Division** on $\mathbb{C}$:

$$\beta/\alpha = \beta(1/\alpha)$$

Definition 8 (Real and imaginary parts)

Let $z = a + ib$. Then $\operatorname{Re} z = a$ is **real part** and $\operatorname{Im} z = b$ is **imaginary part** of z

If ambiguous, write $\operatorname{Re}(z)$ and $\operatorname{Im}(z)$

Definition 9 (Conjugate and Modulus)

Let $z = a + ib \in \mathbb{C}$. Then

- ▶ **Complex conjugate** of z is

$$\bar{z} = a - ib$$

- ▶ **Modulus** (or **absolute value**) of z is

$$|z| = \sqrt{a^2 + b^2} \geq 0$$

Properties of complex numbers

Let $w, z \in \mathbb{C}$, then

▶ $z + \bar{z} = 2\operatorname{Re} z$

▶ $z - \bar{z} = 2i\operatorname{Im} z$

▶ $z\bar{z} = |z|^2$

▶ $\overline{\bar{w} + \bar{z}} = \bar{w} + z$ and $\overline{wz} = \bar{w}\bar{z}$

▶ $\overline{\bar{z}} = z$

▶ $|\operatorname{Re} z| \leq |z|$ and $|\operatorname{Im} z| \leq |z|$

▶ $|\bar{z}| = |z|$

▶ $|wz| = |w| |z|$

▶ $|w + z| \leq |w| + |z|$

[triangle inequality]

Solving quadratic equations

Consider the polynomial

$$P(x) = a_0 + a_1x + a_2x^2$$

where $x, a_0, a_1, a_2 \in \mathbb{R}$. Letting

$$\Delta = a_1^2 - 4a_0a_2$$

you know that if $\Delta > 0$, then

$$P(x) = 0$$

has two distinct *real* solutions,

$$x_1 = \frac{-a_1 - \sqrt{\Delta}}{2a_2} \quad \text{and} \quad x_2 = \frac{-a_1 + \sqrt{\Delta}}{2a_2}$$

if $\Delta = 0$, then there is a (multiplicity 2) unique *real* solution

$$x_1 = \frac{-a_1}{2a_2}$$

while if $\Delta < 0$, there is no solution

Solving quadratic equations with complex numbers

Consider the polynomial

$$P(x) = a_0 + a_1x + a_2x^2$$

where $x, a_0, a_1, a_2 \in \mathbb{R}$. If instead of seeking $x \in \mathbb{R}$, we seek $x \in \mathbb{C}$, then the situation is the same, except when $\Delta < 0$

In the latter case, note that

$$\sqrt{\Delta} = \sqrt{(-1)(-\Delta)} = \sqrt{-1}\sqrt{-\Delta} = i\sqrt{-\Delta}$$

Since $\Delta < 0$, $-\Delta > 0$ and the square root is the usual one

Solving quadratic equations with complex numbers

To summarize, consider the polynomial

$$P(x) = a_0 + a_1x + a_2x^2$$

where $x, a_0, a_1, a_2 \in \mathbb{R}$. Letting

$$\Delta = a_1^2 - 4a_0a_2$$

Then

$$P(x) = 0$$

has two solutions,

$$x_{1,2} = \frac{-a_1 \pm \sqrt{\Delta}}{2a_2}$$

where, if $\Delta < 0$, $x_1, x_2 \in \mathbb{C}$ and take the form

$$x_{1,2} = \frac{-a_1 \pm i\sqrt{-\Delta}}{2a_2}$$

Why this matters

Recall (we will come back to this later) that to find the *eigenvalues* of the matrix

$$A = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix}$$

we seek λ solutions to $\det(A - \lambda\mathbb{I}) = 0$, i.e., λ solutions to

$$|A - \lambda\mathbb{I}| = \begin{vmatrix} a_{11} - \lambda & a_{12} \\ a_{21} & a_{22} - \lambda \end{vmatrix} = (a_{11} - \lambda)(a_{22} - \lambda) - a_{12}a_{21} = 0$$

i.e., λ solutions to

$$\lambda^2 - (a_{11} + a_{22})\lambda + a_{11}a_{22} - a_{12}a_{21} = 0$$

Why this matters (cont.)

Let

$$P(\lambda) = \lambda^2 - (a_{11} + a_{22})\lambda + a_{11}a_{22} - a_{12}a_{21}$$

From previous discussion, letting


$$\begin{aligned}\Delta &= (a_{11} + a_{22})^2 - 4(a_{11}a_{22} - a_{12}a_{21}) \\ &= a_{11}^2 + a_{22}^2 + 2a_{11}a_{22} - 4a_{11}a_{22} + 4a_{12}a_{21} \\ &= a_{11}^2 + a_{22}^2 - 2a_{11}a_{22} + 4a_{12}a_{21} \\ &= (a_{11} - a_{22})^2 + 4a_{12}a_{21}\end{aligned}$$

we have two (potentially equal) solutions to $P(\lambda) = 0$

$$x_{1,2} = \frac{a_{11} + a_{22} \pm \sqrt{\Delta}}{2}$$

that are complex if $\Delta < 0$

Example: $\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$

A robot with a boxy, metallic body stands in the foreground of a vast, arid desert landscape. The robot is facing away from the viewer, looking towards a large, leafless tree that dominates the right side of the background. The ground is sandy and dotted with small rocks. The sky is a pale, hazy blue, and the overall atmosphere is one of isolation and contemplation.

Sets and logic

Complex numbers

Vectors and vector spaces

Linear systems and matrices

A crash course in multivariable calculus

Vectors

A **vector** $\mathbf{v}$ is an ordered n -tuple of real or complex numbers

Denote $\mathbb{F} = \mathbb{R}$ or $\mathbb{C}$ (real or complex numbers). For $v_1, \dots, v_n \in \mathbb{F}$,

$$\mathbf{v} = (v_1, \dots, v_n) \in \mathbb{F}^n$$

is a vector. $v_1, \dots, v_n$ are the **components** of $\mathbf{v}$

If unambiguous, we write v . Otherwise, $\mathbf{v}$ or $\vec{v}$

Vector space

Definition 10 (Vector space)

A **vector space** over $\mathbb{F}$ is a set V together with two binary operations, **vector addition**, denoted $+$, and **scalar multiplication**, that satisfy the relations:

1. $\forall \mathbf{u}, \mathbf{v}, \mathbf{w} \in V, \mathbf{u} + (\mathbf{v} + \mathbf{w}) = (\mathbf{u} + \mathbf{v}) + \mathbf{w}$
2. $\forall \mathbf{v}, \mathbf{w} \in V, \mathbf{v} + \mathbf{w} = \mathbf{w} + \mathbf{v}$
3. $\exists \mathbf{0} \in V$, the zero vector, such that $\mathbf{v} + \mathbf{0} = \mathbf{v}$ for all $\mathbf{v} \in V$
4. $\forall \mathbf{v} \in V$, there exists an element $\mathbf{w} \in V$, the additive inverse of $\mathbf{v}$, such that $\mathbf{v} + \mathbf{w} = \mathbf{0}$
5. $\forall \alpha \in \mathbb{R}$ and $\forall \mathbf{v}, \mathbf{w} \in V, \alpha(\mathbf{v} + \mathbf{w}) = \alpha\mathbf{v} + \alpha\mathbf{w}$
6. $\forall \alpha, \beta \in \mathbb{R}$ and $\forall \mathbf{v} \in V, (\alpha + \beta)\mathbf{v} = \alpha\mathbf{v} + \beta\mathbf{v}$
7. $\forall \alpha, \beta \in \mathbb{R}$ and $\forall \mathbf{v} \in V, \alpha(\beta\mathbf{v}) = (\alpha\beta)\mathbf{v}$
8. $\forall \mathbf{v} \in V, 1\mathbf{v} = \mathbf{v}$

Norms

Definition 11 (Norm)

Let V be a vector space over $\mathbb{F}$, and $\mathbf{v} \in V$ be a vector. The **norm** of $\mathbf{v}$, denoted $\|\mathbf{v}\|$, is a function from V to $\mathbb{R}_+$ that has the following properties:

1. For all $\mathbf{v} \in V$, $\|\mathbf{v}\| \geq 0$ with $\|\mathbf{v}\| = 0$ iff $\mathbf{v} = \mathbf{0}$
2. For all $\alpha \in \mathbb{F}$ and all $\mathbf{v} \in V$, $\|\alpha\mathbf{v}\| = |\alpha| \|\mathbf{v}\|$
3. For all $\mathbf{u}, \mathbf{v} \in V$, $\|\mathbf{u} + \mathbf{v}\| \leq \|\mathbf{u}\| + \|\mathbf{v}\|$

Let V be a vector space (for example, $\mathbb{R}^2$ or $\mathbb{R}^3$)

The **zero element** (or **zero vector**) is the vector $\mathbf{0} = (0, \dots, 0)$

The **additive inverse** of $\mathbf{v} = (v_1, \dots, v_n)$ is $-\mathbf{v} = (-v_1, \dots, -v_n)$

For $\mathbf{v} = (v_1, \dots, v_n) \in V$, the length (or Euclidean norm) of $\mathbf{v}$ is the **scalar**


$$\|\mathbf{v}\| = \sqrt{v_1^2 + \dots + v_n^2}$$

To **normalize** the vector $\mathbf{v}$ consists in considering $\tilde{\mathbf{v}} = \mathbf{v}/\|\mathbf{v}\|$, i.e., the vector in the same direction as $\mathbf{v}$ that has unit length

Standard basis vectors

Vectors $\mathbf{i} = (1, 0, 0)$, $\mathbf{j} = (0, 1, 0)$ and $\mathbf{k} = (0, 0, 1)$ are the **standard basis vectors** of $\mathbb{R}^3$. A vector $\mathbf{v} = (v_1, v_2, v_3)$ can then be written

$$\mathbf{v} = v_1 \mathbf{i} + v_2 \mathbf{j} + v_3 \mathbf{k}$$



For $V(\mathbb{R}^n)$, the standard basis vectors are usually denoted $\mathbf{e}_1, \dots, \mathbf{e}_n$, with

$$\mathbf{e}_k = (\underbrace{0, \dots, 0}_{k-1}, 1, \underbrace{0, \dots, 0}_{n-k+1})$$

Dot product

Definition 12 (Dot product)

Let $\mathbf{a} = (a_1, \dots, a_n) \in \mathbb{R}^n$, $\mathbf{b} = (b_1, \dots, b_n) \in \mathbb{R}^n$. The **dot product** of $\mathbf{a}$ and $\mathbf{b}$ is the **scalar**

$$\mathbf{a} \bullet \mathbf{b} = \sum_{i=1}^n a_i b_i = a_1 b_1 + \dots + a_n b_n$$

The dot product is a special case of **inner product**

Properties of the dot product

Theorem 13

For $\mathbf{a}, \mathbf{b}, \mathbf{c} \in \mathbb{R}^n$ and $\alpha \in \mathbb{R}$,

► $\mathbf{a} \bullet \mathbf{a} = \|\mathbf{a}\|^2$

(so $\mathbf{a} \bullet \mathbf{a} \geq 0$, with $\mathbf{a} \bullet \mathbf{a} = 0$ iff $\mathbf{a} = \mathbf{0}$)

► $\mathbf{a} \bullet \mathbf{b} = \mathbf{b} \bullet \mathbf{a}$

($\bullet$ is commutative)

► $\mathbf{a} \bullet (\mathbf{b} + \mathbf{c}) = \mathbf{a} \bullet \mathbf{b} + \mathbf{a} \bullet \mathbf{c}$

($\bullet$ distributive over $+$)

► $(\alpha \mathbf{a}) \bullet \mathbf{b} = \alpha(\mathbf{a} \bullet \mathbf{b}) = \mathbf{a} \bullet (\alpha \mathbf{b})$

► $\mathbf{0} \bullet \mathbf{a} = 0$

Some results stemming from the dot product

Theorem 14

If θ is the angle between the vectors $\mathbf{a}$ and $\mathbf{b}$, then

$$\mathbf{a} \bullet \mathbf{b} = \|\mathbf{a}\| \|\mathbf{b}\| \cos \theta$$

Corollary 15 (Cauchy-Schwarz inequality)

For any two vectors $\mathbf{a}$ and $\mathbf{b}$, we have

$$|\mathbf{a} \bullet \mathbf{b}| \leq \|\mathbf{a}\| \|\mathbf{b}\|$$

with equality if and only if $\mathbf{a}$ is a scalar multiple of $\mathbf{b}$, or one of them is $\mathbf{0}$.

Theorem 16

$\mathbf{a}$ and $\mathbf{b}$ are orthogonal if and only if $\mathbf{a} \bullet \mathbf{b} = 0$.


Scalar and vector projections

Scalar projection of $\mathbf{v}$ onto $\mathbf{a}$ (or component of $\mathbf{v}$ along $\mathbf{a}$):

$$\text{comp}_{\mathbf{a}} \mathbf{v} = \frac{\mathbf{a} \bullet \mathbf{v}}{\|\mathbf{a}\|}$$

Vector (or orthogonal) projection of $\mathbf{v}$ onto $\mathbf{a}$:

$$\text{proj}_{\mathbf{a}} \mathbf{v} = \left(\frac{\mathbf{a} \bullet \mathbf{v}}{\|\mathbf{a}\|} \right) \frac{\mathbf{a}}{\|\mathbf{a}\|} = \frac{\mathbf{a} \bullet \mathbf{v}}{\|\mathbf{a}\|^2} \mathbf{a}$$



Sets and logic

Complex numbers

Vectors and vector spaces

Linear systems and matrices

A crash course in multivariable calculus

Linear systems and matrices

Linear systems and matrices

Matrix arithmetic

Eigenpairs

Similarity and diagonalisation

Linear independence/Bases/Dimension

Linear systems

Definition 17 (Linear system)

A **linear system** of m equations in n unknowns takes the form

$$\begin{array}{ccccccccc} a_{11}x_1 & + & a_{12}x_2 & + & \cdots & + & a_{1n}x_n & = & b_1 \\ a_{21}x_1 & + & a_{22}x_2 & + & \cdots & + & a_{2n}x_n & = & b_2 \\ \vdots & & \vdots & & \vdots & & \vdots & & \vdots \\ a_{m1}x_1 & + & a_{m2}x_2 & + & \cdots & + & a_{mn}x_n & = & b_n \end{array} \quad (1)$$

The a_{ij} , x_j and b_j could be in $\mathbb{R}$ or $\mathbb{C}$, although here we typically assume they are in $\mathbb{R}$

The aim is to find $x_1, x_2, \dots, x_n$ that satisfy all equations simultaneously

Theorem 18 (Nature of solutions to a linear system)

A linear system can have

- ▶ *no solution*
- ▶ *a unique solution*
- ▶ *infinitely many solutions*

Operations on linear systems

You learned to manipulate linear systems using

- ▶ Gaussian elimination
- ▶ Gauss-Jordan elimination

with the aim to put the system in **row echelon form** (REF) or **reduced row echelon form** (RREF)

Matrices and linear systems

Writing

$$A = \begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{pmatrix}, \quad \mathbf{x} = \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} \quad \text{and} \quad \mathbf{b} = \begin{pmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{pmatrix}$$

where A is an $m \times n$ **matrix**, $\mathbf{x}$ and $\mathbf{b}$ are n (column) **vectors** (or $n \times 1$ matrices), then the linear system in the previous slide takes the form

$$A\mathbf{x} = \mathbf{b}$$

Notation for vectors

We usually assume vectors are column vectors and thus write, e.g.,

$$\mathbf{x} = \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} = (x_1, x_2, \dots, x_n)^T$$

Here, T is the **transpose operator** (more on this soon)

Consider the system

$$A\mathbf{x} = \mathbf{b}$$

If $\mathbf{b} = \mathbf{0}$, the system is **homogeneous** and always has the solution $\mathbf{x} = \mathbf{0}$ and so the “no solution” option in Theorem 18 goes away

Linear systems and matrices

Linear systems and matrices

Matrix arithmetic

Eigenpairs

Similarity and diagonalisation

Linear independence/Bases/Dimension

Definition 19 (Matrix)

An m -by- n or $m \times n$ matrix is a rectangular array of elements of $\mathbb{R}$ or $\mathbb{C}$ with m rows and n columns,

$$A = [a_{ij}] = \begin{pmatrix} a_{11} & \cdots & a_{1n} \\ \vdots & & \vdots \\ a_{m1} & \cdots & a_{mn} \end{pmatrix}$$

We always list indices as “row,column”

We denote $\mathcal{M}_{mn}(\mathbb{F})$ or $\mathbb{F}^{mn}$ the set of $m \times n$ matrices with entries in $\mathbb{F} = \{\mathbb{R}, \mathbb{C}\}$. Often, we omit $\mathbb{F}$ in $\mathcal{M}_{mn}$ if the nature of $\mathbb{F}$ is not important

When $m = n$, we usually write $\mathcal{M}_n$

Basic matrix arithmetic

Let $A \in \mathcal{M}_{mn}$, $B \in \mathcal{M}_{mn}$ be matrices (of the same size) and $c \in \mathbb{F} = \{\mathbb{R}, \mathbb{C}\}$ be a scalar

► **Scalar multiplication**

$$cA = [ca_{ij}]$$

► **Addition**

$$A + B = [a_{ij} + b_{ij}]$$

► **Subtraction** (addition of $-B = (-1)B$ to A)

$$A - B = A + (-1)B = [a_{ij} + (-1)b_{ij}] = [a_{ij} - b_{ij}]$$

► **Transposition** of A gives a matrix $A^T \in \mathcal{M}_{nm}$ with

$$A^T = [a_{ji}], \quad j = 1, \dots, n, \quad i = 1, \dots, m$$

Matrix multiplication

The (matrix) **product** of A and B , AB , requires the “inner dimensions” to match, i.e., the number of columns in A must equal the number of rows in B

Suppose that is the case, i.e., let $A \in \mathcal{M}_{mn}$, $B \in \mathcal{M}_{np}$. Then the i, j entry in $C := AB$ takes the form

$$c_{ij} = \sum_{k=1}^n a_{ik} b_{kj}$$

Recall that the matrix product is not commutative, i.e., in general, $AB \neq BA$ (when both those products are defined, i.e., when $A, B \in \mathcal{M}_n$)

Special matrices

Definition 20 (Zero and identity matrices)

The **zero** matrix is the matrix 0_{mn} whose entries are all zero. The **identity** matrix is a square $n \times n$ matrix $\mathbb{I}_n$ with all entries on the main diagonal equal to one and all off diagonal entries equal to zero

Definition 21 (Symmetric matrix)

A square matrix $A \in \mathcal{M}_n$ is **symmetric** if $\forall i, j = 1, \dots, n, a_{ij} = a_{ji}$. In other words, $A \in \mathcal{M}_n$ is symmetric if $A = A^T$

Making symmetric matrices

Theorem 22

1. *If $A \in \mathcal{M}_n$, then $A + A^T$ is symmetric*
2. *If $A \in \mathcal{M}_{mn}$, then $AA^T \in \mathcal{M}_m$ and $A^T A \in \mathcal{M}_n$ are symmetric*

Proof of Theorem 22

X symmetric $\iff X = X^T$, so use $X =$ the matrix whose symmetric property you want to check

1. True if $A + A^T = (A + A^T)^T$. We have

$$(A + A^T)^T = A^T + (A^T)^T = A^T + A = A + A^T$$

2. AA^T symmetric if $AA^T = (AA^T)^T$. We have

$$(AA^T)^T = (A^T)^T A^T = AA^T$$

$A^T A$ works similarly

Determinants

Definition 23 (Determinant)

Let $A \in \mathcal{M}_n$ with $n \geq 2$. The **determinant** of A is the *scalar*

$$\det(A) = |A| = \sum_{j=1}^n a_{ij} C_{ij}$$

where $C_{ij} = (-1)^{i+j} \det(A_{ij})$ is the (i, j) -**cofactor** of A and A_{ij} is the submatrix of A from which the i th row and j th column have been removed

This is a cofactor expansion along the i th row

This is a recursive formula: it gives result in terms of $n-1$ $\mathcal{M}_{n-1}$ matrices, to which it must in turn be applied, all the way down to

$$\det \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} = a_{11} a_{22} - a_{12} a_{21}$$

Two special matrices and their determinants

Definition 24

$A \in \mathcal{M}_n$ is **upper triangular** if $a_{ij} = 0$ when $i > j$, **lower triangular** if $a_{ij} = 0$ when $j > i$, **triangular** if it is *either* upper or lower triangular and **diagonal** if it is *both* upper and lower triangular

When A diagonal, we often write $A = \text{diag}(a_{11}, a_{22}, \dots, a_{nn})$

Theorem 25

Let $A \in \mathcal{M}_n$ be triangular or diagonal. Then

$$\det(A) = \prod_{i=1}^n a_{ii} = a_{11} a_{22} \cdots a_{nn}$$

Inversion/Singularity

Definition 26 (Matrix inverse)

$A \in \mathcal{M}_n$ is **invertible** (or **nonsingular**) if $\exists A^{-1} \in \mathcal{M}_n$ s.t.

$$AA^{-1} = A^{-1}A = \mathbb{I}$$

A^{-1} is the **inverse** of A . If A^{-1} does not exist, A is **singular**

Theorem 27

Let $A \in \mathcal{M}_n$, $\mathbf{x}, \mathbf{b} \in \mathbb{F}^n$. Then

- ▶ A invertible $\iff \det(A) \neq 0$
- ▶ If A invertible, A^{-1} is unique
- ▶ If A invertible, then $A\mathbf{x} = \mathbf{b}$ has the unique solution $\mathbf{x} = A^{-1}\mathbf{b}$

Revisiting matrix arithmetic

With addition, subtraction, scalar multiplication, multiplication, transposition and inversion, you can perform arithmetic on matrices essentially as on scalar, if you bear in mind a few rules

- ▶ The sizes have to be compatible
- ▶ The order is important since matrix multiplication is not commutative
- ▶ Transposition and inversion change the order of products:

$$(AB)^T = B^T A^T \text{ and } (AB)^{-1} = B^{-1} A^{-1}$$

Linear systems and matrices

Linear systems and matrices

Matrix arithmetic

Eigenpairs

Similarity and diagonalisation

Linear independence/Bases/Dimension

Eigenvalues / Eigenvectors / Eigenpairs

Definition 28

Let $A \in \mathcal{M}_n$. A vector $\mathbf{x} \in \mathbb{F}^n$ such that $\mathbf{x} \neq \mathbf{0}$ is an **eigenvector** of A if $\exists \lambda \in \mathbb{F}$ called an **eigenvalue**, s.t.

$$A\mathbf{x} = \lambda\mathbf{x}$$

A couple $(\lambda, \mathbf{x})$ with $\mathbf{x} \neq \mathbf{0}$ s.t. $A\mathbf{x} = \lambda\mathbf{x}$ is an **eigenpair**

Parallel eigenvectors

Theorem 29

If $(\lambda, \mathbf{x})$ eigenpair, then for $c \neq 0$, $(\lambda, c\mathbf{x})$ also eigenpair

Proof.

Assume $(\lambda, \mathbf{x})$ is an eigenpair of $A \in \mathcal{M}_n$, i.e., $\mathbf{x} \neq \mathbf{0}$ and $A\mathbf{x} = \lambda\mathbf{x}$

Take the nonzero scalar $c \in \mathbb{F}$. Then $A(c\mathbf{x}) = cA\mathbf{x}$ because scalars move freely in vector/matrix products. Therefore $A(c\mathbf{x}) = cA\mathbf{x} = c\lambda\mathbf{x}$ since $A\mathbf{x} = \lambda\mathbf{x}$

So finally, moving c again, $A(c\mathbf{x}) = \lambda(c\mathbf{x})$, i.e., $c\mathbf{x}$ is an eigenvector of A associated to λ



Linear systems and matrices

Linear systems and matrices

Matrix arithmetic

Eigenpairs

Similarity and diagonalisation

Linear independence/Bases/Dimension

Similarity

Definition 30 (Similarity)

$A, B \in \mathcal{M}_n$ are **similar** ($A \sim B$) if $\exists P \in \mathcal{M}_n$ invertible s.t.

$$P^{-1}AP = B$$

Theorem 31 ($\sim$ is an equivalence relation)

$A, B, C \in \mathcal{M}_n$, then

- ▶ $A \sim A$ ($\sim$ **reflexive**)
- ▶ $A \sim B \implies B \sim A$ ($\sim$ **symmetric**)
- ▶ $A \sim B$ and $B \sim C \implies A \sim C$ ($\sim$ **transitive**)

Similarity (cont.)

Theorem 32

$A, B \in \mathcal{M}_n$ with $A \sim B$. Then

- ▶ $\det A = \det B$
- ▶ A invertible $\iff B$ invertible
- ▶ A and B have the same eigenvalues

Diagonalisation

Definition 33 (Diagonalisability)

$A \in \mathcal{M}_n$ is **diagonalisable** if $\exists D \in \mathcal{M}_n$ diagonal s.t. $A \sim D$

In other words, $A \in \mathcal{M}_n$ is diagonalisable if there exists a diagonal matrix $D \in \mathcal{M}_n$ and a nonsingular matrix $P \in \mathcal{M}_n$ s.t. $P^{-1}AP = D$

Could of course write $PAP^{-1} = D$ since P invertible, but $P^{-1}AP$ makes more sense for computations

Theorem 34

$A \in \mathcal{M}_n$ diagonalisable $\iff A$ has n linearly independent eigenvectors

Corollary 35 (Sufficient condition for diagonalisability)

$A \in \mathcal{M}_n$ has all its eigenvalues distinct $\implies A$ diagonalisable

For $P^{-1}AP = D$: in P , put the linearly independent eigenvectors as columns and in D , the corresponding eigenvalues

Linear systems and matrices

Linear systems and matrices

Matrix arithmetic

Eigenpairs

Similarity and diagonalisation

Linear independence/Bases/Dimension

Linear combination and span

Definition 36 (Linear combination)

Let V be a vector space. A **linear combination** of a set $\{\mathbf{v}_1, \dots, \mathbf{v}_k\}$ of vectors in V is a *vector*

$$c_1 \mathbf{v}_1 + \dots + c_k \mathbf{v}_k$$

where $c_1, \dots, c_k \in \mathbb{F}$

Definition 37 (Span)

The set of all linear combinations of a set of vectors $\mathbf{v}_1, \dots, \mathbf{v}_k$ is the **span** of $\{\mathbf{v}_1, \dots, \mathbf{v}_k\}$,

$$\text{span}(\mathbf{v}_1, \dots, \mathbf{v}_k) = \{c_1 \mathbf{v}_1 + \dots + c_k \mathbf{v}_k : c_1, \dots, c_k \in \mathbb{F}\}$$

Finite/infinite-dimensional vector spaces

Theorem 38

The span of a set of vectors in V is the smallest subspace of V containing all the vectors in the set

Definition 39 (Set of vectors spanning a space)

If $\text{span}(\mathbf{v}_1, \dots, \mathbf{v}_k) = V$, we say $\mathbf{v}_1, \dots, \mathbf{v}_k$ **spans** V

Definition 40 (Dimension of a vector space)

A vector space V is **finite-dimensional** if some set of vectors in it spans V . A vector space V is **infinite-dimensional** if it is not finite-dimensional

Linear (in)dependence

Definition 41 (Linear independence/Linear dependence)

A set $\{\mathbf{v}_1, \dots, \mathbf{v}_k\}$ of vectors in a vector space V is **linearly independent** if

$$(c_1 \mathbf{v}_1 + \dots + c_k \mathbf{v}_k = \mathbf{0}) \Leftrightarrow (c_1 = \dots = c_k = 0),$$

where $c_1, \dots, c_k \in \mathbb{F}$. A set of vectors is **linearly dependent** if it is not linearly independent.

If linearly dependent, assume w.l.o.g. that $c_1 \neq 0$, then

$$\mathbf{v}_1 = -\frac{c_2}{c_1} \mathbf{v}_2 - \dots - \frac{c_k}{c_1} \mathbf{v}_k$$

i.e., $\mathbf{v}_1$ is a linear combination of the other vectors in the set

Theorem 42

*Let V be a finite-dimensional vector space. Then the **cardinal** (number of elements) of every linearly independent set of vectors is less than or equal to the number of elements in every spanning set of vectors*

E.g., in $\mathbb{R}^3$, a set with 4 or more vectors is automatically linearly dependent

Basis

Definition 43 (Basis)

Let V be a vector space. A **basis** of V is a set of vectors in V that is both linearly independent and spanning

Theorem 44 (Criterion for a basis)

A set $\{\mathbf{v}_1, \dots, \mathbf{v}_k\}$ of vectors in a vector space V is a basis of $V \iff \forall \mathbf{v} \in V, \mathbf{v}$ can be written uniquely in the form

$$\mathbf{v} = c_1 \mathbf{v}_1 + \dots + c_k \mathbf{v}_k,$$

where $c_1, \dots, c_k \in \mathbb{F}$

More on bases

Theorem 45 (Basis of finite-dimensional vector space)

Every finite-dimensional vector space has a basis

Theorem 46

Any two bases of a finite-dimensional vector space have the same number of vectors

Definition 47 (Dimension)

The **dimension** $\dim V$ of a finite-dimensional vector space V is the number of vectors in any basis of the vector space

Theorem 48 (Dimension of a subspace)

Let V be a finite-dimensional vector space and $U \subset V$ be a subspace of V . Then $\dim U \leq \dim V$


Constructing bases

Theorem 49

Let V be a finite-dimensional vector space. Then every linearly independent set of vectors in V with $\dim V$ elements is a basis of V

Theorem 50

Let V be a finite-dimensional vector space. Then every spanning set of vectors in V with $\dim V$ elements is a basis of V

A robot stands in a vast, arid desert landscape under a hazy, orange-tinted sky. In the background, a large, leafless tree with a complex, branching structure stands prominently. The ground is sandy and dotted with small, dark rocks. The overall atmosphere is desolate and contemplative.

Sets and logic

Complex numbers

Vectors and vector spaces

Linear systems and matrices

A crash course in multivariable calculus


A crash course in multivariable calculus

What is Multivariable Calculus?

Optimization with Constraints

The Lagrangian Method


Conclusion



One dimension

MATH 1500 & 1700 deal with functions of one variable, like $f(x) = x^2$


A 1D function



Multivariable calculus

Multivariable calculus extends this to functions of two or more variables, like $f(x, y) = x^2 + y^2$

A 2D function surface



Partial derivatives


How do we measure the “slope” on a 3D surface?

A **partial derivative** measures the slope in a direction parallel to one of the axes

- ▶ $\frac{\partial f}{\partial x}$ measures height change as we move only in the x direction. Treat y as a constant
- ▶ $\frac{\partial f}{\partial y}$ measures height change as we move only in the y direction. Treat x as a constant

Partial derivatives

Slices for Partial Derivatives



The Steepest path: the gradient

The **gradient**, denoted ∇f , is a vector that combines all the partial derivatives:


$$\nabla f(x, y) = \left(\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y} \right)$$

What does it tell us?

- ▶ **Direction:** it points in the direction of the *steepest ascent*
- ▶ **Magnitude:** its length represents the steepness of that ascent


Follow the gradient

Gradient Field



At a peak or a valley (a local max/min), the ground is flat. So, $\nabla f = (0, 0)$

A crash course in multivariable calculus

A stylized illustration of a robot standing in a desert landscape. The robot is on the left, facing right. In the background, there is a large, leafless tree on the right and mountains in the distance. The ground is sandy with some small rocks and a fallen log in the foreground. The sky is a mix of blue and orange, suggesting a sunset or sunrise.

What is Multivariable Calculus?

Optimization with Constraints

The Lagrangian Method

Conclusion

The real-world problem

Often, we want to maximize or minimize a function, but we don't have unlimited freedom. We have **constraints**

- ▶ Maximize the profit of your company... *subject to a limited budget*
- ▶ Minimize the material used for a can... *that must hold a specific volume*
- ▶ Find the highest point on a mountain... *while staying on a specific trail*


Setting the gradient to zero ($\nabla f = 0$) finds the highest point on the whole mountain, which might not be on our trail!

Visualizing the problem

Imagine our function $f(x, y)$ is the altitude on a map (contour lines)

Our constraint, $g(x, y) = c$, is a specific path we must walk on

Optimization with a Constraint




We are looking for the highest (or lowest) point *along the red path*

The key insight

At the optimal point on the path, the path will be perfectly **tangent** to the contour line of the surface

Tangency at the Optimum



Why? If the path crossed the contour line, you could move along the path to get to a higher (or lower) contour

Mathematically, this tangency means the gradient vectors of the function and the constraint are **parallel**

$$\nabla f = \lambda \nabla g$$

The scalar λ (lambda) is called the **Lagrange multiplier**

A crash course in multivariable calculus

What is Multivariable Calculus?

Optimization with Constraints

The Lagrangian Method

Conclusion



The Lagrangian function

The condition $\nabla f = \lambda \nabla g$ is clever, but solving it can be messy

Instead, we combine our function and constraint into a single, new function called the **Lagrangian**

$$\mathcal{L}(x, y, \lambda) = f(x, y) - \lambda[g(x, y) - c]$$

- ▶ $f(x, y)$ the function we want to optimize
- ▶ $g(x, y) = c$ the constraint we must follow
- ▶ λ the Lagrange multiplier

Finding the unconstrained optimum of $\mathcal{L}$ solves the original constrained problem!

The method – step-by-step

To find the optimum of the Lagrangian $\mathcal{L}(x, y, \lambda)$, we find where its gradient is zero

We take the partial derivative with respect to *all* its variables (x , y , and λ) and set them to zero

$$1. \quad \frac{\partial \mathcal{L}}{\partial x} = \frac{\partial f}{\partial x} - \lambda \frac{\partial g}{\partial x} = 0$$

$$2. \quad \frac{\partial \mathcal{L}}{\partial y} = \frac{\partial f}{\partial y} - \lambda \frac{\partial g}{\partial y} = 0$$

$$3. \quad \frac{\partial \mathcal{L}}{\partial \lambda} = -(g(x, y) - c) = 0 \implies g(x, y) = c$$

The first two equations rearrange to $\nabla f = \lambda \nabla g$ and the third equation is the original constraint

Example: Fencing a Field

Problem: You have 40 meters of fence. What is the largest rectangular area you can enclose?

- ▶ **Maximize Area:** $A(x, y) = xy$
- ▶ **Constraint (Perimeter):** $2x + 2y = 40$

1. Form the Lagrangian:

$$\mathcal{L}(x, y, \lambda) = xy - \lambda(2x + 2y - 40)$$

2. Take Partial Derivatives:

- ▶ $\frac{\partial \mathcal{L}}{\partial x} = y - 2\lambda = 0 \implies y = 2\lambda$
- ▶ $\frac{\partial \mathcal{L}}{\partial y} = x - 2\lambda = 0 \implies x = 2\lambda$
- ▶ $\frac{\partial \mathcal{L}}{\partial \lambda} = -(2x + 2y - 40) = 0$

Example: Solution

From the first two equations, we see that $x = y$

Now, substitute this into the third equation (the constraint):


$$2x + 2(x) = 40$$

$$4x = 40$$

$$x = 10$$

Since $x = y$, we have $y = 10$, i.e., optimal dimensions are 10m by 10m (a square), giving a maximum area of 100 m^2

A crash course in multivariable calculus

A stylized illustration of a humanoid robot standing in a vast, arid desert landscape. The robot is positioned on the left side of the frame, facing right. It has a boxy, mechanical design with visible joints and a small head. In the background, a large, leafless tree stands prominently on the right, its intricate branch structure silhouetted against a hazy, orange-tinted sky. The ground is sandy and dotted with small rocks and shrubs. Distant mountains are visible on the horizon under a soft, glowing light, possibly from a low sun or moon.

What is Multivariable Calculus?

Optimization with Constraints

The Lagrangian Method

Conclusion

What does λ mean?

The Lagrange multiplier λ has a very useful interpretation

It tells you how much the optimal value of your function f will change if you slightly relax the constraint c

$$\lambda = \frac{df_{\text{optimal}}}{dc}$$

In our example: If we had 41 meters of fence instead of 40 (so c changes by 1), how much would the max area increase?

$x = y = 2\lambda$, so $\lambda = x/2 = 10/2 = 5$. The maximum area would increase by approximately 5 m^2