Least squares problems

MATH 2740 – Mathematics of Data Science – Unit U03

Julien Arino
julien.arino@umanitoba.ca

Department of Mathematics @ University of Manitoba

Fall 2026

The University of Manitoba campuses are located on original lands of Anishinaabeg, Ininew, Anisininew, Dakota and Dene peoples, and on the National Homeland of the Red River Métis. We respect the Treaties that were made on these territories, we acknowledge the harms and mistakes of the past, and we dedicate ourselves to move forward in partnership with Indigenous communities in a spirit of Reconciliation and collaboration.

Outline

Getting the Canadian census data

Least squares problems

Example – Canadian population data

Getting the Canadian census data

Least squares problems

Example – Canadian population data

Remark

This is the slide version of `Canada-census-get-plot-data.Rmd` in the folder `CODE` of the course repository

The `html` file generated there (link) is much easier to read than these slides

Canadian census data

We want to consider the evolution of the population of Canada through time

For this, we grab the Canadian census data

Search for (Google) “Canada historical census data csv", since csv (comma separated values) is a very easy format to use with R

Here, we find a csv for 1851 to 1976

We follow the link to Table A2-14, where we find another link, this time to a csv file. This is what we use in R

Grabing the Canadian census data

The function `read.csv` reads in a file (potentially directly from the web)

```
prefix = "https://www150.statcan.gc.ca/n1/en/pub/11-516-x/sectiona/"
page = "A2_14-eng.csv?st=L7vSnqio"
url = paste0(prefix, page)
data_old = read.csv(url)
```

This directly assigns the result to the variable `data`

We then use the function `head` to show the first few lines in the result

```
head(data_old)
```

```
##    X Series.A2.14.
## 1 NA
## 2 NA          Year
## 3 NA
## 4 NA
## 5 NA
## 6 NA
##   Population.of.Canada..by.province..census.dates..1851.to.1976 X.1
## 1                                                                NA
## 2                                                        Canada  NA Newfo
## 3                                                                NA
## 4                                                                NA
## 5                                                             2  NA
## 6                                                                NA
##      X.3 X.4    X.5       X.6    X.7     X.8       X.9 X.10    X.11 X.12
## 1             NA                                          NA           NA
```

Obviously, this does not make a lot of sense. This is normal: take a look at the first few lines in the file. They take the form

```
head(data_old, n = 1)

##    X Series.A2.14.
## 1 NA
##   Population.of.Canada..by.province..census.dates..1851.to.1976 X.1 X.2 X
## 1                                                                NA
##   X.5 X.6 X.7 X.8 X.9 X.10 X.11 X.12 X.13 X.14 X.15 X.16 X.17 X.18 X.19 X
## 1                      NA        NA        NA        NA             NA
```

This happens often: the first few lines are here to describe the information contained in the data set, i.e., the so-called **metadata**

It is easy to deal with this: the function `read.csv` takes the optional argument `skip`, which indicates how many lines to skip at the beginning of the file

The second line is also empty, so let us skip it too

```
data_old = read.csv(url, skip = 2)
head(data_old, n = 2)
```

```
##    X Year Canada X.1 Newfound. Prince X.2   Nova       New Quebec Ontario
## 1 NA              NA      land Edward  NA Scotia Brunswick
## 2 NA              NA           Island  NA
##   Manitoba X.3 Saskat. X.4 Alberta X.5  British X.6     Yukon   Northwest
## 1           NA  chewan  NA          NA Columbia  NA Territory Territories
## 2           NA          NA          NA           NA
##   X.8
## 1  NA
## 2  NA
```

To make things legible, the table authors used 3 rows to write long names (e.g., Prince Edward Island is written over 3 rows)

Note, however, that `read.csv` has rightly picked up on the first row being the column names

Because we are only interested in the total population of the country and the year, let us get rid of the first 4 rows and of all columns except the second (Year) and third (Canada)

```
data_old = data_old[5:dim(data_old)[1], 2:3]
head(data_old, n=4)

##   Year     Canada
## 5 1976 22,992,604
## 6
## 7 1971 21,568,311
## 8 1966 20,014,880
```

Still not perfect...

```
head(data_old, n=3)

##   Year     Canada
## 5 1976 22,992,604
## 6
## 7 1971 21,568,311

tail(data_old, n=2)

##                            Year Canada
## 28 see notes to series A15-53.
## 29                1848 figure.
```

1. there are some empty rows
2. the last few rows need to be removed too, they contain remarks about the data
3. the population counts contain commas
4. it would be better if years were increasing

Fixing these issues

1 and 2 are easy: remark that the Canada column is empty for both issues. Entries in the column are strings. Looking for empty content therefore means looking for empty strings

So to fix 1 and 2, we keep the rows where Canada does not equal the empty string

To get rid of commas, we just need to substitute an empty string for ","

To sort, we find the order for the years and apply it to the entire table

Finally, as remarked above, for now, both the year and the population are considered as strings. To plot, we will have to indicate that these are numbers, not characters

```r
data_old = data_old[which(data_old$Canada != ""),]
data_old$Canada = gsub(",", "", data_old$Canada)
order_data = order(data_old$Year)
data_old = data_old[order_data,]
data_old$Year = as.numeric(data_old$Year)
data_old$Canada = as.numeric(data_old$Canada)
head(data_old)
```

```
##    Year  Canada
## 23 1851 2436297
## 22 1861 3229633
## 21 1871 3689257
## 20 1881 4324810
## 19 1891 4833239
## 17 1901 5371315
```


Row numbers are a little weird

```
row.names(data_old) = 1:dim(data_old)[1]
head(data_old)
```

```
##   Year  Canada
## 1 1851 2436297
## 2 1861 3229633
## 3 1871 3689257
## 4 1881 4324810
## 5 1891 4833239
## 6 1901 5371315
```

Well, that looks about right! Let's see what this looks like in a graph

```r
plot(data_old$Year, data_old$Canada,
    type = "b", lwd = 2,
    xlab = "Year", ylab = "Population")
```



But wait, this is only to 1976..!

Looking around, we find another table here with a download csv link (this one)

StatCan has added Cloudflare bot protection, so automated downloads from R fail. We download it via browser and save to `DATA/98-315-XWE2011001-001.csv`

```
data_new = read.csv("../DATA/98-315-XWE2011001-001.csv",
                    fileEncoding = "latin1")
```

The table is 720KB, so surely there must be more to this than just the population

To get a sense of that, we dump a few random rows

```
data_new[sort(unique(c(1,sample(nrow(data_new), 4)))), ]
```

```
##                 GEOGRAPHY.NAME
## 1                       Canada
## 453  Newfoundland and Labrador
## 3126            Trois-Rivières
## 5586                    London
## 5636                   Windsor
##                                                    CHARACTERISTIC YEAR.S
## 1                                        Population (in thousands)    1956
## 453                    Population by single years of age: 76 years    2011
## 3126                                % of three-person households    2006
## 5586 % of private households containing a couple without children    2011
## 5636                     % of the population aged 0 to 14 years    2011
##        TOTAL FLAG_TOTAL
## 1    16081.0
## 453   3145.0
## 3126    14.3
```

Flat tables

This is a **flat table**: each row contains a piece of information, the columns describe what this piece of information is about

This has *way more* information than we need: we just want the population of Canada and here we get 9960 rows over 213 characteristics

Also, population is expressed in thousands, so once we selected what we want, we need to multiply by 1,000

Selecting rows

We want the rows where the geography is “Canada” and the characteristic is “Population (in thousands)”

Find indices of rows that satisfy the first criterion and those that satisfy the second; intersecting these two sets of indices, we have selected the rows we want

```
idx_CAN = which(data_new$GEOGRAPHY.NAME == "Canada")
idx_char = which(data_new$CHARACTERISTIC == "Population (in thousands)")
idx_keep = intersect(idx_CAN, idx_char)
```

Let us keep only these rows

```
data_new = data_new[idx_keep,]
head(data_new, n = 8)
```

```
##   GEOGRAPHY.NAME            CHARACTERISTIC YEAR.S. TOTAL FLAG_TOTAL
## 1         Canada Population (in thousands)    1956 16081
## 2         Canada Population (in thousands)    1961 18238
## 3         Canada Population (in thousands)    1966 20015
## 4         Canada Population (in thousands)    1971 21568
## 5         Canada Population (in thousands)    1976 22993
## 6         Canada Population (in thousands)    1981 24343
## 7         Canada Population (in thousands)    1986 25309
## 8         Canada Population (in thousands)    1991 27297
```

We want to concatenate this data frame with the one from earlier

To do this, we need the two data frames to have the same number of columns, same column names and same entry types (notice that `YEAR.S.` in `data_new` is a column of characters)


What remains to be done

- Rename the columns in the pruned old data (data_pruned) to `year` and `population`. Personally, I prefer lowercase column names.. and `population` is more informative than `Canada`
- Keep only the relevant columns in `data_new`, rename them accordingly and multiply population by 1,000 there
- Transform year in `data_new` to numbers
- We already have data up to and including 1976 in `data_old`, so get rid of that in `data_new`
- Append the rows of `data_new` to those of `data_pruned`

```r
colnames(data_old) = c("year", "population")
data_new = data_new[,c("YEAR.S.","TOTAL")]
colnames(data_new) = c("year", "population")
data_new$year = as.numeric(data_new$year)
data_new = data_new[which(data_new$year>1976),]
data_new$population = data_new$population*1000

data = rbind(data_old,data_new)
```

```r
plot(data$year, data$population,
    type = "b", lwd = 2,
    xlab = "Year", ylab = "Population")
```



Save the processed data

In case we need the data elsewhere, we save the data to a `csv` file

```
write.csv(data, file = "../CODE/Canada_census.csv")
```

Using `readr` saves the data without row numbers (by default), so we can do this instead

```
readr::write_csv(data, file = "../CODE/Canada_census.csv")
```

Getting the Canadian census data

Least squares problems

Example – Canadian population data

Least squares problems

Initial considerations

The least squares problem

Fitting something more complicated


We just collected the census data for Canada

Suppose we want to predict the population of Canada in 20 years given the historical population growth seen in the previous plot. What can we do?

If there were just two points, we could easily "drive" a line through these two points. However, we have much more than two points, so we will use *fitting*, *i.e.*, try to make a curve come as close to possible to the points

We start with a line, giving rise to **linear least squares**

Least squares approximation – A trivial case



We want to find the equation of a line $y = a + bx$ that goes through these two points, i.e., we seek a and b such that

$$3 = a + b$$
$$5 = a + 2b$$

i.e., they satisfy $y = a + bx$ for $(x, y) = (1, 3)$ and $(x, y) = (2, 5)$

This is a linear system with 2 equations and 2 unknowns a and b

$$\begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} 3 \\ 5 \end{pmatrix}$$

We know from Theorem **??** (Linear algebra in a nutshell) that this system has a unique solution if the matrix


$$M = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix}$$

is invertible


$\det(M) = 1$, so we are good, we'll find a and b easily..

```
A = matrix(c(1,1,1,2), nr = 2, nc = 2, byrow = TRUE)
rhs = matrix(c(3,5), nr = 2, nc =1)
coefs = solve(A,rhs)
coefs
```

```
##      [,1]
## [1,]    1
## [2,]    2
```


y
0
2
4
6
0.0
0.5
1.0
1.5
2.0
2.5
3.0
x

Now let's add another point



These points are clearly not colinear, so there is not one line going through the 3

We end up with an *overdetermined* system

$$
\begin{aligned}
3 &= a + b \\
5 &= a + 2b \\
4 &= a + 3b
\end{aligned}
$$


i.e.,

$$
\begin{pmatrix} 1 & 1 \\ 1 & 2 \\ 1 & 3 \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} 3 \\ 5 \\ 4 \end{pmatrix}
$$

We have verified visually that the points are not colinear, so this system has no solution


(If you had to do it for good, you consider two vectors stemming from these 3 points and compute the angle between them or check that one is a multiple of the other)

So let us instead try to find the line that comes "closest" to the 3 points



Obviously, not sensational..

Let's eyeball it – Less steep and higher y-intercept



The blue line looks better, but how do we quantify this?

How do we find "how far away" we are?

- We could use projections onto the line (which we know minimises the distance)

- However, this will be a problem if we later decide that rather than a straight line, we want to use something more "funky" like a quadratic or an exponential

Compare, for a given value x, the distance between the true value y and the value of y obtained using the curve (line, here) that we use to fit the data

Let (x_i, y_i) be the data points, i.e., here, $(x_1, y_1) = (1, 3)$, $(x_2, y_2) = (2, 5)$ and $(x_3, y_3) = (3, 4)$

Suppose we use a line with equation $y = a + bx$ and that we pick a value for a and b. Then at x_1,

$$\tilde{y}_1 = a + bx_1 = a + b$$

at x_2

$$\tilde{y}_2 = a + bx_2 = a + 2b$$

and at x_3,

$$\tilde{y}_3 = a + bx_3 = a + 3b$$

Consider x_1, for instance. The error we made by using the line with coefficients (a, b) is $\overrightarrow{(x_1, y_1)(x_1, \tilde{y}_1)}$

For future use, let us create a function for $y = a_0 + a_1 x$

```
my_line = function(x, a_0, a_1){
    return(a_0 + a_1*x)
}
```

Functions are super useful when programming

```
my_line(1,2,3)
```


```
## [1] 5
```

```
my_line(a_0 = 2, a_1 = 3, x = 1)
```

```
## [1] 5
```

```
my_line(x = c(1,2,3), a_0 = 2, a_1 = 3)
```

```
## [1]  5  8 11
```


y
0
1
2
3
4
5
6
0.0
0.5
1.0
1.5
2.0
2.5
3.0
3.5
x

Let us return to the error

$$\overrightarrow{(x_1, y_1)(x_1, \tilde{y}_1)}$$

We have

$$\overrightarrow{(x_1, y_1)(x_1, \tilde{y}_1)} = (x_1 - x_1, y_1 - \tilde{y}_1) = (0, y_1 - \tilde{y}_1)$$

Let us call

$$\varepsilon_1 = y_1 - \tilde{y}_1$$

We can compute ε_2 and ε_3 too. And we can then form the **error vector**

$$\mathbf{e} = (\varepsilon_1, \varepsilon_2, \varepsilon_3)^T$$

The norm of $\mathbf{e}$, $\|\mathbf{e}\|$, then tells us how much error we are making for the choice of (a, b) we are using

The norm of $\|\mathbf{e}\|$ tells us how much error we are making for the choice of (a, b) we are using

So our objective is to find (a, b) such that $\|\mathbf{e}\|$ is minimal

We could use various norms, but the Euclidean norm has some very interesting properties, so we use

$$\|\mathbf{e}\| = \sqrt{\varepsilon_1^2 + \varepsilon_2^2 + \varepsilon_3^2}$$

The linear least squares problem

Given a collection of data points $(x_1, y_1), \ldots, (x_n, y_n)$, find the coefficients a, b of the line $y = a + bx$ such that

$$\|\mathbf{e}\| = \sqrt{\varepsilon_1^2 + \cdots + \varepsilon_n^2} = \sqrt{(y_1 - \tilde{y}_1)^2 + \cdots + (y_n - \tilde{y}_n)^2}$$

is minimal, where $\tilde{y}_i = a + bx_i$, for $i = 1, \ldots, n$

Let us first hack a brute force solution! (For the example we have been using this far)

We have our three points in the list `points`, the function `my_line` that computes the value $\tilde{y}$ given x and a, b, so let us make a new function that, given a, b, computes **e**

We'll also pass the points `points`

A function to compute the error

```
error = function(a_0, a_1, points) {
    y_tilde = my_line(points$x, a_0 = a_0, a_1 = a_1)
    e = points$y - y_tilde
    return(sqrt(sum(e^2)))
}
error(a_0 = 2, a_1 = 3, points)

## [1] 7.874008

error(a_0 = 3, a_1 = 0.5, points)

## [1] 1.224745

error(a_0 = 3.1, a_1 = 0.48, points)

## [1] 1.229471
```

We can't be doing this by hand..!

Let's minimise using something cool – *genetic algorithms*

- Genetic algorithms are a stochastic *optimisation* method. There are other types, e.g., gradient descent (deterministic)
- The idea is to use a mechanism mimicking evolution's drive towards higher fitness
- The function value is its fitness
- We try different genes (here, different values of a, b) and evaluate their fitness.. keep the good ones
- We mutate or crossover genes, throw in new ones, etc.
- We keep doing this until we reach a stopping criterion
- We then return the best gene we found

```r
if (!require("GA", quietly = TRUE)) {
  install.packages("GA")
  library(GA)
}
GA = ga(type = "real-valued",
        fitness = function(x) -error(a_0 = x[1], a_1 = x[2], points),
        suggestions = c(a_0 = 2, a_1 = 3),
        lower = c(-10, -10), upper = c(10, 10),
        popSize = 200, maxiter = 150)
```

GA returns an S4 object of class `ga`

```
GA

## An object of class "ga"
##
## Call:
## ga(type = "real-valued", fitness = function(x) -error(a_0 = x[1],     a_1
##
## Available slots:
##  [1] "call"         "type"         "lower"        "upper"        "nBits"
##  [6] "names"        "popSize"      "iter"         "run"          "maxiter
## [11] "suggestions"  "population"   "elitism"      "pcrossover"   "pmutati
## [16] "optim"        "fitness"      "summary"      "bestSol"      "fitness
## [21] "solution"
```

Access slots using @. For instance, the GA finds the coefficients


```
GA@solution
```

```
##             x1        x2
## [1,] 2.999243 0.5004865
```

with fitness value (i.e., $\|\boldsymbol{e}\|$)

```
-GA@fitnessValue
```

```
## [1] 1.224745
```


Fitness value
−5
−10
−15
−20
0
50
100
150
Generation
Best
Mean
Median

- Here, however, we do not have to go brute force: we can reason using mathematics

- We now take a little detour on the math side of things, we will come back to code in a while..

Least squares problems

- Initial considerations
- **The least squares problem**
- Fitting something more complicated

The least squares problem (simplest version)

Definition 1

Given a collection of points $(x_1, y_1), \ldots, (x_n, y_n)$*, find the coefficients* a, b *of the line* $y = a + bx$ *such that*

$$\|\mathbf{e}\| = \sqrt{\varepsilon_1^2 + \cdots + \varepsilon_n^2} = \sqrt{(y_1 - \tilde{y}_1)^2 + \cdots + (y_n - \tilde{y}_n)^2}$$

is minimal, where $\tilde{y}_i = a + bx_i$ *for* $i = 1, \ldots, n$

We just saw how to solve this by brute force using a genetic algorith to minimise $\|e\|$, let us now see how to solve this problem “properly”

For a data point $i = 1, \ldots, n$

$$\varepsilon_i = y_i - \tilde{y}_i = y_i - (a + bx_i)$$

So if we write this for all data points,

$$\begin{aligned} \varepsilon_1 &= y_1 - (a + bx_1) \\ &\vdots \\ \varepsilon_n &= y_n - (a + bx_n) \end{aligned}$$

In matrix form

$$\boldsymbol{e} = \boldsymbol{b} - A\boldsymbol{x}$$

with

$$\boldsymbol{e} = \begin{pmatrix} \varepsilon_1 \\ \vdots \\ \varepsilon_n \end{pmatrix}, A = \begin{pmatrix} 1 & x_1 \\ \vdots & \vdots \\ 1 & x_n \end{pmatrix}, \boldsymbol{x} = \begin{pmatrix} a \\ b \end{pmatrix} \text{ and } \boldsymbol{b} = \begin{pmatrix} y_1 \\ \vdots \\ y_n \end{pmatrix}$$

The least squares problem (reformulated)

Definition 2 (Least squares solutions)

Consider a collection of points $(x_1, y_1), \ldots, (x_n, y_n)$*, a matrix* $A \in \mathcal{M}_{np}$*,* $\boldsymbol{b} \in \mathbb{R}^n$*. A* ***least squares solution*** *of* $A\boldsymbol{x} = \boldsymbol{b}$ *is a vector* $\tilde{\boldsymbol{x}} \in \mathbb{R}^p$ *s.t.*

$$\forall \boldsymbol{x} \in \mathbb{R}^p, \quad \|\boldsymbol{b} - A\tilde{\boldsymbol{x}}\| \leq \|\boldsymbol{b} - A\boldsymbol{x}\|$$

Needed to solve the problem

Definition 3 (Best approximation)

Let V be a vector space, $W \subset V$ and $\mathbf{v} \in V$. The **best approximation** to $\mathbf{v}$ in W is $\tilde{\mathbf{v}} \in W$ s.t.

$$\forall \mathbf{w} \in W, \mathbf{w} \neq \tilde{\mathbf{v}}, \quad \|\mathbf{v} - \tilde{\mathbf{v}}\| < \|\mathbf{v} - \mathbf{w}\|$$

Theorem 4 (Best approximation theorem)

Let V be a vector space with an inner product, $W \subset V$ and $\mathbf{v} \in V$. Then $\text{proj}_W(\mathbf{v})$ is the best approximation to $\mathbf{v}$ in W

Let us find the least squares solution

$\forall \boldsymbol{x} \in \mathbb{R}^p$, $A\boldsymbol{x}$ is a vector in the **column space** of A (the space spanned by the vectors making up the columns of A)

Since $\boldsymbol{x} \in \mathbb{R}^p$, $A\boldsymbol{x} \in \text{col}(A)$

$\implies$ least squares solution of $A\boldsymbol{x} = \boldsymbol{b}$ is a vector $\tilde{\boldsymbol{y}} \in \text{col}(A)$ s.t.

$$\forall \boldsymbol{y} \in \text{col}(A), \quad \|\boldsymbol{b} - \tilde{\boldsymbol{y}}\| \leq \|\boldsymbol{b} - \boldsymbol{y}\|$$

This looks very much like Best approximation and Best approximation theorem

Putting things together

We just stated: The least squares solution of $A\boldsymbol{x} = \boldsymbol{b}$ is a vector $\tilde{\boldsymbol{y}} \in \text{col}(A)$ s.t.

$$\forall \boldsymbol{y} \in \text{col}(A), \quad \|\boldsymbol{b} - \tilde{\boldsymbol{y}}\| \leq \|\boldsymbol{b} - \boldsymbol{y}\|$$

We know (reformulating a tad):

Theorem 5 (Best approximation theorem)

Let V be a vector space with an inner product, $W \subset V$ and $\mathbf{v} \in V$. Then $\text{proj}_W(\mathbf{v}) \in W$ is the best approximation to $\mathbf{v}$ in W, i.e.,

$$\forall \mathbf{w} \in W, \mathbf{w} \neq \text{proj}_W(\mathbf{v}), \quad \|\mathbf{v} - \text{proj}_W(\mathbf{v})\| < \|\mathbf{v} - \mathbf{w}\|$$

$\implies$ $W = \text{col}(A)$, $\boldsymbol{v} = \boldsymbol{b}$ and $\tilde{\boldsymbol{y}} = \text{proj}_{\text{col}(A)}(\mathbf{b})$

So if $\tilde{\boldsymbol{x}}$ is a least squares solution of $A\boldsymbol{x} = \boldsymbol{b}$, then

$$\tilde{\boldsymbol{y}} = A\tilde{\boldsymbol{x}} = \text{proj}_{\text{col}(A)}(\mathbf{b})$$

We have

$$\boldsymbol{b} - A\tilde{\boldsymbol{x}} = \boldsymbol{b} - \text{proj}_{\text{col}(A)}(\mathbf{b}) = \text{perp}_{\text{col}(A)}(\mathbf{b})$$

and it is easy to show that

$$\text{perp}_{\text{col}(A)}(\mathbf{b}) \perp \text{col}(A)$$

So for all columns $\boldsymbol{a}_i$ of A

$$\boldsymbol{a}_i \cdot (\boldsymbol{b} - A\tilde{\boldsymbol{x}}) = 0$$

which we can also write as $\boldsymbol{a}_i^T(\boldsymbol{b} - A\tilde{\boldsymbol{x}}) = 0$

For all columns $\boldsymbol{a}_i$ of A,

$$\boldsymbol{a}_i^T(\boldsymbol{b} - A\tilde{\boldsymbol{x}}) = 0$$

This is equivalent to saying that

$$A^T(\boldsymbol{b} - A\tilde{\boldsymbol{x}}) = \mathbf{0}$$

We have

$$\begin{aligned} A^T(\boldsymbol{b} - A\tilde{\boldsymbol{x}}) = \mathbf{0} &\iff A^T\boldsymbol{b} - A^TA\tilde{\boldsymbol{x}} = \mathbf{0} \\ &\iff A^T\boldsymbol{b} = A^TA\tilde{\boldsymbol{x}} \\ &\iff A^TA\tilde{\boldsymbol{x}} = A^T\boldsymbol{b} \end{aligned}$$

The latter system constitutes the **normal equations** for $\tilde{\boldsymbol{x}}$

Invertibility of A^TA

Theorem 6 (Invertibility of A^TA)

Let $A \in \mathcal{M}_{np}$. Then A has linearly independent columns $\iff$ A^TA invertible

Proof. Since $A^TA \in \mathcal{M}_p$ is square, it is invertible $\iff A^TA\boldsymbol{x} = \boldsymbol{0} \implies \boldsymbol{x} = \boldsymbol{0}$. Also, A has linearly independent columns $\iff A\boldsymbol{x} = \boldsymbol{0} \implies \boldsymbol{x} = \boldsymbol{0}$. Thus, it suffices to show that $A^TA\boldsymbol{x} = \boldsymbol{0} \iff A\boldsymbol{x} = \boldsymbol{0}$.

- ($\Longleftarrow$) If $A\boldsymbol{x} = \boldsymbol{0}$, then multiplying on the left by A^T gives $A^TA\boldsymbol{x} = A^T\boldsymbol{0} = \boldsymbol{0}$.
- ($\Longrightarrow$) Suppose $A^TA\boldsymbol{x} = \boldsymbol{0}$. Multiplying on the left by $\boldsymbol{x}^T$ gives

$$\boldsymbol{x}^TA^TA\boldsymbol{x} = \boldsymbol{x}^T\boldsymbol{0} = 0$$

But $\boldsymbol{x}^TA^TA\boldsymbol{x} = (A\boldsymbol{x})^T(A\boldsymbol{x}) = \|A\boldsymbol{x}\|^2$. Since the norm of a vector is zero if and only if the vector is the zero vector, we have $\|A\boldsymbol{x}\|^2 = 0 \implies A\boldsymbol{x} = \boldsymbol{0}$. □

Least squares theorem

Theorem 7 (Least squares theorem)

$A \in \mathcal{M}_{np}$, $\boldsymbol{b} \in \mathbb{R}^n$. Then

1. *$A\boldsymbol{x} = \boldsymbol{b}$ always has at least one least squares solution $\tilde{\boldsymbol{x}}$*
2. *$\tilde{\boldsymbol{x}}$ least squares solution to $A\boldsymbol{x} = \boldsymbol{b} \iff \tilde{\boldsymbol{x}}$ is a solution to the normal equations $A^T A\tilde{\boldsymbol{x}} = A^T\boldsymbol{b}$*
3. *If $A^T A$ invertible, then the least squares solution is unique and*

$$\tilde{\boldsymbol{x}} = \left(A^T A\right)^{-1} A^T \boldsymbol{b}$$

We have seen 1 and 2, and we just proved 3!

Least squares problems

- Initial considerations
- The least squares problem
- **Fitting something more complicated**

Suppose we want to fit something a bit more complicated..

For instance, instead of the affine function

$$y = a + bx$$

suppose we want to do the quadratic

$$y = a_0 + a_1 x + a_2 x^2$$

or even

$$y = k_0 e^{k_1 x}$$

How do we proceed?

Fitting the quadratic

We have the data points $(x_1, y_1), (x_2, y_2), \ldots, (x_n, y_n)$ and want to fit

$$y = a_0 + a_1 x + a_2 x^2$$

At (x_1, y_1),

$$\tilde{y}_1 = a_0 + a_1 x_1 + a_2 x_1^2$$

$\vdots$

At (x_n, y_n),

$$\tilde{y}_n = a_0 + a_1 x_n + a_2 x_n^2$$

In terms of the error

$$\begin{aligned}\varepsilon_1 &= y_1 - \tilde{y}_1 = y_1 - (a_0 + a_1 x_1 + a_2 x_1^2)\\ &\vdots\\ \varepsilon_n &= y_n - \tilde{y}_n = y_n - (a_0 + a_1 x_n + a_2 x_n^2)\end{aligned}$$

i.e.,

$$\boldsymbol{e} = \boldsymbol{b} - A\boldsymbol{x}$$

where

$$\boldsymbol{e} = \begin{pmatrix}\varepsilon_1\\ \vdots\\ \varepsilon_n\end{pmatrix}, A = \begin{pmatrix}1 & x_1 & x_1^2\\ \vdots & \vdots & \vdots\\ 1 & x_n & x_n^2\end{pmatrix}, \boldsymbol{x} = \begin{pmatrix}a_0\\ a_1\\ a_2\end{pmatrix} \text{ and } \boldsymbol{b} = \begin{pmatrix}y_1\\ \vdots\\ y_n\end{pmatrix}$$

Theorem 7 applies, with here $A \in \mathcal{M}_{n3}$ and $\boldsymbol{b} \in \mathbb{R}^n$

Fitting the exponential

Things are a bit more complicated here

If we proceed as before, we get the system

$$\begin{aligned} y_1 &= k_0 e^{k_1 x_1} \\ &\vdots \\ y_n &= k_0 e^{k_1 x_n} \end{aligned}$$

$e^{k_1 x_i}$ is a nonlinear term, it cannot be put in a matrix

However: take the ln of both sides of the equation

$$\ln(y_i) = \ln(k_0 e^{k_1 x_i}) = \ln(k_0) + \ln(e^{k_1 x_i}) = \ln(k_0) + k_1 x_i$$

If $y_i, k_0 > 0$, then their ln are defined and we're in business..

$$\ln(y_i) = \ln(k_0) + k_1 x_i$$

So the system is

$$\boldsymbol{y} = A\boldsymbol{x} + \boldsymbol{b}$$

with

$$A = \begin{pmatrix} x_1 \\ \vdots \\ x_n \end{pmatrix}, \boldsymbol{x} = \begin{pmatrix} k_1 \end{pmatrix}, \boldsymbol{b} = \begin{pmatrix} \ln(k_0) \end{pmatrix} \text{ and } \boldsymbol{y} = \begin{pmatrix} \ln(y_1) \\ \vdots \\ \ln(y_n) \end{pmatrix}$$

Getting the Canadian census data

Least squares problems

Example – Canadian population data

Example: Canadian census data

Let's consider data from the Canadian census. We first load and visualize it.

```
canada = read.csv("../DATA/Canada_census.csv")
canada = as.data.frame(canada)
plot(canada, main = "Canadian Population over Time",
     xlab = "Year", ylab = "Population")
```

Fitting a linear model

Let's first try to fit a linear model $y = a_0 + a_1 \times \text{year}$.

We construct the matrix A and the vector $\boldsymbol{b}$ (the population data):

```
b = canada$population
A = matrix(ncol = 2, nrow = dim(canada)[1],
           data = c(rep(1, dim(canada)[1]),
                    canada$year))
```

Recall that the normal equations are $A^T A\tilde{\boldsymbol{x}} = A^T \boldsymbol{b}$, where $\tilde{\boldsymbol{x}} = (a_0, a_1)^T$.

Solving the normal equations

We can compute $A^T A$ and solve the normal equations in R:

```
ATA = t(A) %*% A
invATA = solve(ATA)
x_tilde = invATA %*% t(A) %*% b
```

The resulting coefficients $\tilde{\boldsymbol{x}}$ are:


```
##               [,1]
## [1,] -377660589.2
## [2,]     202664.8
```

Visualizing the linear fit

We compute the fitted values and plot them alongside the data:

```
x_axis = seq(canada$year[1], max(canada$year), 1)
y_fitted = x_tilde[1] + x_tilde[2] * x_axis

plot(canada, ylim = c(min(y_fitted), max(canada$population)),
xlab = "Year", ylab = "Population")
lines(x_axis, y_fitted, col = "red", lwd = 2)
```


Population
0.0e+00
1.5e+07
3.0e+07
1850
1900
1950
2000
Year

Fitting a quadratic model

The linear fit misses the curvature in the population growth. Let us try a quadratic model:

$$y = a_0 + a_1 \times \text{year} + a_2 \times \text{year}^2$$

We can set up the matrix A_2:

```
A2 = matrix(ncol = 3, nrow = dim(canada)[1],
            data = c(rep(1, dim(canada)[1]),
                     canada$year,
                     canada$year^2))
```

Numerical issues

Let's try to compute $A_2^T A_2$ and its inverse:

```
ATA2 = t(A2) %*% A2
invATA2 = solve(ATA2)

## Error in `solve.default()`:
## !  system is computationally singular:  reciprocal condition number
= 2.05592e-20
```

The years are around 1900-2000, so year$^2 \approx 4 \times 10^6$. The columns of A_2 have vastly different scales, which makes $A_2^T A_2$ very poorly **conditioned**!

A better conditioned formulation

To avoid numerical issues, we can shift years so they start at 0:

$$\text{shifted_year} = \text{year} - \min(\text{year})$$

We are now fitting $y = a_0 + a_1 \times \text{shifted_year} + a_2 \times \text{shifted_year}^2$.

```
A3 = matrix(ncol = 3, nrow = dim(canada)[1],
            data = c(rep(1, dim(canada)[1]),
                     canada$year - min(canada$year),
                     (canada$year - min(canada$year))^2))
```

This makes the values in the columns of A_3 much closer in magnitude.

Solving the shifted quadratic model

We can now safely compute the normal equations for the shifted model:

```
ATA3 = t(A3) %*% A3
invATA3 = solve(ATA3)
x_tilde = invATA3 %*% t(A3) %*% b
```

The resulting coefficients $\tilde{\boldsymbol{x}}$ are:


```
##             [,1]
## [1,] 2730500.544
## [2,]    3442.989
## [3,]    1208.270
```

Visualizing the quadratic fit

We compute the fitted quadratic curve and plot it:

```
y_fitted_quad = x_tilde[1] +
  x_tilde[2] * (x_axis - min(canada$year)) +
  x_tilde[3] * (x_axis - min(canada$year))^2

plot(canada, ylim = c(min(y_fitted), max(canada$population)),
xlab = "Year", ylab = "Population")
lines(x_axis, y_fitted, col = "red", lwd = 2) # Linear
lines(x_axis, y_fitted_quad, col = "blue", lwd = 2) # Quadratic
```



Population
0.0e+00
2.0e+07
1850
1900
1950
2000
Year

Visualizing the quadratic fit (ggplot2)

```r
df_fit <- data.frame(year = x_axis,
                     linear = y_fitted,
                     quad = y_fitted_quad)

ggplot() +
  geom_point(data = canada, aes(x = year, y = population)) +
  geom_line(data = df_fit, aes(x = year, y = linear), color = "red", linewic
  geom_line(data = df_fit, aes(x = year, y = quad), color = "blue", linewidt
  scale_y_continuous(labels = function(x) paste0(x / 1e6, "M")) +
  labs(x = "Year", y = "Population")
```

`ggplot2` is a graphic package providing a “grammar of graphics”. Nicer output than default `R` plots, but also harder (LLMs help a lot here)


30M
20M
10M
0M
Population
1850
1900
1950
2000
Year

Takeaways

- Matrix formulations of least squares give us a straightforward way to solve polynomial curve fitting problems
- By adding columns corresponding to higher powers (or other transformations), we can easily expand the model
- **Numerical stability is crucial**: always be careful when dealing with polynomials of large numbers. Shifting and scaling your data can prevent matrices from becoming ill-conditioned
- You could shift to the mean or even $[-1, 1]$. Just remember to "unshift" back the years