

# Your Title

Your Name

August 1, 2025

Showcase  
Figure


# Outline

1 Research Context

2 Classical fANOVA

3 Generalized fANOVA

4 Conclusion



References: [1, 2, 5, 7, 6, 4, 3]

# Outline

1 Research Context

2 Classical fANOVA

3 Generalized fANOVA

4 Conclusion

## General Form

$$y(\mathbf{x}) = \sum_{u \subseteq \{1, \dots, N\}} y_u(\mathbf{x}_u) = y_{\emptyset} + y_{\{1\}}(\mathbf{x}_1) + \dots + y_{\{1,2\}}(\mathbf{x}_1, \mathbf{x}_2) + \dots$$

- $y$  : Model output
- $y_u$  : Component functions for subset  $u$
- Assumption:  $X_1, \dots, X_N$  are independent

## Strong Annihilating Conditions

$$\int_{\mathbb{R}} y_u(\mathbf{x}_u) f_{\{i\}}(x_i) d\nu(x_i) = 0 \quad \text{for } i \in u \neq \emptyset.$$

- Ensures unique component functions
- Applies under independent (product-type) input distributions

$$\mathbb{E}[y_u(\mathbf{X}_u)] = 0$$

$$\mathbb{E}[y_u(\mathbf{X}_u)y_v(\mathbf{X}_v)] = 0 \quad (u \neq v)$$

- Zero mean components
- Mutual orthogonality



$$y_{\emptyset} = \int_{\mathbb{R}^N} y(\mathbf{x}) \prod_{i=1}^N f_{\{i\}}(x_i) d\nu(x_i) = \mathbb{E}[y(\mathbf{x})].$$

$$y_u(x_u) = \int y(x) f_{-u}(x_{-u}) dx_{-u} - \sum_{v \subset u} y_v(x_v)$$

- $f_{-u}$  : marginal density of variables not in  $u$
- Components solved sequentially by increasing order





# Example: 2D Function



# Outline

- 1 Research Context
- 2 Classical fANOVA
- 3 Generalized fANOVA**
- 4 Conclusion

# Example with Dependent Inputs ( $\rho = 0.8$ )



## Weak Annihilating Conditions

$$\int_{\mathbb{R}} y_{u,G}(\mathbf{x}_u) f_{\mathbf{x}_u}(\mathbf{x}_u) d\nu(x_i) = 0 \quad \text{for } i \in u \neq \emptyset.$$

- Allows dependent input distributions
- Leads to hierarchical orthogonality

$$\mathbb{E}[y_{u,G}(\mathbf{X}_u)] := \int_{\mathbb{R}^N} y_{u,G}(\mathbf{x}_u) f_{\mathbf{X}}(\mathbf{x}) d\nu(\mathbf{x}) = 0.$$

$$\mathbb{E}[y_{u,G}(\mathbf{X}_u) y_{v,G}(\mathbf{X}_v)] := \int_{\mathbb{R}^N} y_{u,G}(\mathbf{x}_u) y_{v,G}(\mathbf{x}_v) f_{\mathbf{X}}(\mathbf{x}) d\nu(\mathbf{x}) = 0.$$

- Zero mean components remain
- Orthogonality is weaker: hierarchical

$$y_{\emptyset,G} = \int_{\mathbb{R}^N} y(\mathbf{x}) f_{\mathbf{x}}(\mathbf{x}) d\nu(\mathbf{x}) \quad (1)$$

$$\begin{aligned} y_{u,G}(\mathbf{x}_u) &= \int_{\mathbb{R}^{N-|u|}} y(\mathbf{x}_u, \mathbf{x}_{-u}) f_{-u}(\mathbf{x}_{-u}) d\nu(\mathbf{x}_{-u}) - \sum_{v \subsetneq u} y_{v,G}(\mathbf{x}_v) \\ &\quad - \sum_{\substack{\emptyset \neq v \subseteq \{1, \dots, N\} \\ v \cap u \neq \emptyset, v \not\subseteq u}} \int_{\mathbb{R}^{|v \cap u|}} y_{v,G}(\mathbf{x}_{v \cap u}, \mathbf{x}_{v \cap u}) f_{v \cap u}(\mathbf{x}_{v \cap u}) d\nu(\mathbf{x}_{v \cap u}). \end{aligned} \quad (2)$$

- All components solved simultaneously
- Depends on marginal densities and coupling terms

- Coupled system  $\rightarrow$  difficult to obtain analytical solutions
- Use alternative method via Fourier Polynomial ([5])
- Building blocks: mutually orthogonal, zero-mean basis functions  $\psi_{i,j}$ , coefficients  $c_{i,j}$

$$\begin{aligned}y(x_1, x_2) &= a_0 + a_1x_1 + a_2x_2 + a_{11}x_1^2 + a_{22}x_2^2 + a_{12}x_1x_2 \\&= c_0 + c_{1,1} \psi_{1,1}(x_1) + c_{2,1} \psi_{2,1}(x_2) \\&\quad + c_{1,2} \psi_{1,2}(x_1) + c_{2,2} \psi_{2,2}(x_2) + c_{12,11} \psi_{12,11}(x_1, x_2) \\&= \underbrace{c_0}_{y_0} + \underbrace{(c_{1,1} \psi_{1,1}(x_1) + c_{1,2} \psi_{1,2}(x_1))}_{y_1(x_1)} \\&\quad + \underbrace{(c_{2,1} \psi_{2,1}(x_2) + c_{2,2} \psi_{2,2}(x_2))}_{y_2(x_2)} \\&\quad + \underbrace{c_{12,11} \psi_{12,11}(x_1, x_2)}_{y_{12}(x_1, x_2)}.\end{aligned}$$



$$\psi_{\emptyset}(x_1, x_2) = 1,$$

$$\psi_{1,1}(x_1) = x_1,$$

$$\psi_{2,1}(x_2) = x_2,$$

$$\psi_{1,2}(x_1) = x_1^2 - 1,$$

$$\psi_{2,2}(x_2) = x_2^2 - 1,$$

$$\psi_{12,11}(x_1, x_2) = \frac{\rho(x_1^2 + x_2^2)}{1 + \rho^2} - x_1x_2 + \frac{\rho(\rho^2 - 1)}{1 + \rho^2},$$

$$\{y_{u,G}(\mathbf{x}_u) \mid u \subseteq d\} = \arg \min_{\{g_u \in \mathcal{L}^2(\mathbb{R}^{|u|})\}} \int_{\mathbb{R}^N} \left( \sum_{u \subseteq d} g_u(\mathbf{x}_u) - y(\mathbf{x}) \right)^2 f_{\mathbf{x}}(\mathbf{x}) d\nu(\mathbf{x}),$$

subject to hierarchical orthogonality conditions:

$$\forall v \subseteq u, \forall g_v : \int_{\mathbb{R}^N} y_u(\mathbf{x}_u) g_v(\mathbf{x}_v) f_{\mathbf{x}}(\mathbf{x}) d\nu(\mathbf{x}) = 0.$$

$$\mu := \mathbb{E}[y(\mathbf{X})] = y_{\emptyset, G}.$$

$$\begin{aligned}\sigma^2 &:= \mathbb{E} \left[ (y(\mathbf{X}) - \mu_G)^2 \right] \\ &= \mathbb{E} \left[ \left( y_{\emptyset, G} + \sum_u y_{u, G}(\mathbf{x}_u) - y_{\emptyset, G} \right)^2 \right] \\ &= \mathbb{E} \left[ \left( \sum_u y_{u, G}(\mathbf{x}_u) \right)^2 \right] \\ &= \sum_u \mathbb{E} [y_{u, G}^2(\mathbf{x}_u)] + \sum_{u \not\subseteq v, v \not\subseteq u} \mathbb{E} [y_{u, G}(\mathbf{x}_u) y_{v, G}(\mathbf{x}_v)],\end{aligned}$$

# Outline

- 1 Research Context
- 2 Classical fANOVA
- 3 Generalized fANOVA
- 4 Conclusion**



Giles Hooker.

Discovering additive structure in black box functions.

*In Proceedings of the tenth ACM SIGKDD international conference on Knowledge discovery and data mining*, pages 575–580, Seattle WA USA, August 2004. ACM.



Giles Hooker.

Generalized Functional ANOVA Diagnostics for High-Dimensional Functions of Dependent Variables.

*Journal of Computational and Graphical Statistics*, 16(3):709–732, September 2007.



Gunnar König, Eric Günther, and Ulrike von Luxburg.

Disentangling Interactions and Dependencies in Feature Attribution, October 2024.  
[arXiv:2410.23772](https://arxiv.org/abs/2410.23772).



Benjamin Lengerich, Sarah Tan, Chun-Hao Chang, Giles Hooker, and Rich Caruana.  
Purifying Interaction Effects with the Functional ANOVA: An Efficient Algorithm for  
Recovering Identifiable Additive Models.

*In Proceedings of the Twenty Third International Conference on Artificial Intelligence and  
Statistics*, pages 2402–2412. PMLR, June 2020.

ISSN: 2640-3498.



Sharif Rahman.

A Generalized ANOVA Dimensional Decomposition for Dependent Probability Measures.

*SIAM/ASA Journal on Uncertainty Quantification*, 2(1):670–697, 2014.



I. M. Sobol.

Sensitivity Estimates for Nonlinear Mathematical Models.

*Mathematical Modelling and Computational Experiments*, 1:407–414, 1993.



Charles J. Stone.

The Use of Polynomial Splines and Their Tensor Products in Multivariate Function Estimation.

*The Annals of Statistics*, 22(1):118–171, 1994.

Publisher: Institute of Mathematical Statistics.