

Your Title

Your Name

August 1, 2025

Showcase
Figure

Outline


1 Research Context

2 Classical fANOVA

3 Generalized fANOVA

4 Conclusion

5 Extra Slides



References: [1, 2, 5, 7, 6, 4, 3]

Outline

1 Research Context

2 Classical fANOVA

3 Generalized fANOVA

4 Conclusion

5 Extra Slides

- measure space, probability measure, random vector, subvector, complementary vector, pdf
- measure space of square integrable functions
- inner product
- norm

General Form

$$y(\mathbf{x}) = \sum_{u \subseteq \{1, \dots, N\}} y_u(\mathbf{x}_u) = y_{\emptyset} + y_{\{1\}}(\mathbf{x}_1) + \dots + y_{\{1,2\}}(\mathbf{x}_1, \mathbf{x}_2) + \dots$$

- y : Model
- y_u : Component functions for subset u
- Assumption: X_1, \dots, X_N are independent

Strong Annihilating Conditions

$$\int_{\mathbb{R}} y_u(\mathbf{x}_u) f_{\{i\}}(x_i) d\nu(x_i) = 0 \quad \text{for } i \in u \neq \emptyset.$$

- Ensures unique component functions
- Applies under independent (product-type) input distributions

$$\mathbb{E}[y_u(\mathbf{X}_u)] = 0$$

$$\mathbb{E}[y_u(\mathbf{X}_u)y_v(\mathbf{X}_v)] = 0 \quad (u \neq v)$$

- Zero mean components
- Mutual orthogonality

$$y_{\emptyset} = \int_{\mathbb{R}^N} y(\mathbf{x}) \prod_{i=1}^N f_{\{i\}}(x_i) d\nu(x_i) = \mathbb{E}[y(\mathbf{x})].$$

$$y_u(x_u) = \int y(x) f_{-u}(x_{-u}) dx_{-u} - \sum_{v \subsetneq u} y_v(x_v)$$



- f_{-u} : marginal density of variables not in u
- Components solved sequentially by increasing order

Example: Two degree Polynomial

$$y(x_1, x_2) = 2x_1 + x_2^2 + x_1x_2 \quad (1)$$

- Coefficients: $a_1 = 20, a_2 = 0, a_{11} = 0, a_{22} = 10, a_{12} = 10$
- Independent variables: $\rho = 0$

Example: 2D Function



Hoeffding Decomposition

$$y(\mathbf{x}) = \sum_{A \subseteq D} y_A(\mathbf{x}_A), \quad D := \{1, \dots, N\}, \quad (2)$$

where, for each $A \subseteq D$, the component function y_A is defined by:

$$y_A(\mathbf{x}_A) = \sum_{B \subseteq A} (-1)^{|A|-|B|} \mathbb{E}[y(\mathbf{x}) \mid \mathbf{x}_B], \quad (3)$$

where y_u are orthogonal components.

- Classical fANOVA and Hoeffding decomposition yield same components under zero-centered inputs
- Both assume independence of input variables

$$\Pi_{\mathcal{G}} y = \arg \min_{g \in \mathcal{G}} \|y - g\|^2 = \arg \min_{g \in \mathcal{G}} \mathbb{E}[(y(\mathbf{X}) - g(\mathbf{X}))^2]. \quad (4)$$

- \mathcal{G} : linear subspace of \mathcal{L}^2 we project onto
- g all functions in the subspace

$$\begin{aligned}\Pi_{\mathcal{G}_0} y &= \arg \min_{g_0 \in \mathcal{G}_0} \|y - g_0\|^2 \\ &= \arg \min_{a \in \mathbb{R}} \mathbb{E}[(y(\mathbf{X}) - a)^2] \\ &= \mathbb{E}[y(\mathbf{X})] = y_\emptyset.\end{aligned}$$

$$\begin{aligned}(\Pi_{\mathcal{G}_u} y)(\cdot) - \sum_{v \subsetneq u} y_v(\cdot) &= \arg \min_{g_u \in \mathcal{G}_u} \|y - g_u\|^2 - \sum_{v \subsetneq u} y_v(\cdot) \\ &= \arg \min_{g_u \in \mathcal{G}_u} \mathbb{E}[(y(\mathbf{X}) - g_u(\cdot))^2] - \sum_{v \subsetneq u} y_v(\cdot) \\ &= \mathbb{E}[y(\mathbf{X}) \mid X_u = \cdot] - \sum_{v \subsetneq u} y_v(x) = y_u(\cdot).\end{aligned}$$

Outline

1 Research Context



2 Classical fANOVA

3 Generalized fANOVA

4 Conclusion

5 Extra Slides

Example with Dependent Inputs ($\rho = 0.8$)



Weak Annihilating Conditions

$$\int_{\mathbb{R}} y_{u,G}(\mathbf{x}_u) f_{\mathbf{x}_u}(\mathbf{x}_u) d\nu(x_i) = 0 \quad \text{for } i \in u \neq \emptyset.$$

- Allows dependent input distributions
- Leads to hierarchical orthogonality

$$\mathbb{E}[y_{u,G}(\mathbf{X}_u)] := \int_{\mathbb{R}^N} y_{u,G}(\mathbf{x}_u) f_{\mathbf{X}}(\mathbf{x}) d\nu(\mathbf{x}) = 0.$$

$$\mathbb{E}[y_{u,G}(\mathbf{X}_u) y_{v,G}(\mathbf{X}_v)] := \int_{\mathbb{R}^N} y_{u,G}(\mathbf{x}_u) y_{v,G}(\mathbf{x}_v) f_{\mathbf{X}}(\mathbf{x}) d\nu(\mathbf{x}) = 0.$$

- Zero mean components remain
- Orthogonality is weaker: hierarchical

$$y_{\emptyset,G} = \int_{\mathbb{R}^N} y(\mathbf{x}) f_{\mathbf{x}}(\mathbf{x}) d\nu(\mathbf{x}) \quad (5)$$

$$\begin{aligned} y_{u,G}(\mathbf{X}_u) &= \int_{\mathbb{R}^{N-|u|}} y(\mathbf{X}_u, \mathbf{x}_{-u}) f_{-u}(\mathbf{x}_{-u}) d\nu(\mathbf{x}_{-u}) - \sum_{v \subsetneq u} y_{v,G}(\mathbf{X}_v) \\ &\quad - \sum_{\substack{\emptyset \neq v \subseteq \{1, \dots, N\} \\ v \cap u \neq \emptyset, v \not\subseteq u}} \int_{\mathbb{R}^{|v \cap u|}} y_{v,G}(\mathbf{X}_{v \cap u}, \mathbf{x}_{v \cap u}) f_{v \cap u}(\mathbf{x}_{v \cap u}) d\nu(\mathbf{x}_{v \cap u}). \end{aligned} \quad (6)$$

- All components solved simultaneously
- Depends on marginal densities and coupling terms

- Coupled system \rightarrow difficult to obtain analytical solutions
- Use alternative method via Fourier Polynomial ([5])
- Building blocks: mutually orthogonal, zero-mean basis functions $\psi_{i,j}$, coefficients $c_{i,j}$

$$\begin{aligned}y(x_1, x_2) &= a_0 + a_1x_1 + a_2x_2 + a_{11}x_1^2 + a_{22}x_2^2 + a_{12}x_1x_2 \\&= c_0 + c_{1,1} \psi_{1,1}(x_1) + c_{2,1} \psi_{2,1}(x_2) \\&\quad + c_{1,2} \psi_{1,2}(x_1) + c_{2,2} \psi_{2,2}(x_2) + c_{12,11} \psi_{12,11}(x_1, x_2) \\&= \underbrace{c_0}_{y_0} + \underbrace{(c_{1,1} \psi_{1,1}(x_1) + c_{1,2} \psi_{1,2}(x_1))}_{y_1(x_1)} \\&\quad + \underbrace{(c_{2,1} \psi_{2,1}(x_2) + c_{2,2} \psi_{2,2}(x_2))}_{y_2(x_2)} \\&\quad + \underbrace{c_{12,11} \psi_{12,11}(x_1, x_2)}_{y_{12}(x_1, x_2)}.\end{aligned}$$

$$\psi_{\emptyset}(x_1, x_2) = 1,$$

$$\psi_{1,1}(x_1) = x_1,$$

$$\psi_{2,1}(x_2) = x_2,$$

$$\psi_{1,2}(x_1) = x_1^2 - 1,$$

$$\psi_{2,2}(x_2) = x_2^2 - 1,$$

$$\psi_{12,11}(x_1, x_2) = \frac{\rho(x_1^2 + x_2^2)}{1 + \rho^2} - x_1x_2 + \frac{\rho(\rho^2 - 1)}{1 + \rho^2},$$

$$\{y_{u,G}(\mathbf{x}_u) \mid u \subseteq d\} = \arg \min_{\{g_u \in \mathcal{L}^2(\mathbb{R}^{|u|})\}} \int_{\mathbb{R}^N} \left(\sum_{u \subseteq d} g_u(\mathbf{x}_u) - y(\mathbf{x}) \right)^2 f_{\mathbf{x}}(\mathbf{x}) d\nu(\mathbf{x}),$$

subject to hierarchical orthogonality conditions:

$$\forall v \subseteq u, \forall g_v : \int_{\mathbb{R}^N} y_u(\mathbf{x}_u) g_v(\mathbf{x}_v) f_{\mathbf{x}}(\mathbf{x}) d\nu(\mathbf{x}) = 0.$$

$$\mu := \mathbb{E}[y(\mathbf{X})] = y_{\emptyset, G}.$$

$$\begin{aligned}\sigma^2 &:= \mathbb{E} \left[(y(\mathbf{X}) - \mu_G)^2 \right] \\ &= \mathbb{E} \left[\left(y_{\emptyset, G} + \sum_u y_{u, G}(\mathbf{x}_u) - y_{\emptyset, G} \right)^2 \right] \\ &= \mathbb{E} \left[\left(\sum_u y_{u, G}(\mathbf{x}_u) \right)^2 \right] \\ &= \sum_u \mathbb{E} [y_{u, G}^2(\mathbf{x}_u)] + \sum_{u \not\subseteq v, v \not\subseteq u} \mathbb{E} [y_{u, G}(\mathbf{x}_u) y_{v, G}(\mathbf{x}_v)],\end{aligned}$$

Outline

- 1 Research Context
- 2 Classical fANOVA
- 3 Generalized fANOVA
- 4 Conclusion**
- 5 Extra Slides

- Estimation schemes and software implementation
- Extension of Fourier polynomial expansion to other distributions

Outline

1 Research Context

2 Classical fANOVA

3 Generalized fANOVA

4 Conclusion

5 Extra Slides

- Zero mean property: factorized density, Fubini's Theorem, strong annihilating conditions
- Mutual orthogonality: factorized density, Fubini's Theorem, strong annihilating conditions

- Zero mean property: separating x into subvectors, marginal density, Fubini's Theorem, weak annihilating conditions
- Hierarchical orthogonality: set the scene, u is a proper subset of v $u \subsetneq v$, so there is an index in u which is not in v ; divide x_u into subvectors, marginal density, Fubini and weak annihilating conditions
- Weak annihilating becomes strong under independence: assume the weak ones, product density, factor out
- Three integration cases: distinguish between different relationships u and v , depending on the relationship the integral w.r.t. to marginal density simplifies
- Generalized fANOVA components by Rahman: first build constant term; for nonconstant terms use integration cases
- Integration constraint Hooker: show that hierarchical orthogonality is fulfilled if the conditions hold, show that it is not fulfilled if they do not hold; but why exactly these conditions a bit unclear
- Take a look at Sobol's proof again



Giles Hooker.

Discovering additive structure in black box functions.

In Proceedings of the tenth ACM SIGKDD international conference on Knowledge discovery and data mining, pages 575–580, Seattle WA USA, August 2004. ACM.



Giles Hooker.

Generalized Functional ANOVA Diagnostics for High-Dimensional Functions of Dependent Variables.

Journal of Computational and Graphical Statistics, 16(3):709–732, September 2007.



Gunnar König, Eric Günther, and Ulrike von Luxburg.

Disentangling Interactions and Dependencies in Feature Attribution, October 2024.
[arXiv:2410.23772](https://arxiv.org/abs/2410.23772).



Benjamin Lengerich, Sarah Tan, Chun-Hao Chang, Giles Hooker, and Rich Caruana.
Purifying Interaction Effects with the Functional ANOVA: An Efficient Algorithm for
Recovering Identifiable Additive Models.

*In Proceedings of the Twenty Third International Conference on Artificial Intelligence and
Statistics*, pages 2402–2412. PMLR, June 2020.

ISSN: 2640-3498.



Sharif Rahman.

A Generalized ANOVA Dimensional Decomposition for Dependent Probability Measures.

SIAM/ASA Journal on Uncertainty Quantification, 2(1):670–697, 2014.



I. M. Sobol.

Sensitivity Estimates for Nonlinear Mathematical Models.

Mathematical Modelling and Computational Experiments, 1:407–414, 1993.



Charles J. Stone.

The Use of Polynomial Splines and Their Tensor Products in Multivariate Function Estimation.

The Annals of Statistics, 22(1):118–171, 1994.

Publisher: Institute of Mathematical Statistics.