


Functional ANOVA Decomposition

Juliet Fleischer
August 12, 2025



1 Research Context

2 Classical fANOVA


3 Generalized fANOVA


4 Conclusion


Foundations


Foundations → (Generalized) fANOVA














Outline

1 Research Context

2 Classical fANOVA

3 Generalized fANOVA

4 Conclusion

General Form

$$y(\mathbf{x}) = \sum_{u \subseteq \{1, \dots, N\}} y_u(\mathbf{x}_u)$$

General Form

$$y(\mathbf{X}) = \sum_{u \subseteq \{1, \dots, N\}} y_u(\mathbf{X}_u)$$

- y : Model
- y_u : Component functions for subvector \mathbf{X}_u
- $\mathbf{X} = (X_1, \dots, X_N)$: input variables (assumed to be independent in classical fANOVA)

General Form

$$\begin{aligned} y(\mathbf{X}) &= \sum_{u \subseteq \{1, \dots, N\}} y_u(\mathbf{X}_u) \\ &= y_{\emptyset} + (y_{\{1\}}(X_1) + \dots + y_{\{N\}}(X_N)) \end{aligned}$$

- y : Model
- y_u : Component functions for subvector \mathbf{X}_u
- $\mathbf{X} = (X_1, \dots, X_N)$: input variables (assumed to be independent in classical fANOVA)

General Form

$$\begin{aligned} y(\mathbf{X}) &= \sum_{u \subseteq \{1, \dots, N\}} y_u(\mathbf{X}_u) \\ &= y_{\emptyset} + (y_{\{1\}}(X_1) + \dots + y_{\{N\}}(X_N)) \\ &\quad + (y_{\{1,2\}}(X_1, X_2) + y_{\{1,3\}}(X_1, X_3) + \dots) \end{aligned}$$

- y : Model
- y_u : Component functions for subvector \mathbf{X}_u
- $\mathbf{X} = (X_1, \dots, X_N)$: input variables (assumed to be independent in classical fANOVA)

General Form

$$\begin{aligned} y(\mathbf{X}) &= \sum_{u \subseteq \{1, \dots, N\}} y_u(\mathbf{X}_u) \\ &= y_{\emptyset} + (y_{\{1\}}(X_1) + \dots + y_{\{N\}}(X_N)) \\ &\quad + (y_{\{1,2\}}(X_1, X_2) + y_{\{1,3\}}(X_1, X_3) + \dots) \\ &\quad + (y_{\{1,2,3\}}(X_1, X_2, X_3) + \dots) + \dots + y_{\{1, \dots, N\}}(X_1, \dots, X_N) \end{aligned}$$

- y : Model
- y_u : Component functions for subvector \mathbf{X}_u
- $\mathbf{X} = (X_1, \dots, X_N)$: input variables (assumed to be independent in classical fANOVA)

Strong Annihilating Conditions

$$\int_{\mathbb{R}} y_u(\mathbf{x}_u) f_{\{i\}}(x_i) d\nu(x_i) = 0 \quad \text{for } i \in u \neq \emptyset$$

Strong Annihilating Conditions

$$\int_{\mathbb{R}} y_u(\mathbf{x}_u) f_{\{i\}}(x_i) d\nu(x_i) = 0 \quad \text{for } i \in u \neq \emptyset$$

- $f_{\{i\}}$: marginal density of variable X_i
- ν : reference measure on \mathbb{R} ; $d\nu(x_i)$ indicates integration with respect to ν in the variable x_i

Strong Annihilating Conditions

$$\int_{\mathbb{R}} y_u(\mathbf{x}_u) f_{\{i\}}(x_i) d\nu(x_i) = 0 \quad \text{for } i \in u \neq \emptyset$$

- $f_{\{i\}}$: marginal density of variable X_i
- ν : reference measure on \mathbb{R} ; $d\nu(x_i)$ indicates integration with respect to ν in the variable x_i

It follows:

$$\mathbb{E}[y_u(\mathbf{x}_u)] = 0$$

Strong Annihilating Conditions

$$\int_{\mathbb{R}} y_u(\mathbf{x}_u) f_{\{i\}}(x_i) d\nu(x_i) = 0 \quad \text{for } i \in u \neq \emptyset$$

- $f_{\{i\}}$: marginal density of variable X_i
- ν : reference measure on \mathbb{R} ; $d\nu(x_i)$ indicates integration with respect to ν in the variable x_i

It follows:

$$\mathbb{E}[y_u(\mathbf{x}_u)] = 0$$

$$\mathbb{E}[y_u(\mathbf{x}_u) y_v(\mathbf{x}_v)] = 0 \quad (u \neq v)$$

$$y_{\emptyset} = \int_{\mathbb{R}^N} y(\mathbf{x}) \prod_{i=1}^N f_{\{i\}}(x_i) d\nu(x_i) = \mathbb{E}[y(\mathbf{x})]$$

$$y_{\emptyset} = \int_{\mathbb{R}^N} y(\mathbf{x}) \prod_{i=1}^N f_{\{i\}}(x_i) d\nu(x_i) = \mathbb{E}[y(\mathbf{x})]$$

- $f_{\mathbf{x}}(\mathbf{x}) d\nu(\mathbf{x}) = \prod_{i=1}^N f_{\{i\}}(x_i) d\nu(x_i)$
- $f_{\mathbf{x}}(\mathbf{x}) d\nu(\mathbf{x}_{-u}) = \prod_{i=1, i \notin u}^N f_{\{i\}}(x_i) d\nu(x_i)$

$$y_{\emptyset} = \int_{\mathbb{R}^N} y(\mathbf{x}) \prod_{i=1}^N f_{\{i\}}(x_i) d\nu(x_i) = \mathbb{E}[y(\mathbf{x})]$$

$$y_u(\mathbf{x}_u) = \int_{\mathbb{R}^{N-|u|}} y(\mathbf{x}_u, \mathbf{x}_{-u}) \prod_{i=1, i \notin u}^N f_{\{i\}}(x_i) d\nu(x_i) - \sum_{v \subsetneq u} y_v(\mathbf{x}_v)$$

- $f_{\mathbf{x}}(\mathbf{x}) d\nu(\mathbf{x}) = \prod_{i=1}^N f_{\{i\}}(x_i) d\nu(x_i)$
- $f_{\mathbf{x}}(\mathbf{x}) d\nu(\mathbf{x}_{-u}) = \prod_{i=1, i \notin u}^N f_{\{i\}}(x_i) d\nu(x_i)$

$$y_{\emptyset} = \int_{\mathbb{R}^N} y(\mathbf{x}) \prod_{i=1}^N f_{\{i\}}(x_i) d\nu(x_i) = \mathbb{E}[y(\mathbf{x})]$$

$$\begin{aligned} y_u(\mathbf{x}_u) &= \int_{\mathbb{R}^{N-|u|}} y(\mathbf{x}_u, \mathbf{x}_{-u}) \prod_{i=1, i \notin u}^N f_{\{i\}}(x_i) d\nu(x_i) - \sum_{v \subsetneq u} y_v(\mathbf{x}_v) \\ &= \mathbb{E}[y(\mathbf{x}_u, \mathbf{x}_{-u}) \mid \mathbf{x}_u = \mathbf{x}_u] - \sum_{v \subsetneq u} y_v(\mathbf{x}_v) \end{aligned}$$

- $f_{\mathbf{x}}(\mathbf{x}) d\nu(\mathbf{x}) = \prod_{i=1}^N f_{\{i\}}(x_i) d\nu(x_i)$
- $f_{\mathbf{x}}(\mathbf{x}) d\nu(\mathbf{x}_{-u}) = \prod_{i=1, i \notin u}^N f_{\{i\}}(x_i) d\nu(x_i)$

Example of Recursive Form

- $N = 3, u = \emptyset$

Example of Recursive Form

- $N = 3, u = \emptyset$

$$y_{\emptyset} = \int_{\mathbb{R}^3} y(\mathbf{x}) \prod_{i=1}^3 f_{\{i\}}(x_i) d\nu(x_i) = \mathbb{E}[y(\mathbf{x})]$$

Example of Recursive Form

- $N = 3, u = \emptyset$

$$y_{\emptyset} = \int_{\mathbb{R}^3} y(\mathbf{x}) \prod_{i=1}^3 f_{\{i\}}(x_i) d\nu(x_i) = \mathbb{E}[y(\mathbf{x})]$$

- $u = \{1\} \rightarrow v \in \{\emptyset\}$

- $N = 3, u = \emptyset$

$$y_{\emptyset} = \int_{\mathbb{R}^3} y(\mathbf{x}) \prod_{i=1}^3 f_{\{i\}}(x_i) d\nu(x_i) = \mathbb{E}[y(\mathbf{x})]$$

- $u = \{1\} \rightarrow v \in \{\emptyset\}$

$$y_{\{1\}}(x_1) = \int_{\mathbb{R}^2} y(x_1, x_2, x_3) \prod_{i=2}^3 f_{\{i\}}(x_i) d\nu(x_i) - y_{\emptyset} = \mathbb{E}[y(X_1, X_2, X_3) | X_1 = x_1] - y_{\emptyset}$$

- $N = 3, u = \emptyset$

$$y_{\emptyset} = \int_{\mathbb{R}^3} y(\mathbf{x}) \prod_{i=1}^3 f_{\{i\}}(x_i) d\nu(x_i) = \mathbb{E}[y(\mathbf{x})]$$

- $u = \{1\} \rightarrow v \in \{\emptyset\}$

$$y_{\{1\}}(x_1) = \int_{\mathbb{R}^2} y(x_1, x_2, x_3) \prod_{i=2}^3 f_{\{i\}}(x_i) d\nu(x_i) - y_{\emptyset} = \mathbb{E}[y(X_1, X_2, X_3) | X_1 = x_1] - y_{\emptyset}$$

- $u = \{2\} \rightarrow v \in \{\emptyset\}$

- $N = 3, u = \emptyset$

$$y_{\emptyset} = \int_{\mathbb{R}^3} y(\mathbf{x}) \prod_{i=1}^3 f_{\{i\}}(x_i) d\nu(x_i) = \mathbb{E}[y(\mathbf{x})]$$

- $u = \{1\} \rightarrow v \in \{\emptyset\}$

$$y_{\{1\}}(x_1) = \int_{\mathbb{R}^2} y(x_1, x_2, x_3) \prod_{i=2}^3 f_{\{i\}}(x_i) d\nu(x_i) - y_{\emptyset} = \mathbb{E}[y(X_1, X_2, X_3) | X_1 = x_1] - y_{\emptyset}$$

- $u = \{2\} \rightarrow v \in \{\emptyset\}$

$$y_{\{2\}}(x_2) = \int_{\mathbb{R}^2} y(x_1, x_2, x_3) \prod_{i=1, i \neq 2}^3 f_{\{i\}}(x_i) d\nu(x_i) - y_{\emptyset} = \mathbb{E}[y(X_1, X_2, X_3) | X_2 = x_2] - y_{\emptyset}$$

- $u = \{1, 2\} \rightarrow v \in \{\emptyset, \{1\}, \{2\}\}$

- $u = \{1, 2\} \rightarrow v \in \{\emptyset, \{1\}, \{2\}\}$

$$\begin{aligned} y_{\{1,2\}}(x_1, x_2) &= \int_{\mathbb{R}} y(x_1, x_2, x_3) f_{\{3\}}(x_3) d\nu(x_3) - y_{\{1\}}(x_1) - y_{\{2\}}(x_2) - y_{\emptyset} \\ &= \mathbb{E}[y(X_1, X_2, X_3) | X_1 = x_1, X_2 = x_2] - y_{\{1\}}(x_1) - y_{\{2\}}(x_2) - y_{\emptyset} \end{aligned}$$

- $u = \{1, 2\} \rightarrow v \in \{\emptyset, \{1\}, \{2\}\}$

$$\begin{aligned} y_{\{1,2\}}(x_1, x_2) &= \int_{\mathbb{R}} y(x_1, x_2, x_3) f_{\{3\}}(x_3) d\nu(x_3) - y_{\{1\}}(x_1) - y_{\{2\}}(x_2) - y_{\emptyset} \\ &= \mathbb{E}[y(X_1, X_2, X_3) | X_1 = x_1, X_2 = x_2] - y_{\{1\}}(x_1) - y_{\{2\}}(x_2) - y_{\emptyset} \end{aligned}$$

- $u = \{1, 2, 3\} \rightarrow v \in \{\emptyset, \{1\}, \{2\}, \{3\}, \{1, 2\}, \{1, 3\}, \{2, 3\}\}$

- $u = \{1, 2\} \rightarrow v \in \{\emptyset, \{1\}, \{2\}\}$

$$\begin{aligned} y_{\{1,2\}}(x_1, x_2) &= \int_{\mathbb{R}} y(x_1, x_2, x_3) f_{\{3\}}(x_3) d\nu(x_3) - y_{\{1\}}(x_1) - y_{\{2\}}(x_2) - y_{\emptyset} \\ &= \mathbb{E}[y(X_1, X_2, X_3) | X_1 = x_1, X_2 = x_2] - y_{\{1\}}(x_1) - y_{\{2\}}(x_2) - y_{\emptyset} \end{aligned}$$

- $u = \{1, 2, 3\} \rightarrow v \in \{\emptyset, \{1\}, \{2\}, \{3\}, \{1, 2\}, \{1, 3\}, \{2, 3\}\}$

$$\begin{aligned} y_{\{1,2,3\}}(x_1, x_2, x_3) &= y(x_1, x_2, x_3) - y_{\{1\}}(x_1) - y_{\{2\}}(x_2) - y_{\{3\}}(x_3) - y_{\emptyset} \\ &\quad - y_{\{1,2\}}(x_1, x_2) - y_{\{1,3\}}(x_1, x_3) - y_{\{2,3\}}(x_2, x_3) \end{aligned}$$

- $u = \{1, 2\} \rightarrow v \in \{\emptyset, \{1\}, \{2\}\}$

$$\begin{aligned} y_{\{1,2\}}(x_1, x_2) &= \int_{\mathbb{R}} y(x_1, x_2, x_3) f_{\{3\}}(x_3) d\nu(x_3) - y_{\{1\}}(x_1) - y_{\{2\}}(x_2) - y_{\emptyset} \\ &= \mathbb{E}[y(X_1, X_2, X_3) | X_1 = x_1, X_2 = x_2] - y_{\{1\}}(x_1) - y_{\{2\}}(x_2) - y_{\emptyset} \end{aligned}$$

- $u = \{1, 2, 3\} \rightarrow v \in \{\emptyset, \{1\}, \{2\}, \{3\}, \{1, 2\}, \{1, 3\}, \{2, 3\}\}$

$$\begin{aligned} y_{\{1,2,3\}}(x_1, x_2, x_3) &= y(x_1, x_2, x_3) - y_{\{1\}}(x_1) - y_{\{2\}}(x_2) - y_{\{3\}}(x_3) - y_{\emptyset} \\ &\quad - y_{\{1,2\}}(x_1, x_2) - y_{\{1,3\}}(x_1, x_3) - y_{\{2,3\}}(x_2, x_3) \end{aligned}$$

Remark: fANOVA components can be seen from the lens of orthogonal projections.

Example with Independent MVN Input

$$y(x_1, x_2) = 2x_1 + x_2^2 + x_1x_2$$

$$y(x_1, x_2) = 2x_1 + x_2^2 + x_1x_2$$

$$(X_1, X_2)^T \sim \mathcal{N}\left(\begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}\right)$$

$$y(x_1, x_2) = 2x_1 + x_2^2 + x_1x_2$$

$$(X_1, X_2)^T \sim \mathcal{N}\left(\begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}\right)$$

$$X_1 \mid X_2 = x_2 \sim \mathcal{N}(0, 1), \quad X_2 \mid X_1 = x_1 \sim \mathcal{N}(0, 1)$$

$$y(x_1, x_2) = 2x_1 + x_2^2 + x_1x_2$$

$$(X_1, X_2)^T \sim \mathcal{N}\left(\begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}\right)$$

$$X_1 \mid X_2 = x_2 \sim \mathcal{N}(0, 1), \quad X_2 \mid X_1 = x_1 \sim \mathcal{N}(0, 1)$$

Components:

$$y(x_1, x_2) = 2x_1 + x_2^2 + x_1x_2$$

$$(X_1, X_2)^T \sim \mathcal{N}\left(\begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}\right)$$



$$X_1 \mid X_2 = x_2 \sim \mathcal{N}(0, 1), \quad X_2 \mid X_1 = x_1 \sim \mathcal{N}(0, 1)$$

Components:

$$y_{\emptyset} = 1, \quad y_{\{1\}}(x_1) = 2x_1, \quad y_{\{2\}}(x_2) = x_2^2 - 1, \quad y_{\{1,2\}}(x_1, x_2) = x_1x_2$$

Visualization of fANOVA components under Independence

$$y(x_1, x_2) = 2x_1 + x_2^2 + x_1x_2$$





Outline

- 1 Research Context
- 2 Classical fANOVA
- 3 Generalized fANOVA**
- 4 Conclusion

Visualization of fANOVA components under Dependence

$$y(x_1, x_2) = 2x_1 + x_2^2 + x_1x_2, \quad \rho = 0.8$$



Weak Annihilating Conditions

$$\int_{\mathbb{R}} y_{u,G}(\mathbf{x}_u) f_{\mathbf{u}}(\mathbf{x}_u) d\nu(x_i) = 0 \quad \text{for } i \in u \neq \emptyset$$

Weak Annihilating Conditions

$$\int_{\mathbb{R}} y_{u,G}(\mathbf{x}_u) f_{\mathbf{u}}(\mathbf{x}_u) d\nu(x_i) = 0 \quad \text{for } i \in u \neq \emptyset$$

- $f_{\{i\}}$: marginal density of variable X_u (possibly multivariate)

It follows:

Weak Annihilating Conditions

$$\int_{\mathbb{R}} y_{u,G}(\mathbf{x}_u) f_{\mathbf{u}}(\mathbf{x}_u) d\nu(x_i) = 0 \quad \text{for } i \in u \neq \emptyset$$

- $f_{\{i\}}$: marginal density of variable X_u (possibly multivariate)

It follows:

$$\mathbb{E}[y_{u,G}(\mathbf{x}_u)] := \int_{\mathbb{R}^N} y_{u,G}(\mathbf{x}_u) f_{\mathbf{x}}(\mathbf{x}) d\nu(\mathbf{x}) = 0$$

Weak Annihilating Conditions

$$\int_{\mathbb{R}} y_{u,G}(\mathbf{x}_u) f_{\mathbf{u}}(\mathbf{x}_u) d\nu(x_i) = 0 \quad \text{for } i \in u \neq \emptyset$$

- $f_{\{i\}}$: marginal density of variable X_u (possibly multivariate)

It follows:

$$\mathbb{E}[y_{u,G}(\mathbf{x}_u)] := \int_{\mathbb{R}^N} y_{u,G}(\mathbf{x}_u) f_{\mathbf{x}}(\mathbf{x}) d\nu(\mathbf{x}) = 0$$

$$\mathbb{E}[y_{u,G}(\mathbf{x}_u) y_{v,G}(\mathbf{x}_v)] := \int_{\mathbb{R}^N} y_{u,G}(\mathbf{x}_u) y_{v,G}(\mathbf{x}_v) f_{\mathbf{x}}(\mathbf{x}) d\nu(\mathbf{x}) = 0 \quad (v \subsetneq u)$$

$$y_{\emptyset, G} = \int_{\mathbb{R}^N} y(\mathbf{x}) f_{\mathbf{x}}(\mathbf{x}) d\nu(\mathbf{x})$$

$$y_{\emptyset,G} = \int_{\mathbb{R}^N} y(\mathbf{x}) f_{\mathbf{x}}(\mathbf{x}) d\nu(\mathbf{x})$$

$$\begin{aligned} y_{u,G}(\mathbf{x}_u) &= \int_{\mathbb{R}^{N-|u|}} y(\mathbf{x}_u, \mathbf{x}_{-u}) f_{-u}(\mathbf{x}_{-u}) d\nu(\mathbf{x}_{-u}) - \sum_{v \subsetneq u} y_{v,G}(\mathbf{x}_v) \\ &\quad - \sum_{\substack{\emptyset \neq v \subseteq \{1, \dots, N\} \\ v \cap u \neq \emptyset, v \not\subseteq u}} \int_{\mathbb{R}^{|v \cap u|}} y_{v,G}(\mathbf{x}_{v \cap u}, \mathbf{x}_{v \cap u}) f_{v \cap u}(\mathbf{x}_{v \cap u}) d\nu(\mathbf{x}_{v \cap u}) \end{aligned}$$

$$y_{\emptyset, G} = \int_{\mathbb{R}^N} y(\mathbf{x}) f_{\mathbf{x}}(\mathbf{x}) d\nu(\mathbf{x})$$

$$\begin{aligned} y_{u, G}(\mathbf{X}_u) &= \int_{\mathbb{R}^{N-|u|}} y(\mathbf{X}_u, \mathbf{x}_{-u}) f_{-u}(\mathbf{x}_{-u}) d\nu(\mathbf{x}_{-u}) - \sum_{v \subsetneq u} y_{v, G}(\mathbf{X}_v) \\ &\quad - \sum_{\substack{\emptyset \neq v \subseteq \{1, \dots, N\} \\ v \cap u \neq \emptyset, v \not\subseteq u}} \int_{\mathbb{R}^{|v \cap u|}} y_{v, G}(\mathbf{X}_{v \cap u}, \mathbf{x}_{v \cap u}) f_{v \cap u}(\mathbf{x}_{v \cap u}) d\nu(\mathbf{x}_{v \cap u}) \end{aligned}$$

- $u = \{1\} \rightarrow v \subsetneq u \in \{\emptyset\}$

$$y_{\emptyset, G} = \int_{\mathbb{R}^N} y(\mathbf{x}) f_{\mathbf{x}}(\mathbf{x}) d\nu(\mathbf{x})$$

$$\begin{aligned} y_{u, G}(\mathbf{X}_u) &= \int_{\mathbb{R}^{N-|u|}} y(\mathbf{X}_u, \mathbf{x}_{-u}) f_{-u}(\mathbf{x}_{-u}) d\nu(\mathbf{x}_{-u}) - \sum_{v \subsetneq u} y_{v, G}(\mathbf{X}_v) \\ &\quad - \sum_{\substack{\emptyset \neq v \subseteq \{1, \dots, N\} \\ v \cap u \neq \emptyset, v \not\subseteq u}} \int_{\mathbb{R}^{|v \cap u|}} y_{v, G}(\mathbf{X}_{v \cap u}, \mathbf{x}_{v \cap u}) f_{v \cap u}(\mathbf{x}_{v \cap u}) d\nu(\mathbf{x}_{v \cap u}) \end{aligned}$$

- $u = \{1\} \rightarrow v \subsetneq u \in \{\emptyset\}$
- $(\emptyset \neq v \subseteq \{1, \dots, N\}, v \cap u \neq \emptyset, v \not\subseteq u) \in \{\{1, 2\}, \{1, 3\}, \{1, 2, 3\}\}$

- $N = 3, u = \emptyset$

$$y_{\emptyset, G} = \mathbb{E}[y(\mathbf{X})]$$

- $u = \{1\} \rightarrow v \subsetneq u \in \{\emptyset\}$ and

$$(\emptyset \neq v \subseteq \{1, \dots, N\}, v \cap u \neq \emptyset, v \not\subseteq u) \in \{\{1, 2\}, \{1, 3\}, \{1, 2, 3\}\}$$

$$\begin{aligned} y_{\{1\}, G}(\mathbf{X}_u) &= \int_{\mathbb{R}^2} y(x_1, x_2, x_3) f_{\{2,3\}}(x_2, x_3) d\nu(x_2, x_3) - y_{\emptyset, G} \\ &\quad - \int_{\mathbb{R}} y_{\{1,2\}, G}(x_1, x_2) f_{\{2\}}(x_2) d\nu(x_2) - \int_{\mathbb{R}} y_{\{1,3\}, G}(x_1, x_3) f_{\{3\}}(x_3) d\nu(x_3) \\ &\quad - \int_{\mathbb{R}^2} y_{\{1,2,3\}, G}(x_1, x_2, x_3) f_{\{2,3\}}(x_2, x_3) d\nu(x_2, x_3) \end{aligned}$$

- $u = \{1, 2\} \rightarrow v \subsetneq u \in \{\emptyset, \{1\}, \{2\}\}$ and
 $(\emptyset \neq v \subseteq \{1, \dots, N\}, v \cap u \neq \emptyset, v \not\subseteq u) \in \{\{1, 2, 3\}\}$

$$\begin{aligned} y_{\{1,2\},G}(\mathbf{x}_u) &= \int_{\mathbb{R}} y(x_1, x_2, x_3) f_{\{3\}}(x_3) d\nu(x_3) - y_{\emptyset,G} - y_{\{1\},G} - y_{\{2\},G} \\ &\quad - \int_{\mathbb{R}} y_{\{1,2,3\},G}(x_1, x_2, x_3) f_{\{3\}}(x_3) d\nu(x_3) \end{aligned}$$

- $u = \{1, 2, 3\} \rightarrow v \subsetneq u \in \{\emptyset, \{1\}, \{2\}, \{3\}, \{1, 2\}, \{1, 3\}, \{2, 3\}\}$ and
 $\nexists v \subseteq \{1, \dots, N\}$ s.t. $v \neq \emptyset, v \cap u \neq \emptyset, v \not\subseteq u$

$$y_{\{1,2,3\},G}(\mathbf{x}_u) = y(x_1, x_2, x_3) - y_{\emptyset,G}$$

$$- y_{\{1\},G} - y_{\{2\},G} - y_{\{3\},G}$$

$$- y_{\{1,2\},G} - y_{\{1,3\},G} - y_{\{2,3\},G}$$

$$y(x_1, x_2) = a_0 + a_1x_1 + a_2x_2 + a_{11}x_1^2 + a_{22}x_2^2 + a_{12}x_1x_2$$

$$\begin{aligned}y(x_1, x_2) &= a_0 + a_1x_1 + a_2x_2 + a_{11}x_1^2 + a_{22}x_2^2 + a_{12}x_1x_2 \\ &= c_0 + c_{1,1} \psi_{1,1}(x_1) + c_{2,1} \psi_{2,1}(x_2)\end{aligned}$$

$$\begin{aligned}y(x_1, x_2) &= a_0 + a_1 x_1 + a_2 x_2 + a_{11} x_1^2 + a_{22} x_2^2 + a_{12} x_1 x_2 \\&= c_0 + c_{1,1} \psi_{1,1}(x_1) + c_{2,1} \psi_{2,1}(x_2) \\&\quad + c_{1,2} \psi_{1,2}(x_1) + c_{2,2} \psi_{2,2}(x_2) + c_{12,11} \psi_{12,11}(x_1, x_2)\end{aligned}$$

$$\begin{aligned}y(x_1, x_2) &= a_0 + a_1 x_1 + a_2 x_2 + a_{11} x_1^2 + a_{22} x_2^2 + a_{12} x_1 x_2 \\&= c_0 + c_{1,1} \psi_{1,1}(x_1) + c_{2,1} \psi_{2,1}(x_2) \\&\quad + c_{1,2} \psi_{1,2}(x_1) + c_{2,2} \psi_{2,2}(x_2) + c_{12,11} \psi_{12,11}(x_1, x_2) \\&= \underbrace{c_0}_{y_0} + \underbrace{(c_{1,1} \psi_{1,1}(x_1) + c_{1,2} \psi_{1,2}(x_1))}_{y_1(x_1)}\end{aligned}$$

$$\begin{aligned}y(x_1, x_2) &= a_0 + a_1 x_1 + a_2 x_2 + a_{11} x_1^2 + a_{22} x_2^2 + a_{12} x_1 x_2 \\&= c_0 + c_{1,1} \psi_{1,1}(x_1) + c_{2,1} \psi_{2,1}(x_2) \\&\quad + c_{1,2} \psi_{1,2}(x_1) + c_{2,2} \psi_{2,2}(x_2) + c_{12,11} \psi_{12,11}(x_1, x_2) \\&= \underbrace{c_0}_{y_0} + \underbrace{(c_{1,1} \psi_{1,1}(x_1) + c_{1,2} \psi_{1,2}(x_1))}_{y_1(x_1)} \\&\quad + \underbrace{(c_{2,1} \psi_{2,1}(x_2) + c_{2,2} \psi_{2,2}(x_2))}_{y_2(x_2)}\end{aligned}$$

$$\begin{aligned}y(x_1, x_2) &= a_0 + a_1 x_1 + a_2 x_2 + a_{11} x_1^2 + a_{22} x_2^2 + a_{12} x_1 x_2 \\&= c_0 + c_{1,1} \psi_{1,1}(x_1) + c_{2,1} \psi_{2,1}(x_2) \\&\quad + c_{1,2} \psi_{1,2}(x_1) + c_{2,2} \psi_{2,2}(x_2) + c_{12,11} \psi_{12,11}(x_1, x_2) \\&= \underbrace{c_0}_{y_0} + \underbrace{(c_{1,1} \psi_{1,1}(x_1) + c_{1,2} \psi_{1,2}(x_1))}_{y_1(x_1)} \\&\quad + \underbrace{(c_{2,1} \psi_{2,1}(x_2) + c_{2,2} \psi_{2,2}(x_2))}_{y_2(x_2)} \\&\quad + \underbrace{c_{12,11} \psi_{12,11}(x_1, x_2)}_{y_{12}(x_1, x_2)}\end{aligned}$$

For Gaussian input variables Hermite polynomial basis functions are proposed:

$$\psi_{\emptyset}(x_1, x_2) = 1,$$

$$\psi_{1,1}(x_1) = x_1,$$

$$\psi_{2,1}(x_2) = x_2,$$

$$\psi_{1,2}(x_1) = x_1^2 - 1,$$

$$\psi_{2,2}(x_2) = x_2^2 - 1,$$

$$\psi_{12,11}(x_1, x_2) = \frac{\rho(x_1^2 + x_2^2)}{1 + \rho^2} - x_1x_2 + \frac{\rho(\rho^2 - 1)}{1 + \rho^2}$$

- Yields fANOVA components for Gaussian Inputs
- Works for polynomials of degree up to $d = 2$



$$y_{\emptyset,G} = a_0 + a_{11} + a_{22} + \rho a_{12},$$

$$y_{\{1\},G}(x_1) = a_1 x_1 + \left(a_{11} + \frac{\rho}{1 + \rho^2} a_{12} \right) (x_1^2 - 1),$$



$$y_{\{2\},G}(x_2) = a_2 x_2 + \left(a_{22} + \frac{\rho}{1 + \rho^2} a_{12} \right) (x_2^2 - 1),$$

$$y_{\{1,2\},G}(x_1, x_2) = -a_{12} \left(\frac{\rho(x_1^2 + x_2^2)}{1 + \rho^2} - x_1 x_2 + \frac{\rho(\rho^2 - 1)}{1 + \rho^2} \right)$$

$$z(x_1, x_2) = 2x_1 + x_1^2 + 0.5x_1x_2$$



$$z(x_1, x_2) = 2x_1 + x_1^2 + 0.5x_1x_2$$



→ nonzero main effect of X_2 only present under correlation

Variance Decomposition via fANOVA

Not only y , but also its variance can be decomposed:

$$\mu := \mathbb{E}[y(\mathbf{X})] = y_{\emptyset, G}$$

Variance Decomposition via fANOVA

Not only y , but also its variance can be decomposed:

$$\mu := \mathbb{E}[y(\mathbf{X})] = y_{\emptyset, G}$$

$$\sigma^2 := \mathbb{E} \left[(y(\mathbf{X}) - \mu)^2 \right]$$

Variance Decomposition via fANOVA

Not only y , but also its variance can be decomposed:

$$\mu := \mathbb{E}[y(\mathbf{X})] = y_{\emptyset, G}$$

$$\begin{aligned}\sigma^2 &:= \mathbb{E} \left[(y(\mathbf{X}) - \mu)^2 \right] \\ &= \mathbb{E} \left[\left(y_{\emptyset, G} + \sum_u y_{u, G}(\mathbf{x}_u) - y_{\emptyset, G} \right)^2 \right]\end{aligned}$$

Variance Decomposition via fANOVA

Not only y , but also its variance can be decomposed:

$$\mu := \mathbb{E}[y(\mathbf{X})] = y_{\emptyset, G}$$

$$\begin{aligned}\sigma^2 &:= \mathbb{E} \left[(y(\mathbf{X}) - \mu)^2 \right] \\ &= \mathbb{E} \left[\left(y_{\emptyset, G} + \sum_u y_{u, G}(\mathbf{x}_u) - y_{\emptyset, G} \right)^2 \right] \\ &= \mathbb{E} \left[\left(\sum_u y_{u, G}(\mathbf{x}_u) \right)^2 \right]\end{aligned}$$

Variance Decomposition via fANOVA

Not only y , but also its variance can be decomposed:

$$\mu := \mathbb{E}[y(\mathbf{X})] = y_{\emptyset, G}$$

$$\sigma^2 := \mathbb{E} \left[(y(\mathbf{X}) - \mu)^2 \right]$$

$$= \mathbb{E} \left[\left(y_{\emptyset, G} + \sum_u y_{u, G}(\mathbf{x}_u) - y_{\emptyset, G} \right)^2 \right]$$

$$= \mathbb{E} \left[\left(\sum_u y_{u, G}(\mathbf{x}_u) \right)^2 \right]$$

$$= \sum_u \mathbb{E} [y_{u, G}^2(\mathbf{x}_u)] + \sum_{u \not\subseteq v, v \not\subseteq u} \mathbb{E} [y_{u, G}(\mathbf{x}_u) y_{v, G}(\mathbf{x}_v)]$$

Different formulations of generalized fANOVA components exists, e.g.:


$$\{y_{u,G}(\mathbf{x}_u) \mid u \subseteq d\} = \arg \min_{\{g_u \in \mathcal{L}^2(\mathbb{R}^{|u|})\}} \int_{\mathbb{R}^N} \left(\sum_{u \subseteq d} g_u(\mathbf{x}_u) - y(\mathbf{x}) \right)^2 f_{\mathbf{x}}(\mathbf{x}) d\nu(\mathbf{x}),$$

subject to hierarchical orthogonality conditions:

$$\forall v \subseteq u, \forall g_v : \int_{\mathbb{R}^N} y_u(\mathbf{x}_u) g_v(\mathbf{x}_v) f_{\mathbf{x}}(\mathbf{x}) d\nu(\mathbf{x}) = 0.$$

Outline

- 1 Research Context
- 2 Classical fANOVA
- 3 Generalized fANOVA
- 4 Conclusion**



$$\Pi_{\mathcal{G}}y = \arg \min_{g \in \mathcal{G}} \|y - g\|^2 = \arg \min_{g \in \mathcal{G}} \mathbb{E}[(y(\mathbf{X}) - g(\mathbf{X}))^2]$$

- \mathcal{G} : linear subspace of \mathcal{L}^2 we project onto
- g all functions in the subspace

$$\begin{aligned}y_{\emptyset} &= \mathbb{E}[y(\mathbf{X})] \\&= \arg \min_{a \in \mathbb{R}} \mathbb{E}[(y(\mathbf{X}) - a)^2] \\&= \arg \min_{g_0 \in \mathcal{G}_0} \|y - g_0\|^2 = \Pi_{\mathcal{G}_0} y\end{aligned}$$

$$\begin{aligned}y_u(.) &= \mathbb{E}[y(\mathbf{X}) \mid X_u = .] - \sum_{v \subsetneq u} y_v(.) \\&= \arg \min_{g_u \in \mathcal{G}_u} \mathbb{E}[(y(\mathbf{X}) - g_u(.))^2] - \sum_{v \subsetneq u} y_v(.) \\&= (\Pi_{\mathcal{G}_u} y)(.) - \sum_{v \subsetneq u} y_v(.)\end{aligned}$$

Hoeffding Decomposition

$$y(\mathbf{x}) = \sum_{A \subseteq D} y_A(\mathbf{x}_A), \quad D := \{1, \dots, N\}, \quad (1)$$

where, for each $A \subseteq D$, the component function y_A is defined by:

$$y_A(\mathbf{x}_A) = \sum_{B \subseteq A} (-1)^{|A|-|B|} \mathbb{E}[y(\mathbf{x}) \mid \mathbf{x}_B], \quad (2)$$

where y_u are orthogonal components.

- Classical fANOVA and Hoeffding decomposition yield same components under zero-centered inputs
- Both assume independence of input variables

$$y(x_1, x_2) = 2x_1 + x_2^2 + x_1x_2$$

$$y_{\emptyset} = \mathbb{E}[y(X_1, X_2)] = 2 \mathbb{E}[X_1] + \mathbb{E}[X_2^2] + \mathbb{E}[X_1X_2] = 1,$$

$$\begin{aligned} y_{\{1\}}(x_1) &= \sum_{B \subseteq \{1\}} (-1)^{1-|B|} \mathbb{E}[y(\mathbf{X}) | X_B] = -\mathbb{E}[y] + \mathbb{E}[y | X_1 = x_1] \\ &= -1 + (2x_1 + \mathbb{E}[X_2^2] + x_1\mathbb{E}[X_2]) = 2x_1, \end{aligned}$$

$$\begin{aligned} y_{\{2\}}(x_2) &= \sum_{B \subseteq \{2\}} (-1)^{1-|B|} \mathbb{E}[y(\mathbf{X}) | X_B] - \mathbb{E}[y] + \mathbb{E}[y | X_2 = x_2] \\ &= -1 + (2\mathbb{E}[X_1] + x_2^2 + x_2\mathbb{E}[X_1]) = x_2^2 - 1. \end{aligned}$$

$$\begin{aligned}
y_{\{1,2\}}(x_1, x_2) &= \sum_{B \subseteq \{1,2\}} (-1)^{2-|B|} \mathbb{E}[y(\mathbf{x}) | X_B] \\
&= (+1) \mathbb{E}[y] - \mathbb{E}[y | X_1 = x_1] - \mathbb{E}[y | X_2 = x_2] + y(x_1, x_2) \\
&= 1 - (2x_1 + 1) - (x_2^2) + (2x_1 + x_2^2 + x_1x_2) \\
&= x_1x_2.
\end{aligned}$$

$$y(x_1, x_2) = y_{\emptyset} + y_{\{1\}}(x_1) + y_{\{2\}}(x_2) + y_{\{1,2\}}(x_1, x_2) = 1 + 2x_1 + (x_2^2 - 1) + x_1x_2$$

Substituting the basis functions:

$$\begin{aligned}
 y(x_1, x_2) = & \underbrace{c_0}_{y_0} + \underbrace{(c_{1,1} x_1 + c_{1,2} (x_1^2 - 1))}_{y_1(x_1)} \\
 & + \underbrace{(c_{2,1} x_2 + c_{2,2} (x_2^2 - 1))}_{y_2(x_2)} \\
 & + \underbrace{c_{12,11} \left(\frac{\rho(x_1^2 + x_2^2)}{1 + \rho^2} - x_1 x_2 + \frac{\rho(\rho^2 - 1)}{1 + \rho^2} \right)}_{y_{12}(x_1, x_2)}.
 \end{aligned}$$

Find weights to recover original polynomial while fulfilling zero-mean and hierarchical orthogonality:

$$y(x_1, x_2) = a_0 + a_1 x_1 + a_2 x_2 + a_{11} x_1^2 + a_{22} x_2^2 + a_{12} x_1 x_2$$

The corresponding weights can be found via coefficient matching. Start from the interaction term:

$$-c_{12,11} = a_{12} \quad \Rightarrow \quad c_{12,11} = -a_{12}$$

$$c_{1,2} + c_{12,11} \frac{\rho}{1+\rho^2} = a_{11} \quad \Rightarrow \quad c_{1,2} = a_{11} + \frac{\rho}{1+\rho^2} a_{12}$$

$$c_{2,2} + c_{12,11} \frac{\rho}{1+\rho^2} = a_{22} \quad \Rightarrow \quad c_{2,2} = a_{22} + \frac{\rho}{1+\rho^2} a_{12}$$



$$c_{1,1} = a_1$$

$$c_{2,1} = a_2$$

$$c_0 - c_{1,2} - c_{2,2} + c_{12,11} \frac{\rho(\rho^2-1)}{1+\rho^2} = a_0 \quad \Rightarrow \quad c_0 = a_0 + a_{11} + a_{22} + \rho a_{12}$$



$$h(x_1, x_2) = x_1 + 2x_2 + x_1x_2 \quad \rho = 0$$

(3)



$$h(x_1, x_2) = x_1 + 2x_2 + x_1x_2 \quad \rho = 0.5$$

(4)



Example: Only Linear Terms




Figure: $q(x_1, x_2) = 1.5x_1 + 3.5x_2$






Figure: $q(x_1, x_2) = -2x_1 + 4x_2$



Example: Only Main Terms under Independence

$$y(x_1, x_2) = -2x_1 - 2x_2 + x_1^2 + x_2^2 \quad \rho = 0$$





Example: Only Interaction Term under Dependence

$$y(x_1, x_2) = x_1 x_2 \quad \rho = -0.5$$



Example: Interaction under Independence

$$y(x_1, x_2) = x_1 x_2 \quad \rho = 0$$



Strong annihilating conditions hold, so:

$$\begin{aligned}\mathbb{E}[y_u(\mathbf{X}_u)] &:= \int_{\mathbb{R}^N} y_u(\mathbf{x}_u) f_{\mathbf{X}}(\mathbf{x}) d\nu(\mathbf{x}) \\ &= \int_{\mathbb{R}^{|u|}} y_u(\mathbf{x}_u) f_{\mathbf{u}}(\mathbf{x}_u) d\nu(\mathbf{x}_u) \\ &= \int_{\mathbb{R}^{|u|}} y_u(\mathbf{x}_u) \prod_{j \in u} f_{\{j\}}(x_j) d\nu(\mathbf{x}_u) \\ &= \int_{\mathbb{R}^{|u|-1}} \int_{\mathbb{R}} y_u(\mathbf{x}_u) f_{\{i\}}(x_i) d\nu(x_i) \prod_{j \in u, j \neq i} f_{\{j\}}(x_j) d\nu(x_{u \setminus \{i\}}) = 0.\end{aligned}$$

Proof of Orthogonality for Classical Components

- $u \neq v$, so pick $i \in u \setminus v$
- $y_v(\mathbf{x}_v)$ is independent of x_i
- strong annihilating conditions hold by assumption

$$\int_{\mathbb{R}} y_u(\mathbf{x}_u) f_{\{i\}}(x_i) d\nu(x_i) = 0 \quad \text{for all fixed } \mathbf{x}_{u \setminus \{i\}}.$$

Hence,

$$\begin{aligned} \mathbb{E}[y_u(\mathbf{X}_u) y_v(\mathbf{X}_v)] &= \int_{\mathbb{R}^N} y_u(\mathbf{x}_u) y_v(\mathbf{x}_v) f_{\mathbf{X}}(\mathbf{x}) d\nu(\mathbf{x}) \\ &= \int_{\mathbb{R}^N} y_u(\mathbf{x}_u) y_v(\mathbf{x}_v) \prod_{j=1}^N f_{\{j\}}(x_j) d\nu(\mathbf{x}) \\ &= \int_{\mathbb{R}^{N-1}} \left(\int_{\mathbb{R}} y_u(\mathbf{x}_u) f_{\{i\}}(x_i) d\nu(x_i) \right) y_v(\mathbf{x}_v) \prod_{j \neq i} f_{\{j\}}(x_j) d\nu(\mathbf{x}_{-i}) = 0. \end{aligned}$$

We assume the weak annihilating conditions hold, then:

$$\begin{aligned}\mathbb{E}[y_{u,G}(\mathbf{X}_u)] &:= \int_{\mathbb{R}^N} y_{u,G}(\mathbf{x}_u) f_{\mathbf{X}}(\mathbf{x}) d\nu(\mathbf{x}) \\&= \int_{\mathbb{R}^{|u|}} y_{u,G}(\mathbf{x}_u) \left(\int_{\mathbb{R}^{N-|u|}} f_{\mathbf{X}}(\mathbf{x}) d\nu(\mathbf{x}_{-u}) \right) d\nu(\mathbf{x}_u) \\&= \int_{\mathbb{R}^{|u|}} y_{u,G}(\mathbf{x}_u) f_u(\mathbf{x}_u) d\nu(\mathbf{x}_u) \\&= \int_{\mathbb{R}^{|u|-1}} \left(\int_{\mathbb{R}} y_{u,G}(\mathbf{x}_u) f_u(\mathbf{x}_u) d\nu(x_i) \right) \prod_{j \in u, j \neq i} d\nu(\mathbf{x}_j) \\&= 0.\end{aligned}$$

Proof of Hierarchical Orthogonality

For any two subsets $\emptyset \neq u \subseteq \{1, \dots, N\}$ and $\emptyset \neq v \subseteq \{1, \dots, N\}$, where $v \subsetneq u$, the subset $u = v \cup (u \setminus v)$. Let $i \in (u \setminus v) \subseteq u$. Then we obtain:

$$\begin{aligned}\mathbb{E}[y_{u,G}(\mathbf{X}_u) y_{v,G}(\mathbf{X}_v)] &:= \int_{\mathbb{R}^N} y_{u,G}(\mathbf{x}_u) y_{v,G}(\mathbf{x}_v) f_{\mathbf{X}}(\mathbf{x}) d\nu(\mathbf{x}) \\&= \int_{\mathbb{R}^{|u|}} y_{u,G}(\mathbf{x}_u) y_{v,G}(\mathbf{x}_v) \left(\int_{\mathbb{R}^{N-|u|}} f_{\mathbf{X}}(\mathbf{x}) d\nu(\mathbf{x}_{-u}) \right) d\nu(\mathbf{x}_u) \\&= \int_{\mathbb{R}^{|u|}} y_{u,G}(\mathbf{x}_u) y_{v,G}(\mathbf{x}_v) f_u(\mathbf{x}_u) d\nu(\mathbf{x}_u) \\&= \int_{\mathbb{R}^{|v|}} y_{v,G}(\mathbf{x}_v) \int_{\mathbb{R}^{|u \setminus v|}} y_{u,G}(\mathbf{x}_u) f_u(\mathbf{x}_u) d\nu(\mathbf{x}_{u \setminus v}) d\nu(\mathbf{x}_v) \\&= \int_{\mathbb{R}^{|v|}} y_{v,G}(\mathbf{x}_v) \int_{\mathbb{R}^{|u \setminus v|-1}} \left(\int_{\mathbb{R}} y_{u,G}(\mathbf{x}_u) f_u(\mathbf{x}_u) d\nu(\mathbf{x}_i) \right) \\&\quad \times \prod d\nu(x_j) d\nu(\mathbf{x}_v) = 0.\end{aligned}$$

$$S_u = \frac{\text{Var}(\mathbb{E}[y(\mathbf{X}) \mid \mathbf{X}_u = .])}{\text{Var}(y(\mathbf{X}))},$$

where

- $y(\mathbf{X})$ is the probabilistic model, which is decomposed
- $\mathbb{E}[y(\mathbf{X}) \mid \mathbf{X}_u = .]$ is the fANOVA component y_u

- https://docs.google.com/spreadsheets/d/1K5ECL6hDPDnHwM_k342xa29H-vHWzdk27PTgDHUwfFE/edit?usp=sharing - Table with fANOVA-related literature



Wassily Hoeffding.

A Class of Statistics with Asymptotically Normal Distribution.

The Annals of Mathematical Statistics, 19(3):293–325, 1948.

Publisher: Institute of Mathematical Statistics.



Giles Hooker.

Discovering additive structure in black box functions.

In Proceedings of the tenth ACM SIGKDD international conference on Knowledge discovery and data mining, pages 575–580, Seattle WA USA, August 2004. ACM.



Giles Hooker.

Generalized Functional ANOVA Diagnostics for High-Dimensional Functions of Dependent Variables.

Journal of Computational and Graphical Statistics, 16(3):709–732, September 2007.



Linwei Hu, Vijayan N. Nair, Agus Sudjianto, Aijun Zhang, Jie Chen, and Zebin Yang.
Interpretable Machine Learning Based on Functional ANOVA Framework: Algorithms and Comparisons.

Applied Stochastic Models in Business and Industry, 41(1):e2916, January 2025.



Gunnar König, Eric Günther, and Ulrike von Luxburg.
Disentangling Interactions and Dependencies in Feature Attribution, October 2024.
[arXiv:2410.23772](https://arxiv.org/abs/2410.23772).



Benjamin Lengerich, Sarah Tan, Chun-Hao Chang, Giles Hooker, and Rich Caruana.
Purifying Interaction Effects with the Functional ANOVA: An Efficient Algorithm for Recovering Identifiable Additive Models.

In Proceedings of the Twenty Third International Conference on Artificial Intelligence and Statistics, pages 2402–2412. PMLR, June 2020.
ISSN: 2640-3498.



Thomas Muehlenstaedt, Olivier Roustant, Laurent Carraro, and Sonja Kuhnt.
Data-driven Kriging models based on FANOVA-decomposition.
Statistics and Computing, 22(3):723–738, May 2012.



Sharif Rahman.
A Generalized ANOVA Dimensional Decomposition for Dependent Probability Measures.
SIAM/ASA Journal on Uncertainty Quantification, 2(1):670–697, 2014.



I. M. Sobol.
Sensitivity Estimates for Nonlinear Mathematical Models.
Mathematical Modelling and Computational Experiments, 1:407–414, 1993.