

Your Title

Your Name

July 31, 2025


Showcase
Figure

1 Research Context

2 Classical fANOVA

3 Generalized fANOVA

4 Conclusion



References: [1, 2, 5, 7, 6, 4, 3]

Outline

1 Research Context

2 Classical fANOVA

3 Generalized fANOVA

4 Conclusion

General Form

$$y(\mathbf{x}) = \sum_{u \subseteq \{1, \dots, N\}} y_u(\mathbf{x}_u)$$

- y : Model output
- y_u : Component functions for subset u

Strong Annihilating Conditions

$$\int_{\mathbb{R}} y_u(\mathbf{x}_u) f_{\{i\}}(x_i) d\nu(x_i) = 0 \quad \text{for } i \in u \neq \emptyset.$$

- Ensures unique component functions
- Applies under independent (product-type) input distributions

$$\mathbb{E}[y_u(\mathbf{X}_u)] = 0$$

$$\mathbb{E}[y_u(\mathbf{X}_u)y_v(\mathbf{X}_v)] = 0 \quad (u \neq v)$$



- Zero mean components
- Mutual orthogonality

$$y_{\emptyset} = \int_{\mathbb{R}^N} y(\mathbf{x}) \prod_{i=1}^N f_{\{i\}}(x_i) d\nu(x_i) = \mathbb{E}[y(\mathbf{x})].$$

$$y_u(x_u) = \int y(x) f_{-u}(x_{-u}) dx_{-u} - \sum_{v \subset u} y_v(x_v)$$

- f_{-u} : marginal density of variables not in u
- Components solved sequentially by increasing order



Example: 2D Function



Outline

- 1 Research Context
- 2 Classical fANOVA
- 3 Generalized fANOVA**
- 4 Conclusion

Example with Dependent Inputs ($\rho = 0.8$)



Weak Annihilating Conditions

$$\int_{\mathbb{R}} y_{u,G}(\mathbf{x}_u) f_{\mathbf{x}_u}(\mathbf{x}_u) d\nu(x_i) = 0 \quad \text{for } i \in u \neq \emptyset.$$

- Allows dependent input distributions
- Leads to hierarchical orthogonality

$$\mathbb{E}[y_{u,G}(\mathbf{X}_u)] := \int_{\mathbb{R}^N} y_{u,G}(\mathbf{x}_u) f_{\mathbf{X}}(\mathbf{x}) d\nu(\mathbf{x}) = 0.$$

$$\mathbb{E}[y_{u,G}(\mathbf{X}_u) y_{v,G}(\mathbf{X}_v)] := \int_{\mathbb{R}^N} y_{u,G}(\mathbf{x}_u) y_{v,G}(\mathbf{x}_v) f_{\mathbf{X}}(\mathbf{x}) d\nu(\mathbf{x}) = 0.$$

- Zero mean components remain
- Orthogonality is weaker: hierarchical

$$y_{\emptyset,G} = \int_{\mathbb{R}^N} y(\mathbf{x}) f_{\mathbf{x}}(\mathbf{x}) d\nu(\mathbf{x}) \quad (1)$$

$$\begin{aligned} y_{u,G}(\mathbf{x}_u) &= \int_{\mathbb{R}^{N-|u|}} y(\mathbf{x}_u, \mathbf{x}_{-u}) f_{-u}(\mathbf{x}_{-u}) d\nu(\mathbf{x}_{-u}) - \sum_{v \subsetneq u} y_{v,G}(\mathbf{x}_v) \\ &\quad - \sum_{\substack{\emptyset \neq v \subseteq \{1, \dots, N\} \\ v \cap u \neq \emptyset, v \not\subseteq u}} \int_{\mathbb{R}^{|v \cap u|}} y_{v,G}(\mathbf{x}_{v \cap u}, \mathbf{x}_{v \cap u}) f_{v \cap u}(\mathbf{x}_{v \cap u}) d\nu(\mathbf{x}_{v \cap u}). \end{aligned} \quad (2)$$

- All components solved simultaneously
- Depends on marginal densities and coupling terms

- Coupled system \rightarrow difficult to obtain analytical solutions
- Use alternative method via Fourier Polynomial ([5])
- Building blocks: mutually orthogonal, zero-mean basis functions $\psi_{i,j}$, coefficients $c_{i,j}$

$$\begin{aligned}y(x_1, x_2) &= a_0 + a_1x_1 + a_2x_2 + a_{11}x_1^2 + a_{22}x_2^2 + a_{12}x_1x_2 \\&= c_0 + c_{1,1} \psi_{1,1}(x_1) + c_{2,1} \psi_{2,1}(x_2) \\&\quad + c_{1,2} \psi_{1,2}(x_1) + c_{2,2} \psi_{2,2}(x_2) + c_{12,11} \psi_{12,11}(x_1, x_2) \\&= \underbrace{c_0}_{y_0} + \underbrace{(c_{1,1} \psi_{1,1}(x_1) + c_{1,2} \psi_{1,2}(x_1))}_{y_1(x_1)} \\&\quad + \underbrace{(c_{2,1} \psi_{2,1}(x_2) + c_{2,2} \psi_{2,2}(x_2))}_{y_2(x_2)} \\&\quad + \underbrace{c_{12,11} \psi_{12,11}(x_1, x_2)}_{y_{12}(x_1, x_2)}.\end{aligned}$$

$$\psi_{\emptyset}(x_1, x_2) = 1,$$

$$\psi_{1,1}(x_1) = x_1,$$

$$\psi_{2,1}(x_2) = x_2,$$

$$\psi_{1,2}(x_1) = x_1^2 - 1,$$

$$\psi_{2,2}(x_2) = x_2^2 - 1,$$

$$\psi_{12,11}(x_1, x_2) = \frac{\rho(x_1^2 + x_2^2)}{1 + \rho^2} - x_1x_2 + \frac{\rho(\rho^2 - 1)}{1 + \rho^2},$$

$$\{y_{u,G}(\mathbf{x}_u) \mid u \subseteq d\} = \arg \min_{\{g_u \in \mathcal{L}^2(\mathbb{R}^{|u|})\}} \int_{\mathbb{R}^N} \left(\sum_{u \subseteq d} g_u(\mathbf{x}_u) - y(\mathbf{x}) \right)^2 f_{\mathbf{x}}(\mathbf{x}) d\nu(\mathbf{x}),$$

subject to hierarchical orthogonality conditions:

$$\forall v \subseteq u, \forall g_v : \int_{\mathbb{R}^N} y_u(\mathbf{x}_u) g_v(\mathbf{x}_v) f_{\mathbf{x}}(\mathbf{x}) d\nu(\mathbf{x}) = 0.$$

$$\mu := \mathbb{E}[y(\mathbf{X})] = y_{\emptyset, G}.$$

$$\begin{aligned}\sigma^2 &:= \mathbb{E} \left[(y(\mathbf{X}) - \mu_G)^2 \right] \\ &= \mathbb{E} \left[\left(y_{\emptyset, G} + \sum_u y_{u, G}(\mathbf{x}_u) - y_{\emptyset, G} \right)^2 \right] \\ &= \mathbb{E} \left[\left(\sum_u y_{u, G}(\mathbf{x}_u) \right)^2 \right] \\ &= \sum_u \mathbb{E} [y_{u, G}^2(\mathbf{x}_u)] + \sum_{u \not\subseteq v, v \not\subseteq u} \mathbb{E} [y_{u, G}(\mathbf{x}_u) y_{v, G}(\mathbf{x}_v)],\end{aligned}$$

Outline

- 1 Research Context
- 2 Classical fANOVA
- 3 Generalized fANOVA
- 4 Conclusion**



Giles Hooker.

Discovering additive structure in black box functions.

In Proceedings of the tenth ACM SIGKDD international conference on Knowledge discovery and data mining, pages 575–580, Seattle WA USA, August 2004. ACM.



Giles Hooker.

Generalized Functional ANOVA Diagnostics for High-Dimensional Functions of Dependent Variables.

Journal of Computational and Graphical Statistics, 16(3):709–732, September 2007.



Gunnar König, Eric Günther, and Ulrike von Luxburg.

Disentangling Interactions and Dependencies in Feature Attribution, October 2024.
[arXiv:2410.23772](https://arxiv.org/abs/2410.23772).



Benjamin Lengerich, Sarah Tan, Chun-Hao Chang, Giles Hooker, and Rich Caruana.
Purifying Interaction Effects with the Functional ANOVA: An Efficient Algorithm for
Recovering Identifiable Additive Models.

*In Proceedings of the Twenty Third International Conference on Artificial Intelligence and
Statistics*, pages 2402–2412. PMLR, June 2020.

ISSN: 2640-3498.



Sharif Rahman.

A Generalized ANOVA Dimensional Decomposition for Dependent Probability Measures.

SIAM/ASA Journal on Uncertainty Quantification, 2(1):670–697, 2014.



I. M. Sobol.

Sensitivity Estimates for Nonlinear Mathematical Models.

Mathematical Modelling and Computational Experiments, 1:407–414, 1993.



Charles J. Stone.

The Use of Polynomial Splines and Their Tensor Products in Multivariate Function Estimation.

The Annals of Statistics, 22(1):118–171, 1994.

Publisher: Institute of Mathematical Statistics.