

Bachelor's Thesis

---

# Functional ANOVA Decomposition for Model Interpretability

---

Department of Statistics  
Ludwig-Maximilians-Universität München

**Juliet Fleischer**

Munich, July 31<sup>th</sup>, 2025



Submitted in partial fulfillment of the requirements for the degree of B. Sc.  
Supervised by Prof. Dr. Thomas Nagler

## Abstract

This article studies the functional ANOVA decomposition (fANOVA) in the context of model interpretability. We begin by introducing the classical fANOVA, which assumes independent inputs, and illustrate its equivalence to the Hoeffding decomposition under zero-centered variables with an example. We then unify different notations under the concept of orthogonal projections and briefly present the variance decomposition. Next, we extend fANOVA to settings with dependent inputs, discussing two different formalizations and highlighting why one is more suitable for deriving an explicit solution in an exemplary decomposition, while the other remains primarily theoretical. Finally, we adopt an applied perspective, visualizing the decomposition of various functions and providing a conceptual overview of current estimation approaches.

# Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Background and Related Work</b>	<b>3</b>
<b>3</b>	<b>Formalization of fANOVA</b>	<b>5</b>
3.1	Classical fANOVA . . . . .	6
3.1.1	Construction of the fANOVA Terms . . . . .	8
3.1.2	Example: Independent Multivariate Normal Inputs . . . . .	8
3.1.3	Equality to Hoeffding Decomposition . . . . .	10
3.1.4	fANOVA via Projection . . . . .	12
3.1.5	Variance Decomposition . . . . .	14
3.2	Generalized fANOVA . . . . .	17
3.2.1	Conditions for Generalized fANOVA . . . . .	18
3.2.2	Construction of the Generalized fANOVA Terms . . . . .	21
3.2.3	Generalization via Projection . . . . .	25
3.2.4	Generalized Variance Decomposition . . . . .	27
3.2.5	Example: Dependent Multivariate Normal Inputs . . . . .	28
<b>4</b>	<b>Visualization and Estimation</b>	<b>33</b>
4.1	Comparison of Decompositions . . . . .	33
4.2	Comparison of Functions . . . . .	35
4.2.1	Scenario: Linear . . . . .	35
4.2.2	Scenario: Linear and Quadratic . . . . .	35
4.2.3	Scenario: Interaction . . . . .	36
4.2.4	Scenario: Full . . . . .	38
4.3	Estimation of fANOVA components . . . . .	40
<b>5</b>	<b>Conclusion</b>	<b>43</b>
<b>A</b>	<b>Appendix</b>	<b>V</b>
<b>B</b>	<b>Electronic appendix</b>	<b>VIII</b>

# 1 Introduction

With the rise of machine learning (ML) and increasingly more complex probabilistic models, interpretability has become a major concern for practitioners and researchers alike. One of the foundational mathematical methods supporting the goal of interpretability is the functional ANOVA decomposition (fANOVA).

At its core, the fANOVA decomposition provides a method which allows decomposing integrable functions into a sum of orthogonal components. It is not only useful in interpretability of black box models (Hooker (2004), Molnar (2025)), but also in areas, such as uncertainty quantification of complex systems Rahman (2014), non-parametric statistical modelling (see for example Stone et al. (1997)), sensitivity analysis Sobol (1993)), and many more fields. Given this wide range of applications, fANOVA is an essential concept worth understanding in depth.

However, learning about fANOVA is not straightforward. A problem is the mix of formalizations and definitions around the method, partly due to its long history and the different streams of science that have used it. This already starts with the name of the method. It has been called decomposition into summands of different order (Sobol, 1993), ANOVA representation (Sobol, 2001), functional ANOVA decomposition (Hooker, 2004), ANOVA dimensional decomposition (Rahman, 2014), or Hoeffding-Sobol decomposition (Chastaing et al., 2012) – in this thesis we will refer to it as the fANOVA decomposition. The variation does not stop at naming. Different authors formalize the decomposition using different notation, slightly different sets of assumptions, and either interpret fANOVA from a probabilistic perspective, using expectations, or from a more deterministic mathematical viewpoint, using integrals. While these approaches are mathematically equivalent and can be unified under the concept of orthogonal projections, this connection is often not obvious when first encountering the literature.

Given this state of affairs, there is a clear need for a comprehensive overview of fANOVA-related work and for a unification of the various notations and definitions that ultimately express the same concepts. Bringing clarity into the fANOVA landscape is more relevant than ever as the method has recently attracted renewed attention in interpretable machine learning (IML) literature (see for example Hu et al. (2025)), yet the theoretical foundation is often mentioned only briefly or left implicit.

This thesis addresses that gap by providing an accessible and intuitive introduction to the fANOVA decomposition while remaining mathematically rigorous. It can be viewed as a handbook of the fANOVA decomposition that will help researchers and practitioners to understand the mathematical background of this method as well as its more applied aspects.

This work is organized as follows: It starts with background and related work (section 2). This is followed by the central part in which we give the formal definition of the classical and generalized fANOVA decomposition (section 3). Next we illustrate characteristics of the method based on analytical examples, before briefly outlining current estimation schemes (section 4) and concluding with a discussion and possible future research directions.

## 2 Background and Related Work

The literature around fANOVA can be grouped into several thematic clusters. Each highlights a different angle on why fANOVA has proven useful and points to why a unified presentation is needed.

The underlying principle of the hierarchical, additive decomposition of a function dates back to Hoeffding (1948). In his seminal work on U-statistics, he introduced the Hoeffding decomposition. Though originally framed around estimators, this decomposition laid the groundwork for fANOVA by showing how a symmetric function can be written as a sum of orthogonal components of increasing dimensionality.

Independently, Sobol (1993) proved that any square integrable function on the unit hypercube can be decomposed into a sum of orthogonal and zero-centered components. The foundational work on fANOVA shows, that it is rooted in rigorous mathematical theory, and provides a principled way to break down complex multivariate functions into interpretable, orthogonal parts.

A second strand of work explores how fANOVA underlies non-parametric modeling. Takemura (1983) introduced tensor-analysis of ANOVA decompositions, laying the theoretical foundation. Stone (1994) applied fANOVA ideas to polynomial splines and generalized additive models. Gu (2013) extended this into smoothing-spline ANOVA frameworks for flexible regression estimation. Their work demonstrates, that fANOVA not only provides a theoretical decomposition, but also serves as a basis for widely-used non-parametric statistical models featuring additive structure and controlled interactions.

Perhaps the most well-known application of fANOVA is in variance-based sensitivity analysis. Sobol’s original decomposition led directly to a variance decomposition, on which Sobol indices are based. Work from Owen (2013, 2014) modernized this framework, introducing efficient estimation strategies and generalized indices suited to quasi-Monte Carlo methods. Borgonovo et al. (2022) further advanced the field with mixture-based generalizations of fANOVA for uncertainty quantification.

Classical fANOVA requires independent input variables, which is a strong assumption in many real-world applications. Therefore, a stream of literature is concerned with the generalization of fANOVA to dependent variables. While Hooker (2007) was the first to present a generalized fANOVA framework, many other researchers were inspired by his work to create modifications of this (Rahman, 2014, Chastaing et al., 2012, El Idrissi et al., 2025). We see the generalization as central part of the basis of the fANOVA decomposition and therefore will also present it in this thesis.

A recent cluster of literature studies fANOVA for model interpretability. There is work of Lengerich et al. (2020), König et al. (2024), Choi et al. (2025) that all enhance in-

interpretability by using fANOVA to identify and disentangle variable interactions. Then there is work done in the explicit context of IML, where fANOVA can be used as a model-agnostic tool (Hooker, 2004, Fumagalli et al., 2025) or as foundational principle to build inherently interpretable models (Hu et al., 2025). fANOVA-based interpretability methods is probably the most novel field of fANOVA in which research is actively ongoing. Finally, there are specific domains of statistics, such as geostatistics, where fANOVA-based Kriging models are designed (Muehlenstaedt et al., 2012) or complex functions arising in computational finance are studied (Liu and Owen, 2006).

### 3 Formalization of fANOVA

The formal setup is based on Rahman (2014) and Chastaing et al. (2012).

Let  $\mathbb{N}, \mathbb{N}_0, \mathbb{R}$ , and  $\mathbb{R}_0^+$  denote the sets of positive integer (natural), nonnegative integer, real, and nonnegative real numbers, respectively. Throughout this thesis, we represent the  $k$ -dimensional Euclidean space by  $\mathbb{R}^k$  and the set of all  $k \times k$  real-valued matrices by  $\mathbb{R}^{k \times k}$ .

Let  $(\Omega, \mathcal{F}, \nu)$  be a measure space, where  $\Omega$  is a sample space,  $\mathcal{F}$  is a  $\sigma$ -algebra on  $\Omega$  and  $\nu : \mathcal{F} \rightarrow [0, 1]$  is a probability measure.  $\mathcal{B}^N$  is the Borel  $\sigma$ -algebra on  $\mathbb{R}^N$ ,  $N \in \mathbb{N}$ .  $\mathbf{X} = (X_1, \dots, X_N) : (\Omega, \mathcal{F}) \rightarrow (\mathbb{R}^N, \mathcal{B}^N)$  denotes an  $\mathbb{R}^N$ -valued random vector.

We assume that the probability distribution of  $\mathbf{X}$  is continuous and completely defined by the joint probability density function  $f_{\mathbf{X}} : \mathbb{R}^N \rightarrow \mathbb{R}_0^+$ .

Let  $u$  denote a subset of  $\{1, \dots, N\}$  with the complementary set  $-u := \{1, \dots, N\} \setminus u$  and cardinality  $0 \leq |u| \leq N$ . We denote strict inclusion of a subset by  $\subsetneq$  and  $\subseteq$  allows for equality.  $\mathbf{X}_u = (X_{i_1}, \dots, X_{i_{|u|}}), u \neq \emptyset, 1 \leq i_1 < \dots < i_{|u|} \leq N$  is a sub-vector of  $\mathbf{X}$  and  $\mathbf{X}_{-u} = \mathbf{X}_{\{1, \dots, N\} \setminus u}$  is the complementary subvector.

The marginal density function of  $\mathbf{X}_u$  is  $f_u(\mathbf{x}_u) := \int_{\mathbb{R}^{N-|u|}} f_{\mathbf{X}}(\mathbf{x}) d\nu(\mathbf{x}_{-u})$  for a given set  $\emptyset \neq u \subseteq \{1, \dots, N\}$ .

Let  $y(\mathbf{X}) := y(X_1, \dots, X_N)$  be a real-valued, measurable transformation on  $(\Omega, \mathcal{F})$ , which represents a probabilistic model with random variables as inputs. The Hilbert space of square-integrable functions  $y$  with respect to the induced generic measure  $f_{\mathbf{X}}(\mathbf{x}) d\nu(\mathbf{x})$  supported on  $\mathbb{R}^N$  is given by:

$$\mathcal{L}^2(\Omega, \mathcal{F}, \nu) = \{y : \Omega \rightarrow \mathbb{R} \text{ s.t. } \mathbb{E}[y^2(\mathbf{X})] < \infty\}.$$

The inner product defined as:

$$\langle y, g \rangle = \int_{\mathbb{R}^N} y(\mathbf{x}) g(\mathbf{x}) f_{\mathbf{X}} d\nu(\mathbf{x}) = \mathbb{E}[y(\mathbf{X}) g(\mathbf{X})], \quad \forall y, g \in \mathcal{L}^2.$$

The norm, denoted as  $\|\cdot\|$ , is defined by:

$$\|y\| = \sqrt{\langle y, y \rangle} = \sqrt{\int_{\mathbb{R}^N} y^2(\mathbf{x}) d\nu(\mathbf{x})} = \sqrt{\mathbb{E}[y(\mathbf{X})^2]}, \quad \forall y \in \mathcal{L}^2.$$

We start by defining the fANOVA decomposition in a general form, which is independent of distribution assumptions about the input variables or anything of the sort. The decomposition consists of  $2^N$  terms and its specific form is determined by the assumptions about the input variables and integration measure.



**Definition 3.1.** Let  $y$  denote a mathematical model with input vector  $\mathbf{X} := (X_1, \dots, X_N)$ . The functional ANOVA decomposition of  $y$  takes the form:

$$y(\mathbf{X}) = \sum_{u \subseteq \{1, \dots, N\}} y_u(\mathbf{X}_u). \quad (1)$$

### 3.1 Classical fANOVA

For his original fANOVA decomposition, Sobol only considered function defined on the unit hypercube, but later work shows that it is no problem to work within the measure space  $(\mathbb{R}^N, \mathcal{B}^N, \nu)$ . In any case, we assume that the coordinates  $X_1, \dots, X_N$  are independent of each other. Under independence, we work with a product-type probability measure of  $\mathbf{X}$  given by  $f_{\mathbf{X}}(\mathbf{x}) d\nu(\mathbf{x}) = \prod_{i=1}^N f_{\{i\}}(x_i) d\nu(x_i)$ , where  $f_{\{i\}} : \mathbb{R} \rightarrow \mathbb{R}_0^+$  is the marginal probability density function of  $X_i$  defined on  $(\Omega_i, \mathcal{F}_i, \nu_i)$  with a bounded or an unbounded support on  $\mathbb{R}$ .

Given this setup, we formulate a condition, proposed by Rahman (2014), which we would like to hold for the fANOVA terms to be well-defined and interpretable.

**Condition 3.1** (Strong annihilating conditions, (Rahman, 2014)). *For the classical fANOVA decomposition we require, that all the nonconstant component functions  $y_u$  integrate to zero w.r.t. the marginal probability density of each random variable in  $u$ , i.e.*

$$\int_{\mathbb{R}} y_u(\mathbf{x}_u) f_{\{i\}}(x_i) d\nu(x_i) = 0 \quad \text{for } i \in u \neq \emptyset. \quad (2)$$

**Proposition 3.1.** *Given the strong annihilating conditions, the nonconstant fANOVA components are centered around zero. This means for all  $\emptyset \neq u \subseteq \{1, \dots, N\}$  it holds that:*

$$\int_{\mathbb{R}^N} y_u(\mathbf{x}_u) f_{\mathbf{X}}(\mathbf{x}) d\nu(\mathbf{x}) := \mathbb{E}[y_u(\mathbf{X}_u)] = 0. \quad (3)$$

*Proof.*

$$\begin{aligned} \mathbb{E}[y_u(\mathbf{X}_u)] &:= \int_{\mathbb{R}^N} y_u(\mathbf{x}_u) f_{\mathbf{X}}(\mathbf{x}) d\nu(\mathbf{x}) \\ &= \int_{\mathbb{R}^{|u|}} y_u(\mathbf{x}_u) f_{\mathbf{X}_u}(\mathbf{x}_u) d\nu(\mathbf{x}_u) \\ &= \int_{\mathbb{R}^{|u|}} y_u(\mathbf{x}_u) \prod_{j \in u} f_{\{j\}}(x_j) d\nu(\mathbf{x}_u) \\ &= \int_{\mathbb{R}^{|u|-1}} \int_{\mathbb{R}} y_u(\mathbf{x}_u) f_{\{i\}}(x_i) d\nu(x_i) \prod_{j \in u, j \neq i} f_{\{j\}}(x_j) d\nu(x_{u \setminus \{i\}}) = 0 \end{aligned}$$

□

**Proposition 3.2.** *Given the strong annihilating conditions, the fANOVA terms are orthogonal to each other. If two sets of indices are not completely equivalent, i.e.  $\emptyset \neq u \subseteq \{1, \dots, N\}$ ,  $\emptyset \neq v \subseteq \{1, \dots, N\}$ , and  $u \neq v$ , then it holds that:*

$$\int_{\mathbb{R}^N} y_u(\mathbf{x}_u) y_v(\mathbf{x}_v) f_{\mathbf{X}}(\mathbf{x}) d\nu(\mathbf{x}) = \mathbb{E}[y_u(\mathbf{X}_u) y_v(\mathbf{X}_v)] = 0. \quad (4)$$

*Proof.* Since  $u \neq v$ , there exists at least one index contained in exactly one of the sets. Without loss of generality, we pick  $i \in u \setminus v$ . Then  $y_v(\mathbf{x}_v)$  is independent of  $x_i$ , and by the strong annihilating conditions

$$\int_{\mathbb{R}} y_u(\mathbf{x}_u) f_{\{i\}}(x_i) d\nu(x_i) = 0 \quad \text{for all fixed } \mathbf{x}_{u \setminus \{i\}}.$$

Hence,

$$\begin{aligned} \mathbb{E}[y_u(\mathbf{X}_u) y_v(\mathbf{X}_v)] &= \int_{\mathbb{R}^N} y_u(\mathbf{x}_u) y_v(\mathbf{x}_v) f_{\mathbf{X}}(\mathbf{x}) d\nu(\mathbf{x}) \\ &= \int_{\mathbb{R}^N} y_u(\mathbf{x}_u) y_v(\mathbf{x}_v) \prod_{j=1}^N f_{\{j\}}(x_j) d\nu(\mathbf{x}) \\ &= \int_{\mathbb{R}^{N-1}} \left( \int_{\mathbb{R}} y_u(\mathbf{x}_u) f_{\{i\}}(x_i) d\nu(x_i) \right) y_v(\mathbf{x}_v) \prod_{j \neq i} f_{\{j\}}(x_j) d\nu(\mathbf{x}_{-i}) \\ &= 0. \end{aligned}$$

□

As we have seen, the fANOVA terms are “fully orthogonal” to each other, meaning not only terms of different order are orthogonal to each other but also terms of the same order are. Zero-mean and orthogonality are desirable and important properties because they ensure that a fANOVA term can be interpreted as isolated effect of the specific variable or isolated effect of an interaction. The term  $y_{\{1\}}$ , for example, captures the isolated main effects of  $X_1$ ; there is no other effect mixed into it, which  $X_1$  might have through interactions with other variables. At the same time, the term  $y_{\{1,2\}}$  captures the interaction effect of  $X_1$  and  $X_2$ , while the solo effect of  $X_1$  is already captured by  $y_{\{1\}}$  and does not merge into  $y_{\{1,2\}}$ . From the lens of interpretability, this distinguishes the fANOVA decomposition from methods such as partial dependence (PD) or Shapley values.

### 3.1.1 Construction of the fANOVA Terms

The individual fANOVA terms for the variables with indices in  $u$  are constructed by integrating the original function  $y(\mathbf{X})$  w.r.t. all variables except for the ones in  $u$ , and subtracting the lower order terms. Intuitively the integral is averaging the original function over all other variables except the ones of interest, which leaves us with a function of the variables of interest only. Subtracting lower order terms corresponds to accounting for effects that are already explained by other variables or interactions so that we obtain the isolated effects.

Since  $u = \emptyset$  for the constant term, we integrate w.r.t. all variables:

$$y_{\emptyset} = \int_{\mathbb{R}^N} y(\mathbf{x}) \prod_{i=1}^N f_{\{i\}}(x_i) d\nu(x_i) = \mathbb{E}[y(\mathbf{X})]. \quad (5)$$

For all other effects  $\emptyset \neq u \in \{1, \dots, N\}$  we can calculate:

$$y_u(\mathbf{X}_u) = \int_{\mathbb{R}^{N-|u|}} y(\mathbf{X}_u, \mathbf{x}_{-u}) \prod_{i=1, i \notin u}^N f_{\{i\}}(x_i) d\nu(x_i) - \sum_{v \subsetneq u} y_v(\mathbf{X}_v). \quad (6)$$

Notice that this definition relies on a product-type measure rooted in the independence assumption. We will see what changes when we let go of this assumption in the second part of this section.

As suggested earlier, the fANOVA components offer a clear interpretation of the model, decomposing it into main effects, two-way interaction effects, and so on. This is why fANOVA decomposition has received increasing attention in the IML literature, holding the potential for a global model-agnostic explanation method of black box models.

### 3.1.2 Example: Independent Multivariate Normal Inputs

Throughout this thesis, we will use the following simple function as a running example:

$$h(x_1, x_2) = a + x_1 + 2x_2 + x_1x_2, \quad (7)$$

which contains both main effects and an interaction term.

We assume the input vector

$$\mathbf{X} = (X_1, X_2)^T$$

follows a bivariate standard normal distribution

$$\mathbf{X} \sim \mathcal{N}\left(\begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 & \rho \\ \rho & 1 \end{pmatrix}\right).$$

This general formulation includes both independent inputs ( $\rho = 0$ ) and correlated inputs ( $\rho \neq 0$ ).

From properties of the multivariate normal distribution, the marginal distributions are

$$X_1 \sim \mathcal{N}(0, 1), \quad X_2 \sim \mathcal{N}(0, 1),$$

and the conditional distributions are given by

$$X_1 \mid X_2 = x_2 \sim \mathcal{N}(\rho x_2, 1 - \rho^2), \quad X_2 \mid X_1 = x_1 \sim \mathcal{N}(\rho x_1, 1 - \rho^2).$$

This example will allow us to compute and compare the classical and generalized fANOVA decompositions for different correlation structures.

The classical fANOVA decomposition we covered so far assumes independence, i.e.,  $\rho = 0$ . Here,  $X_1$  and  $X_2$  are independent and standard normal, so the conditional means vanish, and the classical fANOVA decomposition simplifies considerably. Computing the constant component via expectation gives:

$$\begin{aligned} h_\emptyset &= \mathbb{E}[h(X_1, X_2)] \\ &= \mathbb{E}[a + X_1 + 2X_2 + X_1X_2] \\ &= a + \mathbb{E}[X_1] + 2\mathbb{E}[X_2] + \mathbb{E}[X_1]\mathbb{E}[X_2] = a. \end{aligned}$$

Under zero-mean constraint and independence, the main effects and the interaction effect can be computed as follows:

$$\begin{aligned} h_{\{1\}}(x_1) &= \mathbb{E}_{X_2}[h(x_1, X_2)] - h_\emptyset \\ &= \mathbb{E}_{X_2}[a + x_1 + 2X_2 + x_1X_2] - a \\ &= x_1 + 2\mathbb{E}_{X_2}[X_2] + x_1\mathbb{E}_{X_2}[X_2] = x_1, \\ h_{\{2\}}(x_2) &= \mathbb{E}_{X_1}[h(X_1, x_2)] - h_\emptyset \\ &= \mathbb{E}_{X_1}[a + X_1 + 2x_2 + X_1x_2] - a \\ &= \mathbb{E}_{X_1}[X_1] + 2x_2 + x_2\mathbb{E}_{X_1}[X_1] = 2x_2, \\ h_{\{1,2\}}(x_1, x_2) &= \mathbb{E}[h(x_1, x_2)] - h_\emptyset - h_{\{1\}}(x_1) - h_{\{2\}}(x_2) \\ &= a + x_1 + 2x_2 + x_1x_2 - a - x_1 - 2x_2 = x_1x_2. \end{aligned}$$

It comes as no surprise that in this simple case the fANOVA decomposition does not provide any additional insights, as the isolated effects can be directly seen from the function. We show this simple example nevertheless to illustrate at which step which assumption is used. This will make clearer what breaks down when we generalize to dependent variables.

### 3.1.3 Equality to Hoeffding Decomposition

As we mentioned in section 2 the Hoeffding decomposition (Hoeffding, 1948) laid the groundwork for the fANOVA decomposition. Here, we want to point out that both decompositions yield the same component functions under the assumption of independent and zero-centered inputs. Though we provide no formal proof, we want to illustrate this with our running example from Equation 7.

**Definition 3.2** (Hoeffding decomposition). *Let  $y$  denote a real-valued function on  $\mathbb{R}^N$  with independent inputs  $X_1, \dots, X_N$ . The Hoeffding decomposition of  $y$  takes the form (Il Idrissi et al., 2025):*

$$y(\mathbf{X}) = \sum_{A \subseteq D} y_A(\mathbf{X}_A), \quad D := \{1, \dots, N\}, \quad (8)$$

where, for each  $A \subseteq D$ , the component function  $y_A$  is defined by

$$y_A(\mathbf{X}_A) = \sum_{B \subseteq A} (-1)^{|A|-|B|} \mathbb{E}[y(\mathbf{X}) | \mathbf{X}_B]. \quad (9)$$

We can now apply the Hoeffding decomposition to our running example from Equation 7. For  $A = \emptyset$  there is only one subset  $B = \emptyset$ . Substituting this into Definition 3.2 we obtain:

$$h_{\emptyset} = (-1)^{0-0} \mathbb{E}[h(\mathbf{X})] = \mathbb{E}[h(X_1, X_2)].$$

We compute

$$\mathbb{E}[h(X_1, X_2)] = a + \mathbb{E}[X_1] + 2\mathbb{E}[X_2] + \mathbb{E}[X_1]\mathbb{E}[X_2] = a + \mu_1 + 2\mu_2 + \mu_1\mu_2.$$

Next, the subsets of  $A = \{1\}$  are  $B = \emptyset$  and  $B = \{1\}$ , so

$$h_{\{1\}}(X_1) = (-1)^{1-0} \mathbb{E}[h(\mathbf{X})] + (-1)^{1-1} \mathbb{E}[h(\mathbf{X}) | X_1] = -\mathbb{E}[h] + \mathbb{E}[h(\mathbf{X}) | X_1].$$

Since  $X_1$  is independent of  $X_2$ , the conditional expectation is:

$$\mathbb{E}[h(\mathbf{X}) | X_1] = a + X_1 + 2\mu_2 + X_1\mu_2,$$

so the final expression is given by:

$$h_{\{1\}}(X_1) = -(a + \mu_1 + 2\mu_2 + \mu_1\mu_2) + (a + X_1 + 2\mu_2 + X_1\mu_2) = (1 + \mu_2)(X_1 - \mu_1).$$

The subsets of  $A = \{2\}$  are  $B = \emptyset$  and  $B = \{2\}$ , so

$$h_{\{2\}}(X_2) = (-1)^{1-0} \mathbb{E}[h(\mathbf{X})] + (-1)^{1-1} \mathbb{E}[h(\mathbf{X})|X_2] = -\mathbb{E}[h(\mathbf{X})] + \mathbb{E}[h(\mathbf{X})|X_2].$$

Under independence the conditional expectation is:

$$\mathbb{E}[h(\mathbf{X})|X_2] = a + \mu_1 + 2X_2 + \mu_1X_2,$$

which yields the final expression:

$$h_{\{2\}}(X_2) = -(a + \mu_1 + 2\mu_2 + \mu_1\mu_2) + (a + \mu_1 + 2X_2 + \mu_1X_2) = (2 + \mu_1)(X_2 - \mu_2).$$

Finally, the subsets of  $A = \{1, 2\}$  are  $B = \emptyset$ ,  $B = \{1\}$ ,  $B = \{2\}$ ,  $B = \{1, 2\}$ . We obtain:

$$\begin{aligned} h_{\{1,2\}}(X_1, X_2) &= (-1)^{2-0} \mathbb{E}[h(\mathbf{X})] + (-1)^{2-1} \mathbb{E}[h(\mathbf{X})|X_1] + (-1)^{2-1} \mathbb{E}[h(\mathbf{X})|X_2] + (-1)^{2-2} \mathbb{E}[h(\mathbf{X})|X_1, X_2] \\ &= \mathbb{E}[h(\mathbf{X})] - \mathbb{E}[h(\mathbf{X})|X_1] - \mathbb{E}[h(\mathbf{X})|X_2] + \mathbb{E}[h(\mathbf{X})|X_1, X_2]. \end{aligned}$$

We already know:

$$\mathbb{E}[h(\mathbf{X})|X_1, X_2] = h(X_1, X_2) = a + X_1 + 2X_2 + X_1X_2.$$

Thus:

$$\begin{aligned} h_{\{1,2\}}(X_1, X_2) &= (a + \mu_1 + 2\mu_2 + \mu_1\mu_2) - (a + X_1 + 2\mu_2 + \mu_2X_1) - (a + \mu_1 + 2X_2 + \mu_1X_2) \\ &\quad + (a + X_1 + 2X_2 + X_1X_2) \\ &= X_1X_2 - \mu_2X_1 - \mu_1X_2 + \mu_1\mu_2 \\ &= (X_1 - \mu_1)(X_2 - \mu_2). \end{aligned}$$

Bringing everything together, the Hoeffding decomposition of  $h(X_1, X_2)$  with general means  $\mu_1 = \mathbb{E}[X_1]$  and  $\mu_2 = \mathbb{E}[X_2]$  is:

$$h(X_1, X_2) = h_\emptyset + h_1(X_1) + h_2(X_2) + h_{12}(X_1, X_2),$$

with

$$\begin{aligned} h_{\emptyset} &= a + \mu_1 + 2\mu_2 + \mu_1\mu_2, \\ h_1(X_1) &= (1 + \mu_2)(X_1 - \mu_1), \\ h_2(X_2) &= (2 + \mu_1)(X_2 - \mu_2), \\ h_{\{1,2\}}(X_1, X_2) &= (X_1 - \mu_1)(X_2 - \mu_2). \end{aligned}$$

Under the special case of centered input variables, so  $\mathbb{E}[X_1] = \mathbb{E}[X_2] = 0$ , the decomposition simplifies to:

$$h_{\emptyset} = a, \quad h_{\{1\}}(X_1) = X_1, \quad h_{\{2\}}(X_2) = 2X_2, \quad h_{\{1,2\}}(X_1, X_2) = X_1X_2,$$

which coincides with the fANOVA terms calculated earlier. The principle of the Hoeffding decomposition is therefore the same as for the fANOVA decomposition, but the fANOVA decomposition is stated in a recursive form, which makes it clear that each effect accounts for the contribution of lower order terms. In addition, the fANOVA terms are themselves zero-centered by construction.

### 3.1.4 fANOVA via Projection

In the following we revisit the fANOVA decomposition from the view of orthogonal projections. For this section the parallel between the (conditional) expected value and orthogonal projections formulated in Van der Vaart (1998) is crucial. Having this perspective on the fANOVA decomposition helps in bridging different notations of the method (e.g. via expected value or via integral) and also supports in understanding the generalization of fANOVA later in this section. First we define generally what an orthogonal projection is, and then we will use the idea in the context of fANOVA.

**Definition 3.3** (Orthogonal Projection). *Let  $\mathcal{G} \subset \mathcal{L}^2$  denote a linear subspace. The projection of  $y$  onto  $\mathcal{G}$  is defined by the function  $\Pi_{\mathcal{G}}y$  which minimizes the distance to  $y$  in  $\mathcal{L}^2$  (Nagler, 2024a):*

$$\Pi_{\mathcal{G}}y = \arg \min_{g \in \mathcal{G}} \|y - g\|^2 = \arg \min_{g \in \mathcal{G}} \mathbb{E}[(y(\mathbf{X}) - g(\mathbf{X}))^2]. \quad (10)$$

When we define the constant term  $y_{\emptyset}$  our goal is to best approximate the original function  $y$  by a constant function. In other words, we want to minimize the squared difference between  $y$  and a constant function  $g_0(x) = a$  over all possible constant functions. The solution is the orthogonal projection of  $y$  onto the linear subspace of all constant functions  $\mathcal{G}_0 = \{g(x) = a; a \in \mathbb{R}\}$ .

In a probabilistic context, we want to minimize the expected squared different between the random variables  $y(\mathbf{X})$  and  $a$ , which turns out to be equivalent to the expected value of the random variable (Van der Vaart, 1998). So intuitively, in the absence of any additional information, the expected value is our best approximation of  $y$ . More formally we can write for the constant term  $y_\emptyset$ :

$$\begin{aligned}\Pi_{\mathcal{G}_0} y &= \arg \min_{g_0 \in \mathcal{G}_0} \|y - g_0\|^2 \\ &= \arg \min_{a_0 \in \mathbb{R}} \mathbb{E}[(y(\mathbf{X}) - a)^2] \\ &= \mathbb{E}[y(\mathbf{X})] = y_\emptyset\end{aligned}$$

The main effect  $y_{\{i\}}(x_i)$  is the projection of  $y$  onto the subspace of all functions that only depend on  $x_i$ , i.e.  $\mathcal{G}_i = \{g(x) = g_{\{i\}}(x_i)\}$ . There is no need for additional constraints since subtracting lower order terms ensures that orthogonality and zero-mean are fulfilled. The conditional expected value of  $\mathbb{E}[y(\mathbf{X}) \mid X_i = x_i]$  is the solution to the minimization problem (Van der Vaart, 1998), and the conditional expected value is also a way to express the fANOVA terms (Muehlenstaedt et al., 2012). Thus, for the main effect  $y_{\{i\}}(x_i)$  we can write:

$$\begin{aligned}(\Pi_{\mathcal{G}_i} y)(\cdot) - y_\emptyset &= \arg \min_{g_i \in \mathcal{G}_i} \|y - g_{\{i\}}\|^2 - y_\emptyset \\ &= \arg \min_{g_i \in \mathcal{G}_i} \mathbb{E}[(y(\mathbf{X}) - g_{\{i\}}(X_i))^2] - y_\emptyset \\ &= \mathbb{E}[y(\mathbf{X}) \mid X_i = \cdot] - y_\emptyset = y_{\{i\}}(\cdot)\end{aligned}$$

The two-way interaction effect  $y_{\{i,j\}}(\cdot, \cdot)$  is the projection of  $y$  onto the subspace of all functions that depend on  $x_i$  and  $x_j$ . i.e.  $\mathcal{G}_{i,j} = \{g(x) = g_{\{i,j\}}(x_i, x_j)\}$ . Again, we account for lower-order effects by subtracting the constant term and all main effects:

$$\begin{aligned}(\Pi_{\mathcal{G}_{i,j}} y)(\cdot, \cdot) - (y_\emptyset + y_{\{i\}}(\cdot) + y_{\{j\}}(\cdot)) &= \arg \min_{g_{\{i,j\}} \in \mathcal{G}_{i,j}} \|y - g_{\{i,j\}}\|^2 - (y_\emptyset + y_{\{i\}}(\cdot) + y_{\{j\}}(\cdot)) \\ &= \arg \min_{g_{\{i,j\}} \in \mathcal{G}_{i,j}} \mathbb{E}[(y(\mathbf{X}) - g_{\{i,j\}}(\cdot, \cdot))^2] - (y_\emptyset + y_{\{i\}}(\cdot) + y_{\{j\}}(\cdot)) \\ &= \mathbb{E}[y(\mathbf{X}) \mid X_j = \cdot, X_i = \cdot] - (y_\emptyset + y_{\{i\}}(\cdot) + y_{\{j\}}(\cdot)) = y_{\{i,j\}}(\cdot, \cdot)\end{aligned}$$

In general, we can write for a subset of indices  $u \subseteq \{1, \dots, N\}$  and the subspace  $\mathcal{G}_u =$



$\{g(\mathbf{x}) = g_u(\mathbf{x}_u)\}$ :

$$\begin{aligned} (\Pi_{\mathcal{G}_u} y)(\cdot) - \sum_{v \subsetneq u} y_v(\cdot) &= \arg \min_{g_u \in \mathcal{G}_u} \|y - g_u\|^2 - \sum_{v \subsetneq u} y_v(\cdot) \\ &= \arg \min_{g_u \in \mathcal{G}_u} \mathbb{E}[(y(\mathbf{X}) - g_u(\cdot))^2] - \sum_{v \subsetneq u} y_v(\cdot) \\ &= \mathbb{E}[y(\mathbf{X}) | X_u = \cdot] - \sum_{v \subsetneq u} y_v(x) = y_u(\cdot), \end{aligned}$$

which means that we project  $y$  onto the subspace spanned by the own terms of the fANOVA component to be defined, while accounting for all lower-order terms.

On this note, we want to highlight that instead of subtracting the lower order terms from the projection, it is just as valid to first subtract lower order terms and project  $y$  on what is left. We can find both formulations in the literature. For example, Muehlenstaedt et al. (2012) subtracts from the projection and defines:

$$\begin{aligned} y_u(\mathbf{x}_u) &:= \mathbb{E}[y(\mathbf{X}) | \mathbf{X}_u = \mathbf{x}_u] - \sum_{v \subsetneq u} y_v(\mathbf{x}) \\ &= \int_{\mathbb{R}^{N-|u|}} y(\mathbf{x}) d\nu(\mathbf{x}_{-u}) - \sum_{v \subsetneq u} y_v(\mathbf{x}). \end{aligned}$$

Hooker (2004) takes the alternative view and defines the fANOVA components via the integral, which can be rewritten as the expected value:

$$\begin{aligned} y_u(\mathbf{x}_u) &:= \int_{\mathbb{R}^{N-|u|}} (y(\mathbf{x}) - \sum_{v \subsetneq u} y_v(\mathbf{x})) d\nu(\mathbf{x}_{-u}) \\ &= \mathbb{E}[y(\mathbf{X}) - \sum_{v \subsetneq u} y_v(\mathbf{x}) | \mathbf{X}_u = \mathbf{x}_u]. \end{aligned}$$

The first equivalence in each formulation is simply the definition in each original paper, while the second equivalence holds under the assumption of independent inputs.

### 3.1.5 Variance Decomposition

Studying the second moments of a function through the lens of the fANOVA decomposition can be useful, especially with regard to the construction of Sobol indices. This requires us to observe the second moment statistics of the decomposition. We already established that:

$$\mu := \mathbb{E}[y(\mathbf{X})] = y_\emptyset.$$

We can also compute the variance of  $y(\mathbf{X})$  via the fANOVA decomposition. We write the

sum over  $u$  for the sum over  $\emptyset \neq u \subseteq \{1, \dots, N\}$  and the sum over  $u \neq v$  for the sum over  $\emptyset \neq u \subseteq \{1, \dots, N\}, \emptyset \neq v \subseteq \{1, \dots, N\}, u \neq v$ . The variance of  $y$  is then given by:

$$\begin{aligned}
 \sigma^2 &:= \mathbb{E}[(y(\mathbf{X}) - \mu_G)^2] \\
 &= \mathbb{E}\left[\left(y_{\emptyset,G} + \sum_u y_{u,G}(\mathbf{X}_u) - y_{\emptyset,G}\right)^2\right] \\
 &= \mathbb{E}\left[\left(\sum_u y_{u,G}(\mathbf{X}_u)\right)^2\right] \\
 &= \sum_u \mathbb{E}[y_{u,G}^2(\mathbf{X}_u)] + 2\mathbb{E}\left[\sum_{u \neq v} y_u(\mathbf{X}_u)y_v(\mathbf{X}_v)\right] \\
 &= \sum_u \mathbb{E}[y_u^2(\mathbf{X}_u)],
 \end{aligned} \tag{11}$$

where all the cross-terms vanish due to the orthogonality of the fANOVA terms, i.e.  $\mathbb{E}[y_u(\mathbf{X}_u)y_v(\mathbf{X}_v)] = 0$  for  $u \neq v$ . This means that the variance of  $y(\mathbf{X})$  can be decomposed into the sum of the variances of the fANOVA terms.

We can verify that the variance decomposition holds for our example. This means, we verify that the following expression is true:

$$\text{Var}(h(X_1, X_2)) = \mathbb{E}[h_{\{1\}}^2(X_1)] + \mathbb{E}[h_{\{2\}}^2(X_2)] + \mathbb{E}[h_{\{1,2\}}^2(X_1, X_2)]$$

for the function

$$h(X_1, X_2) = a + X_1 + 2X_2 + X_1X_2,$$

where  $X_1, X_2$  are independent with zero-mean and unit variance.

We start with the left-hand side and compute the variance of  $h(X_1, X_2)$ .

$$\begin{aligned}
 \text{Var}(h(X_1, X_2)) &= \text{Var}(a + X_1 + 2X_2 + X_1X_2) \\
 &= \text{Var}(a) + \text{Var}(X_1) + 4\text{Var}(X_2) + \underbrace{\text{Var}(X_1X_2)}_{(*)} + 2\text{Cov}(X_1, 2X_2) \\
 &= 0 + 1 + 4 \cdot 1 + 1 \cdot 1 + 2 \cdot 0 = 6, \\
 (*) \quad \text{Var}(X_1X_2) &= \mathbb{E}[X_1^2X_2^2] - (\mathbb{E}[X_1X_2])^2 \\
 &= \mathbb{E}[X_1^2] \mathbb{E}[X_2^2] - (\mathbb{E}[X_1] \mathbb{E}[X_2])^2 \\
 &= (\text{Var}(X_1) + \mathbb{E}[X_1]^2) (\text{Var}(X_2) + \mathbb{E}[X_2]^2) - 0 \\
 &= 1 \cdot 1 = 1.
 \end{aligned}$$

For  $(\star)$  we used the fact that independence of any measurable map of  $X_1$  and  $X_2$  follows from independence of  $X_1$  and  $X_2$  for the second line, and the definition of the variance for the third line.

Next, we go in the other direction and start from the sum of the variances of the fANOVA terms. Their variances are given by:

$$\mathbb{E}[h_{\{1\}}^2(X_1)] = \mathbb{E}[X_1^2] = 1, \quad \mathbb{E}[h_{\{2\}}^2(X_2)] = \mathbb{E}[(2X_2)^2] = 4, \quad \mathbb{E}[h_{\{1,2\}}^2(X_1, X_2)] = \mathbb{E}[(X_1X_2)^2] = 1.$$

Therefore, summing everything up we obtain:

$$\mathbb{E}[h_{\{1\}}^2(X_1)] + \mathbb{E}[h_{\{2\}}^2(X_2)] + \mathbb{E}[h_{\{1,2\}}^2(X_1, X_2)] = 1 + 4 + 1 = 6 = \text{Var}(h(X_1, X_2)).$$

### 3.2 Generalized fANOVA

For the classical fANOVA we make the assumption of independent inputs, which is often violated in practice. In the remainder of this section, we therefore investigate what happens, when we allow for dependency between variables. First, let us test with our running example:

$$h(X_1, X_2) = a + X_1 + 2X_2 + X_1X_2,$$

Now  $\rho \neq 0$ , while keeping everything else the same. When we follow the same logic as in the classical case we obtain the following terms:

$$\begin{aligned} \tilde{h}_\emptyset &= \mathbb{E}[h(X_1, X_2)] = a + \mathbb{E}[X_1] + 2\mathbb{E}[X_2] + \mathbb{E}[X_1X_2] \\ &= a + \mathbb{E}[X_1X_2] = a + (\text{Cov}(X_1, X_2) + \mathbb{E}[X_1]\mathbb{E}[X_2]) \\ &= a + \rho \\ \tilde{h}_{\{1\}}(x_1) &= \mathbb{E}[h(X_1, X_2)|X_1 = x_1] - \tilde{h}_\emptyset \\ &= \mathbb{E}[a + X_1 + 2X_2 + X_1X_2|X_1 = x_1] - (a + \rho) \\ &= a + x_1 + 2\mathbb{E}[X_2|X_1 = x_1] + x_1\mathbb{E}[X_2|X_1 = x_1] - a - \rho \\ &= x_1 + \rho(2x_1 + x_1^2 - 1) \\ \tilde{h}_{\{2\}}(x_2) &= \mathbb{E}[h(X_1, X_2) | X_2 = x_2] - \tilde{h}_\emptyset \\ &= \mathbb{E}[a + X_1 + 2X_2 + X_1X_2 | X_2 = x_2] - (a + \rho) \\ &= a + 2x_2 + x_2\mathbb{E}[X_1 | X_2 = x_2] - a - \rho \\ &= 2x_2 + \rho(x_2 + x_2^2 - 1) \\ \tilde{h}_{\{1,2\}}(x_1, x_2) &= h(x_1, x_2) - \tilde{h}_\emptyset - \tilde{h}_{\{1\}}(x_1) - \tilde{h}_{\{2\}}(x_2) \\ &= a + x_1 + 2x_2 + x_1x_2 - (a + \rho) - (x_1 + \rho(2x_1 + x_1^2 - 1)) - (2x_2 + \rho(x_2 + x_2^2 - 1)) \\ &= x_1x_2 - 2\rho x_1 - \rho x_2 - \rho x_1^2 - \rho x_2^2 + \rho \end{aligned}$$

The fANOVA components are characterized by two central properties: zero-mean and orthogonality, which follow from ???. When we check if the components  $\tilde{h}_\emptyset, \tilde{h}_{\{1\}}, \tilde{h}_{\{2\}}, \tilde{h}_{\{1,2\}}$  satisfy these properties, we find out that all components are zero-centered, but not all are orthogonal to each other. We can, for example, immediately see that checking orthogonality between  $\tilde{h}_{\{1\}}, \tilde{h}_{\{1,2\}}$  will yield the expectation over the constant term  $\rho^2$  exactly once, meaning even if all the other expectations cancel out, this constant will remain and

the entire expression will be unequal to zero:

$$\begin{aligned}\mathbb{E}(\tilde{h}_{\{1\}}(X_1)\tilde{h}_{\{1,2\}}(X_1, X_2)) &= \mathbb{E}[(X_1 + 2\rho X_1 + \rho X_1^2 - \rho) \\ &\quad \cdot (X_1 X_2 - 2\rho X_1 - \rho X_2 - \rho X_1^2 - \rho X_2^2 + \rho)] \\ &= \mathbb{E}[X_1^2 X_2] \dots - \mathbb{E}[\rho^2] \neq 0.\end{aligned}$$

It turns out that naively computing the “fANOVA decomposition” under dependent inputs, results in components that lack orthogonality, which is a crucial property for interpretability. What we performed in this example is only a fANOVA-type decomposition, but not the true fANOVA decomposition for dependent inputs. This shows the need for a more involved approach for a generalization of this method.

### 3.2.1 Conditions for Generalized fANOVA

Stone (1994) inspired the pioneering work of Hooker (2007) who offers a first solution to the problem of dependent inputs in fANOVA. Work by Chastaing et al. (2012) and Rahman (2014) build on his framework with modifications and extensions.

The generalized fANOVA decomposition still follows the overarching form in Equation 1. However, we no longer work with a product-type probability measure but now  $f_{\mathbf{X}} : \mathbb{R} \rightarrow \mathbb{R}_0^+$  denotes an arbitrary probability density function and we now consider the marginal probability measure  $f_u(\mathbf{x}_u)d\nu(\mathbf{x}_u)$  supported on  $\mathbb{R}^{|u|}$ .

Instead of enforcing the strong annihilating conditions for desirable properties of the components, Rahman (2014) proposed to formulate a milder version. The milder version fulfills the same function as the strong annihilating conditions in the classical case but works with the joint density of the variables of interest, instead of the individual marginal probability density functions.

**Condition 3.2** (Weak annihilating conditions, (Rahman, 2014)). *For the generalized fANOVA decomposition we require, that all the nonconstant fANOVA terms of variables in  $u$  integrate to zero w.r.t. the joint probability density function of variables in  $u$ :*

$$\int_{\mathbb{R}} y_{u,G}(\mathbf{x}_u) f_{\mathbf{X}_u}(\mathbf{x}_u) d\nu(x_i) = 0 \quad \text{for } i \in u \neq \emptyset \quad (12)$$

If components are defined under the weak annihilating conditions, we can ensure that they have zero-mean and satisfy a milder form of orthogonality - hierarchical orthogonality, which means that components of different order are orthogonal to each other while components of the same order are not. Hierarchical orthogonality is the best we can do when independence cannot be assumed.

**Proposition 3.3.** *Given the weak annihilating conditions, the generalized fANOVA components  $y_{u,G}$ , with  $\emptyset \neq u \subseteq \{1, \dots, N\}$ , are centered around zero:*

$$\mathbb{E}[y_{u,G}(\mathbf{X}_u)] := \int_{\mathbb{R}^N} y_{u,G}(\mathbf{x}_u) f_{\mathbf{X}}(\mathbf{x}) d\nu(\mathbf{x}) = 0. \quad (13)$$

*Proof.* For any subset  $\emptyset \neq u \subseteq \{1, \dots, N\}$ , let  $i \in u$ . We assume that the weak annihilating conditions hold. Then

$$\begin{aligned} \mathbb{E}[y_{u,G}(\mathbf{X}_u)] &:= \int_{\mathbb{R}^N} y_{u,G}(\mathbf{x}_u) f_{\mathbf{X}}(\mathbf{x}) d\nu(\mathbf{x}) \\ &= \int_{\mathbb{R}^{|u|}} y_{u,G}(\mathbf{x}_u) \left( \int_{\mathbb{R}^{N-|u|}} f_{\mathbf{X}}(\mathbf{x}) d\nu(\mathbf{x}_{-u}) \right) d\nu(\mathbf{x}_u) \\ &= \int_{\mathbb{R}^{|u|}} y_{u,G}(\mathbf{x}_u) f_u(\mathbf{x}_u) d\nu(\mathbf{x}_u) \\ &= \int_{\mathbb{R}^{|u|-1}} \left( \int_{\mathbb{R}} y_{u,G}(\mathbf{x}_u) f_u(\mathbf{x}_u) d\nu(\mathbf{x}_i) \right) \prod_{j \in u, j \neq i} d\nu(\mathbf{x}_j) \\ &= 0, \end{aligned}$$

where we make use of Fubini's theorem and the last line follows from using the weak annihilating conditions (??).  $\square$

**Proposition 3.4.** *Given the weak annihilating conditions, the fANOVA components are hierarchically orthogonal. This means that for two components  $y_{u,G}$  and  $y_{v,G}$  with  $u \subsetneq v$ ,  $\emptyset \neq u \subseteq \{1, \dots, N\}$ ,  $\emptyset \neq v \subseteq \{1, \dots, N\}$  it holds that:*

$$\mathbb{E}[y_{u,G}(\mathbf{X}_u) y_{v,G}(\mathbf{X}_v)] := \int_{\mathbb{R}^N} y_{u,G}(\mathbf{x}_u) y_{v,G}(\mathbf{x}_v) f_{\mathbf{X}}(\mathbf{x}) d\nu(\mathbf{x}) = 0. \quad (14)$$

*Proof.* For any two subsets  $\emptyset \neq u \subseteq \{1, \dots, N\}$  and  $\emptyset \neq v \subseteq \{1, \dots, N\}$ , where  $v \subsetneq u$ ,

the subset  $u = v \cup (u \setminus v)$ . Let  $i \in (u \setminus v) \subseteq u$ . Then

$$\begin{aligned}
 \mathbb{E}[y_{u,G}(\mathbf{X}_u) y_{v,G}(\mathbf{X}_v)] &:= \int_{\mathbb{R}^N} y_{u,G}(\mathbf{x}_u) y_{v,G}(\mathbf{x}_v) f_{\mathbf{X}}(\mathbf{x}) d\nu(\mathbf{x}) \\
 &= \int_{\mathbb{R}^{|u|}} y_{u,G}(\mathbf{x}_u) y_{v,G}(\mathbf{x}_v) \left( \int_{\mathbb{R}^{N-|u|}} f_{\mathbf{X}}(\mathbf{x}) d\nu(\mathbf{x}_{-u}) \right) d\nu(\mathbf{x}_u) \\
 &= \int_{\mathbb{R}^{|u|}} y_{u,G}(\mathbf{x}_u) y_{v,G}(\mathbf{x}_v) f_u(\mathbf{x}_u) d\nu(\mathbf{x}_u) \\
 &= \int_{\mathbb{R}^{|v|}} y_{v,G}(\mathbf{x}_v) \int_{\mathbb{R}^{|u \setminus v|}} y_{u,G}(\mathbf{x}_u) f_u(\mathbf{x}_u) d\nu(\mathbf{x}_{u \setminus v}) d\nu(\mathbf{x}_v) \\
 &= \int_{\mathbb{R}^{|v|}} y_{v,G}(\mathbf{x}_v) \int_{\mathbb{R}^{|u \setminus v|-1}} \left( \int_{\mathbb{R}} y_{u,G}(\mathbf{x}_u) f_u(\mathbf{x}_u) d\nu(\mathbf{x}_i) \right) \prod_{\substack{j \in (u \setminus v) \\ j \neq i}} d\nu(\mathbf{x}_j) d\nu(\mathbf{x}_v) \\
 &= 0.
 \end{aligned}$$

Repeatedly using Fubini's theorem and the weak annihilating conditions the equality to zero follows.  $\square$

A key contribution from Hooker (2007) and Rahman (2014) is that they construct a generalization of the fANOVA decomposition method as a whole, not only parts, such as the Sobol indices. This means it is important that Rahman's generalized statements are coherent with the classical fANOVA decomposition.

**Proposition 3.5.** *The weak annihilating conditions become the strong annihilating conditions under independence assumption.*

*Proof.* Assume that the random variables  $\{X_j\}_{j \in u}$  are independent. Then we can factorize the marginal density  $f_u(\mathbf{x}_u)$  as

$$f_u(\mathbf{x}_u) = \prod_{j \in u} f_{\{j\}}(x_j).$$

Now consider the weak annihilating condition (??) for some  $i \in u \neq \emptyset$ :

$$\int_{\mathbb{R}} y_{u,G}(\mathbf{x}_u) f_u(\mathbf{x}_u) d\nu(\mathbf{x}_i) = 0.$$

Since we assume independence, we can substitute the joint marginal density with the product of the marginal densities:

$$\int_{\mathbb{R}} y_{u,G}(\mathbf{x}_u) \left( \prod_{j \in u} f_{\{j\}}(x_j) \right) d\nu(\mathbf{x}_i) = 0.$$

For fixed  $x_j$  with  $j \neq i$ , the terms  $f_{\{j\}}(x_j)$  are constant with respect to  $x_i$ , and can therefore be pulled out of the integral:

$$\left( \prod_{j \in u, j \neq i} f_{\{j\}}(x_j) \right) \int_{\mathbb{R}} y_{u,G}(\mathbf{x}_u) f_{\{i\}}(x_i) d\nu(x_i) = 0.$$

As product of probability density functions the prefactor is strictly positive for all  $x_j$  with  $j \neq i$ . Therefore, the integral must be zero for the equality to hold:

$$\int_{\mathbb{R}} y_{u,G}(\mathbf{x}_u) f_{\{i\}}(x_i) d\nu(x_i) = 0,$$

which are the strong annihilating conditions from ??.

□

### 3.2.2 Construction of the Generalized fANOVA Terms

Recall the construction of the classical fANOVA components (Equation 6). The equation tells us that the non-constant classical fANOVA components are defined via the integral of the original function w.r.t. to the product-type probability density function, minus effects by other terms. So ideally for a well-aligned generalization, we would want that the general fANOVA terms can be understood in a similar way, as the integral of  $y$  w.r.t. a *smartly chosen* probability density function, minus effects explained by other terms. This is exactly what Rahman (2014) accomplishes. To understand this, we first need to distinguish three cases of integration that will occur in the construction of the generalized components.

**Proposition 3.6.** *Consider the generalized fANOVA components  $y_{v,G}$ ,  $\emptyset \neq v \subseteq \{1, \dots, N\}$ , of a square-integrable function  $y : \mathbb{R}^N \rightarrow \mathbb{R}$ . When integrated w.r.t. the probability measure  $f_{-u}(\mathbf{x}_{-u}) d\nu(\mathbf{x}_{-u})$ ,  $u \subseteq \{1, \dots, N\}$ , one can distinguish between three cases:*

$$\begin{aligned} & \int_{\mathbb{R}^{N-|u|}} y_{v,G}(\mathbf{x}_v) f_{-u}(\mathbf{x}_{-u}) d\nu(\mathbf{x}_{-u}) \\ &= \begin{cases} \int_{\mathbb{R}^{|v \cap u|}} y_{v,G}(\mathbf{x}_v) f_{v \cap u}(\mathbf{x}_{v \cap u}) d\nu(\mathbf{x}_{v \cap u}) & \text{if } v \cap u \neq \emptyset \text{ and } v \not\subseteq u, \\ y_{v,G}(\mathbf{x}_v) & \text{if } v \cap u \neq \emptyset \text{ and } v \subseteq u, \\ 0 & \text{if } v \cap u = \emptyset. \end{cases} \end{aligned} \quad (15)$$

*Proof.* Let  $u \subseteq \{1, \dots, N\}$  and  $\emptyset \neq v \subseteq \{1, \dots, N\}$ . Rahman (2014) distinguishes between three types of relationships between  $v$  and  $u$ . Before analysing the first case, note



that for any such  $u$  and  $v$ , it is possible to write


$$(v \cap -u) \subseteq -u \quad \text{and} \quad -u = (-u \setminus (v \cap -u)) \cup (v \cap -u),$$

which will be used in the integral decomposition below.

**Case 1:**  $v \cap u \neq \emptyset$  and  $v \not\subseteq u$  Rahman (2014) uses the decomposition of  $-u$  stated above to decompose the integration over  $\mathbf{x}_{-u}$  as:

$$\begin{aligned} & \int_{\mathbb{R}^{N-|u|}} y_{v,G}(\mathbf{x}_v) f_{-u}(\mathbf{x}_{-u}) d\nu(\mathbf{x}_{-u}) \\ &= \int_{\mathbb{R}^{|v \cap -u|}} y_{v,G}(\mathbf{x}_v) \left( \int_{\mathbb{R}^{N-|u|-|v \cap -u|}} f_{-u}(\mathbf{x}_{-u \setminus (v \cap -u)}, \mathbf{x}_{v \cap -u}) d\nu(\mathbf{x}_{-u \setminus (v \cap -u)}) \right) d\nu(\mathbf{x}_{v \cap -u}) \\ &= \int_{\mathbb{R}^{|v \cap -u|}} y_{v,G}(\mathbf{x}_v) f_{v \cap -u}(\mathbf{x}_{v \cap -u}) d\mathbf{x}_{v \cap -u}, \end{aligned}$$


where the inner integral integrates out all variables in  $-u \setminus (v \cap -u)$ , resulting in the marginal density  $f_{v \cap -u}(\mathbf{x}_{v \cap -u})$ .



**Case 2:**  $v \cap u \neq \emptyset$  and  $v \subseteq u$ . Since the sets  $v$  and  $-u$  are then completely disjoint,  $y_{v,G}(\mathbf{x}_v)$  is independent of  $\mathbf{x}_{-u}$  and can be pulled out of the integral:

$$\int_{\mathbb{R}^{N-|u|}} y_{v,G}(\mathbf{x}_v) f_{-u}(\mathbf{x}_{-u}) d\nu(\mathbf{x}_{-u}) = y_{v,G}(\mathbf{x}_v) \int_{\mathbb{R}^{N-|u|}} f_{-u}(\mathbf{x}_{-u}) d\nu(\mathbf{x}_{-u}) = y_{v,G}(\mathbf{x}_v),$$


which works because  $f_{-u}$  integrates to one.



**Case 3:**  $v \cap u = \emptyset$ . In this case, we have  $v \subseteq -u$ , so  $v \cap -u = v$ . Thus, Rahman (2014) writes:

$$\begin{aligned}
 \int_{\mathbb{R}^{N-|u|}} y_{v,G}(\mathbf{x}_v) f_{-u}(\mathbf{x}_{-u}) d\nu(\mathbf{x}_{-u}) &= \int_{\mathbb{R}^{|v|}} y_{v,G}(\mathbf{x}_v) \left( \int_{\mathbb{R}^{N-|u|-|v|}} f_{-u}(\mathbf{x}_{-u}) d\nu(\mathbf{x}_{-u \setminus v}) \right) d\nu(\mathbf{x}_v) \\
 &= \int_{\mathbb{R}^{|v|}} y_{v,G}(\mathbf{x}_v) f_v(\mathbf{x}_v) d\nu(\mathbf{x}_v) \\
 &= \int_{\mathbb{R}^{|v|-1}} \left( \int_{\mathbb{R}} y_{v,G}(\mathbf{x}_v) f_v(\mathbf{x}_v) d\nu(x_i) \right) \prod_{\substack{j \in v \\ j \neq i}} d\nu(x_j) \\
 &= 0,
 \end{aligned}$$

while the integral is split in such a way that one recognizes the marginal density  $f_v$  and the zero-mean property from Condition 3.1 is used.



□

As we will see in the following, we will encounter all of these three integration cases from Proposition 3.6 in the definition of the generalized fANOVA components à la Rahman (2014). In Proposition 3.6 we also already see that the smartly chosen probability density function is  $f_{-u}(\mathbf{x}_{-u})$ .

**Proposition 3.7.** *The generalized fANOVA component functions  $y_{u,G}(\mathbf{x}_u)$ ,  $u \subseteq \{1, \dots, N\}$  of a square-integrable function  $y : \mathbb{R}^N \rightarrow \mathbb{R}$  for a given probability measure  $f_{\mathbf{X}}(\mathbf{x}) d\nu(\mathbf{x})$  of  $\mathbf{X} \in \mathbb{R}^N$  can be recursively defined via the following set of equations:*

$$y_{\emptyset,G} = \int_{\mathbb{R}^N} y(\mathbf{x}) f_{\mathbf{X}}(\mathbf{x}) d\nu(\mathbf{x}) \quad (16)$$

$$\begin{aligned}
 y_{u,G}(\mathbf{X}_u) &= \int_{\mathbb{R}^{N-|u|}} y(\mathbf{X}_u, \mathbf{x}_{-u}) f_{-u}(\mathbf{x}_{-u}) d\nu(\mathbf{x}_{-u}) - \sum_{v \subsetneq u} y_{v,G}(\mathbf{X}_v) \\
 &\quad - \sum_{\substack{\emptyset \neq v \subseteq \{1, \dots, N\} \\ v \cap u \neq \emptyset, v \not\subseteq u}} \int_{\mathbb{R}^{|v \cap -u|}} y_{v,G}(\mathbf{X}_{v \cap u}, \mathbf{x}_{v \cap -u}) f_{v \cap -u}(\mathbf{x}_{v \cap -u}) d\nu(\mathbf{x}_{v \cap -u}). \quad (17)
 \end{aligned}$$

*Proof.* Rahman (2014) begins by integrating both sides of the generalized fANOVA decomposition

$$y(\mathbf{x}) = \sum_{v \subseteq \{1, \dots, N\}} y_{v,G}(\mathbf{x}_v)$$

w.r.t.  $f_{-u}(\mathbf{x}_{-u}) d\nu(\mathbf{x}_{-u})$ , replacing  $\mathbf{X}$  by  $\mathbf{x}$ , and changing the dummy index from  $u$  to  $v$ . This yields:

$$\int_{\mathbb{R}^{N-|u|}} y(\mathbf{x}) f_{-u}(\mathbf{x}_{-u}) d\nu(\mathbf{x}_{-u}) = \sum_{v \subseteq \{1, \dots, N\}} \int_{\mathbb{R}^{N-|u|}} y_{v,G}(\mathbf{x}_v) f_{-u}(\mathbf{x}_{-u}) d\nu(\mathbf{x}_{-u}).$$

**Case:**  $u = \emptyset$ . We set  $u = \emptyset$ , so  $-u = \{1, \dots, N\}$  and  $f_{-u}(\mathbf{x}_{-u}) d\nu(\mathbf{x}_{-u}) = f_{\mathbf{X}}(\mathbf{x}) d\nu(\mathbf{x})$ . The above integral can then be written as:

$$\begin{aligned} \int_{\mathbb{R}^N} y(\mathbf{x}) f_{\mathbf{X}}(\mathbf{x}) d\nu(\mathbf{x}) &= \sum_{v \subseteq \{1, \dots, N\}} \int_{\mathbb{R}^N} y_{v,G}(\mathbf{x}_v) f_{\mathbf{X}}(\mathbf{x}) d\nu(\mathbf{x}) \\ &= y_{\emptyset,G} + \sum_{\emptyset \neq v \subseteq \{1, \dots, N\}} \int_{\mathbb{R}^N} y_{v,G}(\mathbf{x}_v) f_{\mathbf{X}}(\mathbf{x}) d\nu(\mathbf{x}) \\ &= y_{\emptyset,G} + \sum_{\emptyset \neq v \subseteq \{1, \dots, N\}} \mathbb{E}[y_{v,G}(\mathbf{X}_v)] = y_{\emptyset,G}, \end{aligned}$$

where the last sum vanishes under the weak annihilating condition.

**Case:**  $\emptyset \neq u \subseteq \{1, \dots, N\}$ . Returning to the integrated decomposition

$$\int_{\mathbb{R}^{N-|u|}} y(\mathbf{x}) f_{-u}(\mathbf{x}_{-u}) d\nu(\mathbf{x}_{-u}) = \sum_{v \subseteq \{1, \dots, N\}} \int_{\mathbb{R}^{N-|u|}} y_{v,G}(\mathbf{x}_v) f_{-u}(\mathbf{x}_{-u}) d\nu(\mathbf{x}_{-u}),$$

and applying Proposition 3.6 to evaluate each term in the sum according to the relationship between  $v$  and  $u$  yields four cases:

**(A)**  $v \cap u \neq \emptyset$  and  $v \not\subseteq u$ :

This corresponds to case 1 of Proposition 3.6. The integral becomes:

$$\sum_{\substack{\emptyset \neq v \subseteq \{1, \dots, N\} \\ v \cap u \neq \emptyset, v \not\subseteq u}} \int_{\mathbb{R}^{|v \cap u|}} y_{v,G}(\mathbf{x}_v) f_{v \cap u}(\mathbf{x}_{v \cap u}) d\nu(\mathbf{x}_{v \cap u}).$$

**(B)**  $v \subseteq u$ :

This is contained in case 2 of Proposition 3.6. The integrals reduce to the component

functions themselves:

$$\sum_{v \subsetneq u} y_{v,G}(\mathbf{x}_v).$$

(C)  $v = u$ :

This is also contained in case 2 of Proposition 3.6. The integral becomes:

$$y_{u,G}(\mathbf{x}_u).$$

(D)  $v \cap u = \emptyset$ :

This is case 3 of Proposition 3.6. These terms vanish:

$$\sum_{\substack{v \subseteq \{1, \dots, N\} \\ v \cap u = \emptyset}} 0 = 0.$$

Putting everything together:

$$\begin{aligned} & \int_{\mathbb{R}^{N-|u|}} y(\mathbf{x}) f_{-u}(\mathbf{x}_{-u}) d\nu(\mathbf{x}_{-u}) \\ &= y_{u,G}(\mathbf{x}_u) + \sum_{v \subsetneq u} y_{v,G}(\mathbf{x}_v) + \sum_{\substack{\emptyset \neq v \subseteq \{1, \dots, N\} \\ v \cap u \neq \emptyset, v \not\subset u}} \int_{\mathbb{R}^{|v \cap -u|}} y_{v,G}(\mathbf{x}_v) f_{v \cap -u}(\mathbf{x}_{v \cap -u}) d\nu(\mathbf{x}_{v \cap -u}). \end{aligned}$$

Rearranging gives the almost final expression for  $y_{u,G}(\mathbf{x}_u)$ :

$$\begin{aligned} y_{u,G}(\mathbf{x}_u) &= \int_{\mathbb{R}^{N-|u|}} y(\mathbf{x}) f_{-u}(\mathbf{x}_{-u}) d\nu(\mathbf{x}_{-u}) \\ &\quad - \sum_{v \subsetneq u} y_{v,G}(\mathbf{x}_v) \\ &\quad - \sum_{\substack{\emptyset \neq v \subseteq \{1, \dots, N\} \\ v \cap u \neq \emptyset, v \not\subset u}} \int_{\mathbb{R}^{|v \cap -u|}} y_{v,G}(\mathbf{x}_v) f_{v \cap -u}(\mathbf{x}_{v \cap -u}) d\nu(\mathbf{x}_{v \cap -u}). \end{aligned}$$

As a final step, Rahman (2014) writes  $v = (v \cap u) \cup (v \cap -u)$  to obtain the expression of Proposition 3.7.  $\square$

### 3.2.3 Generalization via Projection

Hooker (2007) approaches his generalization of the fANOVA decomposition from the angle of orthogonal projections. Instead of the more recursive definition of the components functions as in Rahman (2014), he defines the fANOVA components as a joint set which

simultaneously minimizes the squared difference to the original function  $y$  under certain constraints. The constraints he sets for the optimization problem should ensure that the generalized components satisfy the desired properties of zero-mean and hierarchical orthogonality. The generalized fANOVA terms  $\{y_u(x_u) | u \subseteq d\}$  jointly satisfy:

$$\{y_{u,G}(\mathbf{x}_u) | u \subseteq d\} = \arg \min_{\{g_u \in \mathcal{L}^2(\mathbb{R}^{|u|})\}} \int_{\mathbb{R}^N} \left( \sum_{u \subseteq d} g_u(\mathbf{x}_u) - y(\mathbf{x}) \right)^2 f_{\mathbf{X}}(\mathbf{x}) d\nu(\mathbf{x}) \quad (18)$$

under the hierarchical orthogonality conditions:

$$\forall v \subseteq u, \forall g_v : \int_{\mathbb{R}^N} y_u(\mathbf{x}_u) g_v(\mathbf{x}_v) f_{\mathbf{X}}(\mathbf{x}) d\nu(\mathbf{x}) = 0. \quad (19)$$

In Equation 18 we recognize a projection. We are simultaneously finding the set of components functions  $g_u$  that minimize the weighted squared difference to the original function  $y$  (under the given integral constraint), which is exactly the definition of a projection of  $y$  onto a specific subspace  $\mathcal{G}$  (Definition 3.3).

However, the constraint in Equation 19 is infeasible to enforce in practice. Therefore, Hooker formulated the following proposition, which ensures hierarchical orthogonality of the fANOVA components and thus forms the building block of his approach. It can be compared to Condition 3.2 in Rahman (2014).

**Proposition 3.8.** *The hierarchical orthogonality of the fANOVA components is ensured if and only if the following integral condition holds:*

$$\forall u \subseteq N, \forall i \in u : \int_{\mathbb{R}^N} y_u(\mathbf{x}_u) f_{\mathbf{X}}(\mathbf{x}) d\nu(x_i) d\nu(\mathbf{x}_{-u}) = 0. \quad (20)$$

*Proof.* The proof is organized in two parts. First, Hooker shows that, the hierarchical orthogonality is true, if the integral conditions hold. Second, he shows that hierarchical orthogonality breaks down if the integral conditions are not true.

For the first part, assume that Equation 3.8 holds. Let  $i \in u \setminus v$ , then  $y_v(\mathbf{x}_v)$  is independent of  $x_i$  and  $\mathbf{x}_{-u}$ , so we can write:

$$\begin{aligned} \langle y_u, y_v \rangle &:= \int_{\mathbb{R}^N} y_v(\mathbf{x}_v) y_u(\mathbf{x}_u) f_{\mathbf{X}}(\mathbf{x}) d\nu(\mathbf{x}) \\ &= y_v(\mathbf{x}_v) \int_{\mathbb{R}^N} y_u(\mathbf{x}_u) f_{\mathbf{X}}(\mathbf{x}) d\nu(\mathbf{x}) \\ &= 0. \end{aligned}$$

For the second part, assume that there exists a subset  $u$  and an index  $i$  for which hierarchical orthogonality does not hold, i.e.

$$\int_{\mathbb{R}^N} y_u(\mathbf{x}_u) f_{\mathbf{X}}(\mathbf{x}) d\nu(x_i) d\nu(\mathbf{x}_{-u}) \neq 0 \quad \text{for some } i, u.$$

Further, assume that hierarchical orthogonality holds for all subsets  $v \neq u$  and indices  $j \neq i$ . Hooker then constructs a fANOVA term  $y_v$  with lower order than  $y_u$ , which is not orthogonal to  $y_u$ . He sets  $v = u \setminus \{i\}$ , so  $y_v$  is one order lower than  $y_u$  and defined as:

$$y_v(\mathbf{x}_v) := \int_{\mathbb{R}^N} f_u(\mathbf{x}_u) f_{\mathbf{X}}(\mathbf{x}) d\nu(x_i) d\nu(\mathbf{x}_{-u}).$$

$y_v$  is a valid fANOVA component, which is unequal to zero by assumption of hierarchical orthogonality being false, while it itself satisfies hierarchical orthogonality by assumption:

$$\forall j \in v, \quad \int_{\mathbb{R}^N} y_v(\mathbf{x}_v) f_{\mathbf{X}}(\mathbf{x}) d\nu(x_j) d\nu(\mathbf{x}_{-v}) = 0.$$

Hooker then shows that  $y_v$  is not orthogonal to  $y_u$ :

$$\begin{aligned} \langle y_u, y_v \rangle &= \int_{\mathbb{R}^N} y_u(\mathbf{x}_u) y_v(\mathbf{x}_v) f_{\mathbf{X}}(\mathbf{x}) d\nu(\mathbf{x}) \\ &= \int_{\mathbb{R}^N} y_u(\mathbf{x}_u) \left( \int_{\mathbb{R}^N} y_u(\mathbf{x}_u) f_{\mathbf{X}}(\mathbf{x}) d\nu(x_i) d\nu(\mathbf{x}_{-u}) \right) f_{\mathbf{X}}(\mathbf{x}) d\nu(\mathbf{x}) \\ &= \int_{\mathbb{R}^N} \left( \int_{\mathbb{R}^N} y_u(\mathbf{x}_u) f_{\mathbf{X}}(\mathbf{x}) d\nu(x_i) d\nu(\mathbf{x}_{-u}) \right)^2 d\nu(\mathbf{x}_{u \setminus \{i\}}) \\ &\neq 0. \end{aligned}$$

□

Hooker approaches his generalization through the lens of projections while Rahman gives a form that tries to imitate the classical fANOVA terms. A crucial parallel of both versions which distinguishes them from the classical case is that their components are defined in dependence of each other (Proposition 3.7, ??). This makes it in general difficult to compute the generalized fANOVA components analytically, even for simple functions.

### 3.2.4 Generalized Variance Decomposition

Given that the fANOVA decomposition changes under dependent inputs, we briefly make an adjustment to the second-moment statistics of the generalized fANOVA decomposition.

The mean of  $y$  remains unchanged and is still given by the constant term  $y_{\emptyset,G}$ , i.e.

$$\mu_G := \mathbb{E}[y(\mathbf{X})] = y_{\emptyset,G}.$$

In contrast, the variance decomposition does not simplify in the same way as Equation 11, as cross-terms of the same order do not vanish under hierarchical orthogonality. For  $\emptyset \neq u \subseteq \{1, \dots, N\}$ ,  $\emptyset \neq v \subseteq \{1, \dots, N\}$ ,  $u \not\subseteq v$  we restate from above:

$$\begin{aligned} \sigma^2 &:= \mathbb{E}[(y(\mathbf{X}) - \mu_G)^2] \\ &= \mathbb{E}\left[\left(y_{\emptyset,G} + \sum_u y_{u,G}(\mathbf{X}_u) - y_{\emptyset,G}\right)^2\right] \\ &= \mathbb{E}\left[\left(\sum_u y_{u,G}(\mathbf{X}_u)\right)^2\right] \\ &= \sum_u \mathbb{E}[y_{u,G}^2(\mathbf{X}_u)] + \sum_{u \not\subseteq v, v \not\subseteq u} \mathbb{E}[y_{u,G}(\mathbf{X}_u)y_{v,G}(\mathbf{X}_v)], \end{aligned}$$

while the first sum in the final line goes over all nonempty subsets  $u$  and the second sum goes over all pairs of subsets  $(u, v)$  where neither is a subset of the other one. Conceptually this means that the first term is the sum of the variances of the components, while the second term is the sum of the covariances between components that are not hierarchically orthogonal. The indices under the second component capture precisely the cross-terms that do not vanish under hierarchical orthogonality. As we saw earlier, cross-terms of the same hierarchy also cancel out under the orthogonality assumption of the classical fANOVA.

### 3.2.5 Example: Dependent Multivariate Normal Inputs

Before ending this section, it remains to answer how the true generalized fANOVA decomposition looks like for our running example (Equation 7). While the interdependence of the generalized components makes it difficult to arrive at an analytical solution, Rahman (2014) provides a way to obtain the closed-form solution for any polynomial of maximum two degree under normally distributed input variables.

The approach in Rahman (2014) is based on a Fourier-polynomial expansion, which expresses each generalized fANOVA component as a weighted sum of basis functions. This shifts the problem from directly determining the fANOVA component functions to identifying suitable basis functions that can represent the fANOVA terms. Rahman chooses Hermite polynomials as these basis functions because they are, by construction,

zero-centered and hierarchically orthogonal. These properties ensure that the resulting fANOVA components are also zero-centered and hierarchically orthogonal. The remaining challenge is then to determine the weights associated with the basis functions, which can be obtained through coefficient matching. A polynomial of degree two has the general form

$$y(x_1, x_2) = a_0 + a_1x_1 + a_2x_2 + a_{11}x_1^2 + a_{22}x_2^2 + a_{12}x_1x_2. \quad (21)$$

Any such polynomial may be expressed as a sum of weighted basis functions of the form (Nagler, 2024b):

$$\begin{aligned} y(x_1, x_2) = & c_0 + c_{1,1} \psi_{1,1}(x_1) + c_{2,1} \psi_{2,1}(x_2) \\ & + c_{1,2} \psi_{1,2}(x_1) + c_{2,2} \psi_{2,2}(x_2) \\ & + c_{12,11} \psi_{12,11}(x_1, x_2), \end{aligned}$$

where the  $\psi_{i,j}$  are the basis functions with corresponding weights  $c_0, \dots, c_{12,11} \in \mathbb{R}$ . The idea is to carefully construct a set of basis functions which fulfill zero-mean and hierarchical orthogonality. Then the expansion in these basis functions is already the fANOVA decomposition of a quadratic polynomial, i.e.

$$\begin{aligned} y(x_1, x_2) &= a_0 + a_1x_1 + a_2x_2 + a_{11}x_1^2 + a_{22}x_2^2 + a_{12}x_1x_2 \\ &= c_0 + c_{1,1} \psi_{1,1}(x_1) + c_{2,1} \psi_{2,1}(x_2) \\ &\quad + c_{1,2} \psi_{1,2}(x_1) + c_{2,2} \psi_{2,2}(x_2) + c_{12,11} \psi_{12,11}(x_1, x_2) \\ &= \underbrace{c_0}_{y_0} + \underbrace{(c_{1,1} \psi_{1,1}(x_1) + c_{1,2} \psi_{1,2}(x_1))}_{y_1(x_1)} \\ &\quad + \underbrace{(c_{2,1} \psi_{2,1}(x_2) + c_{2,2} \psi_{2,2}(x_2))}_{y_2(x_2)} \\ &\quad + \underbrace{c_{12,11} \psi_{12,11}(x_1, x_2)}_{y_{12}(x_1, x_2)}. \end{aligned}$$

Derived from the probability density of a multivariate normal distribution, Rahman (2014) chooses multivariate Hermite polynomials. We use a slightly simplified version of the proposed basis functions to find an explicit solution for our running example<sup>1</sup>. The basis

---

<sup>1</sup>We omit the scaling factor, which means the basis functions are not orthonormal anymore but still orthogonal.



functions we work with are:

$$\begin{aligned}
 \psi_{\emptyset}(x_1, x_2) &= 1, \\
 \psi_{1,1}(x_1) &= x_1, \\
 \psi_{2,1}(x_2) &= x_2, \\
 \psi_{1,2}(x_1) &= x_1^2 - 1, \\
 \psi_{2,2}(x_2) &= x_2^2 - 1, \\
 \psi_{12,11}(x_1, x_2) &= \frac{\rho(x_1^2 + x_2^2)}{1 + \rho^2} - x_1x_2 + \frac{\rho(\rho^2 - 1)}{1 + \rho^2},
 \end{aligned}$$

where  $\rho$  is the correlation coefficient between  $X_1$  and  $X_2$ . So this formula will work for dependent as well as independent inputs.

What remains it to find the coefficients  $c_0, c_{1,1}, \dots, c_{12,11}$  such that the weighted sum of the basis functions truly recovers the original polynomial in Equation 21. To find the correct weights, we plug in the basis functions and rearrange terms to recognize the groups more easily:

$$\begin{aligned}
 y(x_1, x_2) &= c_0 + c_{1,1}x_1 + c_{2,1}x_2 + c_{1,2}(x_1^2 - 1) + c_{2,2}(x_2^2 - 1) \\
 &\quad + c_{12,11} \left( \frac{\rho(x_1^2 + x_2^2)}{1 + \rho^2} - x_1x_2 + \frac{\rho(\rho^2 - 1)}{1 + \rho^2} \right) \\
 &= \left( c_0 - c_{1,2} - c_{2,2} + c_{12,11} \frac{\rho(\rho^2 - 1)}{1 + \rho^2} \right) + c_{1,1} x_1 + c_{2,1} x_2 \\
 &\quad + \left( c_{1,2} + c_{12,11} \frac{\rho}{1 + \rho^2} \right) x_1^2 + \left( c_{2,2} + c_{12,11} \frac{\rho}{1 + \rho^2} \right) x_2^2 - c_{12,11} x_1x_2.
 \end{aligned}$$

Now we can use monomial matching to find the coefficients. It is best to start with the interaction term and work backwards from there to the constant term, plugging in the current solutions along the way:

$$\begin{aligned}
 -c_{12,11} &= a_{12} & \Rightarrow & \quad c_{12,11} = -a_{12} \\
 c_{1,2} + c_{12,11} \frac{\rho}{1 + \rho^2} &= a_{11} & \Rightarrow & \quad c_{1,2} = a_{11} + \frac{\rho}{1 + \rho^2} a_{12} \\
 c_{2,2} + c_{12,11} \frac{\rho}{1 + \rho^2} &= a_{22} & \Rightarrow & \quad c_{2,2} = a_{22} + \frac{\rho}{1 + \rho^2} a_{12} \\
 c_{1,1} &= a_1 \\
 c_{2,1} &= a_2 \\
 c_0 - c_{1,2} - c_{2,2} + c_{12,11} \frac{\rho(\rho^2 - 1)}{1 + \rho^2} &= a_0 & \Rightarrow & \quad c_0 = a_0 + a_{11} + a_{22} + \rho a_{12}
 \end{aligned}$$

Hence, the generalized fANOVA decomposition of a two-degree polynomial is given by:

$$\begin{aligned}
 y(x_1, x_2) &= c_0 + c_{1,1} \psi_{1,1}(x_1) + c_{2,1} \psi_{2,1}(x_2) \\
 &\quad + c_{1,2} \psi_{1,2}(x_1) + c_{2,2} \psi_{2,2}(x_2) + c_{12,11} \psi_{12,11}(x_1, x_2) \\
 &= \underbrace{(a_0 + a_{11} + a_{22} + \rho a_{12})}_{c_0} + \underbrace{a_1}_{c_{1,1}} x_1 + \underbrace{a_2}_{c_{2,1}} x_2 \\
 &\quad + \underbrace{\left(a_{11} + \frac{\rho}{1 + \rho^2} a_{12}\right)}_{c_{1,2}} (x_1^2 - 1) + \underbrace{\left(a_{22} + \frac{\rho}{1 + \rho^2} a_{12}\right)}_{c_{2,2}} (x_2^2 - 1) \\
 &\quad + \underbrace{(-a_{12})}_{c_{12,11}} \left( \frac{\rho(x_1^2 + x_2^2)}{1 + \rho^2} - x_1 x_2 + \frac{\rho(\rho^2 - 1)}{1 + \rho^2} \right) \\
 &= (a_0 + a_{11} + a_{22} + \rho a_{12}) + a_1 x_1 + a_2 x_2 \\
 &\quad + \left(a_{11} + \frac{\rho}{1 + \rho^2} a_{12}\right) (x_1^2 - 1) + \left(a_{22} + \frac{\rho}{1 + \rho^2} a_{12}\right) (x_2^2 - 1) \\
 &\quad - a_{12} \left( \frac{\rho(x_1^2 + x_2^2)}{1 + \rho^2} - x_1 x_2 + \frac{\rho(\rho^2 - 1)}{1 + \rho^2} \right),
 \end{aligned}$$

with individual components:

$$\begin{aligned}
 y_\emptyset &= a_0 + a_{11} + a_{22} + \rho a_{12}, \\
 y_{\{1\}}(x_1) &= a_1 x_1 + \left(a_{11} + \frac{\rho}{1 + \rho^2} a_{12}\right) (x_1^2 - 1), \\
 y_{\{2\}}(x_2) &= a_2 x_2 + \left(a_{22} + \frac{\rho}{1 + \rho^2} a_{12}\right) (x_2^2 - 1), \\
 y_{\{1,2\}}(x_1, x_2) &= -a_{12} \left( \frac{\rho(x_1^2 + x_2^2)}{1 + \rho^2} - x_1 x_2 + \frac{\rho(\rho^2 - 1)}{1 + \rho^2} \right).
 \end{aligned} \tag{22}$$

This set of component functions is true under the assumption of Gaussian inputs. The basis representation is still correct for other distribution assumptions in the sense that it recovers the original function; however, the component function would not be hierarchically orthogonal anymore.

With this we are able to give the fANOVA terms for our running example in a generalized form, which allows for dependent input variables, assumed to be Gaussian.

For  $h(x_1, x_2) = x_1 + 2x_2 + x_1 x_2$  we have  $a_0 = 0, a_1 = 1, a_2 = 2, a_{11} = 0, a_{22} = 0, a_{12} = 1,$

and therefore obtain with Equation 22:

$$\begin{aligned}h_{\emptyset} &= \rho, \\h_{\{1\}}(x_1) &= x_1 + \frac{\rho}{1 + \rho^2}(x_1^2 - 1), \\h_{\{2\}}(x_2) &= 2x_2 + \frac{\rho}{1 + \rho^2}(x_2^2 - 1), \\h_{\{1,2\}}(x_1, x_2) &= -\left(\frac{\rho(x_1^2 + x_2^2)}{1 + \rho^2} - x_1x_2 + \frac{\rho(\rho^2 - 1)}{1 + \rho^2}\right).\end{aligned}$$

## 4 Visualization and Estimation

In the final section of this thesis, we will explore the fANOVA decomposition visually. This will provide a better understanding for how fANOVA components behave in different scenarios. We will first revisit our running example and then explore some other functions.

### 4.1 Comparison of Decompositions

Recall our running example (Equation 7):

$$h(x_1, x_2) = x_1 + 2x_2 + x_1x_2,$$

with polynomial coefficients:  $a_0 = 0$ ,  $a_1 = 1$ ,  $a_2 = 2$ ,  $a_{11} = 0$ ,  $a_{22} = 0$ ,  $a_{12} = 1$ . Under independent inputs ( $\rho = 0$ ), the fANOVA components are given by:

$$\begin{aligned} h_{\emptyset} &= 0, \\ h_{\{1\}}(x_1) &= x_1 \\ h_{\{2\}}(x_2) &= 2x_2 \\ h_{\{1,2\}}(x_1, x_2) &= x_1x_2, \end{aligned}$$

visualized in Figure 1. As expected, we observe simple linear functions and a regular symmetric contour plot.




Figure 1: Main effects (left) and interaction effect (right) of the fANOVA decomposition for  $h(x_1, x_2) = x_1 + 2x_2 + x_1x_2$  with independent inputs.

Now we assume  $\rho = 0.5$ . In attempt to compute the fANOVA terms under dependent inputs, we calculated the following effects, which do not satisfy the fANOVA property of orthogonality. Nevertheless, is it interesting to compare their visualization in Figure 2 to

the one of the true generalized fANOVA components in Figure 3. The main effects are parabolic, and the interaction term seems to be non-symmetric.




Figure 2: Main effects (left) and interaction effect (right) of a fANOVA-type decomposition for  $h(x_1, x_2) = x_1 + 2x_2 + x_1x_2$  with dependent inputs,  $\rho = 0.5$ .

The true fANOVA components under  $\rho = 0.5$  are given by:

$$\begin{aligned}y_0 &= 0.5, \\ y_1(x_1) &= x_1 + 0.4(x_1^2 - 1) = x_1 + 0.4x_1^2 - 0.4, \\ y_2(x_2) &= 2x_2 + 0.4(x_2^2 - 1) = 2x_2 + 0.4x_2^2 - 0.4, \\ y_{12}(x_1, x_2) &= -\left(0.4(x_1^2 + x_2^2) - x_1x_2 - 0.3\right) \\ &= -0.4x_1^2 - 0.4x_2^2 + x_1x_2 + 0.3.\end{aligned}$$

These are visualized in Figure 3. Interestingly, the parabolic form of the main effects is similar between both decompositions, but the interaction effects diverge notably.

Our running example (Equation 7) included linear effects of both input variables and the interaction term. For the remainder of this section, we will explore other representative scenarios, we can build within the scaffold of a bivariate two-degree polynomial.




Figure 3: Main effects (left) and interaction effect (right) of the generalized fANOVA decomposition for  $h(x_1, x_2) = x_1 + 2x_2 + x_1x_2$  with dependent inputs,  $\rho = 0.5$ .

## 4.2 Comparison of Functions

### 4.2.1 Scenario: Linear

First, we consider two-degree polynomials of the form:

$$q(x_1, x_2) = a_1x_1 + a_2x_2.$$

We can immediately read off the fANOVA components or use the general set of fANOVA components for a two-degree polynomial in Equation 22 which simplify for  $g_1$  to:

$$q_{\{1\}}(x_1) = a_1x_1,$$

$$q_{\{2\}}(x_2) = a_2x_2.$$

The function  $q$  can solely be described by linear main effects (Figure 4). Since no interaction effect is present varying  $\rho$  has no impact on the main effects.

### 4.2.2 Scenario: Linear and Quadratic

Slightly more complex is a two-degree polynomials, which allows for effects of linear and quadratic nature:

$$p(x_1, x_2) = a_1x_1 + a_2x_2 + a_{11}x_1^2 + a_{22}x_2^2.$$




Figure 4: Main fANOVA components for linear terms with different coefficients. The components are given by:  $y_1(x_1) = a_1x_1$ ,  $y_2(x_2) = a_2x_2$ .

The fANOVA components for  $p$  are given by:

$$\begin{aligned}
 p_{\emptyset} &= a_{11} + a_{22}, \\
 p_{\{1\}}(x_1) &= a_1x_1 + a_{11}(x_1^2 - 1), \\
 p_{\{2\}}(x_2) &= a_2x_2 + a_{22}(x_2^2 - 1).
 \end{aligned}$$

The main effects are parabolas now. In Figure 5, we vary the coefficients  $a_1$ ,  $a_2$ ,  $a_{11}$ , and  $a_{22}$ , while the interaction term is still absent. The coefficients of the quadratic terms determine whether the parabola is facing downwards or upwards; when  $a_{11}$  and  $a_{22}$  are both negative or both positive the parabola is open downwards or upwards respectively, and when they have opposite signs the parabolas are open in different directions. Alongside the quadratic coefficients, the linear ones  $a_1$  and  $a_2$  influence how stretched or compressed the parabola is.

#### 4.2.3 Scenario: Interaction

Next, we consider a model, which solely consists of an interaction term:

$$w(x_1, x_2) = a_{12}x_1x_2.$$




Figure 5: Main effects for different coefficient combinations of mixed main effects. The components are given by:  $y_1(x_1) = a_1x_1 + a_{11}(x_1^2 - 1)$ ,  $y_2(x_2) = a_2x_2 + a_{22}(x_2^2 - 1)$ .



The fANOVA components for  $w$  are given by:

$$\begin{aligned} w_{\emptyset} &= a_{12}\rho, \\ w_{\{1\}}(x_1) &= a_{12}\frac{\rho}{1+\rho}(x_1^2 - 1), \\ w_{\{2\}}(x_2) &= a_{12}\frac{\rho}{1+\rho}(x_2^2 - 1), \\ w_{\{1,2\}}(x_1, x_2) &= -a_{12}\left(\frac{\rho(x_1^2 + x_2^2)}{1+\rho^2} - x_1x_2 + \frac{\rho(\rho^2 - 1)}{1+\rho^2}\right). \end{aligned}$$

The main effects  $y_1$  and  $y_2$ , as well as the interaction term  $y_{12}$ , are influenced by  $\rho$  and  $a_{12}$ . In our example we keep  $a_{12} = 2$  fixed and show the interaction effect as a contour plot for varying  $\rho$  with the corresponding main effects next to it Figure 6. The main effects have the same form for every case of  $\rho$  and  $a_{12}$  and thus overlap. This example is simple yet interesting because it shows that in the case where the true function consists solely of an interaction term, fANOVA still attributes something to the isolated effect of each variable. Only when the variables are uncorrelated, all the effect is attributed to the interaction term. This functionality hints to why Lengerich et al. (2020) build an algorithm around fANOVA to purify interaction effects<sup>2</sup>.

#### 4.2.4 Scenario: Full

Finally, a full example, including all main and interaction effects:

$$z(x_1, x_2) = a_1x_1 + a_2x_2 + a_{11}x_1^2 + a_{22}x_2^2 + a_{12}x_1x_2.$$

Now the fANOVA components are given by Equation 22, where  $a_0 = 0$ . We can vary the coefficients as well as  $\rho$ .

When the true function has no interaction term, as in our first two scenarios, varying  $\rho$  is uninteresting because there is no way it could influence the form of the main effects. In this full scenario, however, there is an interaction term present, and therefore it is most interesting to compare pairs of coefficient sets under  $\rho = 0$  versus  $\rho \neq 0$ . With this we want to essentially ask how effects are distorted by performing the classical fANOVA decomposition when a true interaction effect is present and variables exhibit dependency. In Figure 7 we make this comparison for a weak linear correlation between variables and in Figure 8 we show the same for a strong linear correlation between variables. Similar to

<sup>2</sup>Because they see a pure interaction as an effect which cannot be attributed to lower order terms; this means when identifying interactions we want to attribute all we can to lower order terms and what is left is the true interaction effect.






Figure 6: Main effects (left column) and interaction contour plots (right column) for different values of  $\rho$ . The effects are given by  $y_1(x_1) = a_{12} \frac{\rho}{1+\rho} (x_1^2 - 1)$ ,  $y_2(x_2) = a_{12} \frac{\rho}{1+\rho} (x_2^2 - 1)$ ,  $y_{12}(x_1, x_2) = -a_{12} \left( \frac{\rho(x_1^2 + x_2^2)}{1+\rho^2} - x_1 x_2 + \frac{\rho(\rho^2 - 1)}{1+\rho^2} \right)$ .

the visualization of our running example in Figure 3, we see that main effects are distorted slightly, while interaction effects look substantially different under dependent inputs.



(a) Main and interaction effects for (1)  $a_1 = 2, a_2 = 0, a_{11} = 1, a_{22} = 0, a_{12} = 0.5, \rho = 0.3$ .




(b) Main and interaction effects for (2)  $a_1 = 0, a_2 = 2, a_{11} = 0, a_{22} = 1, a_{12} = -0.5, \rho = 0$ .

Figure 7: Main effects (left) and interaction contours (right) for four different coefficient sets.


### 4.3 Estimation of fANOVA components

This was all rather theoretical and the examples we used foster understanding, but they are toy examples and in reality the true function is unknown and more complex. So to become a more widely used, established interpretability method, an estimation scheme is inevitable. We already encountered one estimation scheme proposed by Rahman (2014) when computing the generalized fANOVA components for our running example; we refer to section 3 the conceptual idea behind it.

In Hooker (2004) an estimation framework based on partial dependence is proposed, which makes use of the formulation of fANOVA via projections. To obtain the component estimate for  $y_u$ , Hooker proposed to estimate the projections of  $y$  onto the subspace of variables spanned by  $u$  empirically. One does so by first estimating the conditional



(a) Main and interaction effects for (3)  $a_1 = -2$ ,  $a_2 = -2$ ,  $a_{11} = 1$ ,  $a_{22} = -1$ ,  $a_{12} = 1$ ,  $\rho = -0.8$ .



(b) Main and interaction effects for (4)  $a_1 = -2$ ,  $a_2 = 2$ ,  $a_{11} = -1$ ,  $a_{22} = 1$ ,  $a_{12} = -1$ ,  $\rho = 0$ .

Figure 8: Main effects (left) and interaction contours (right) for four different coefficient sets.

expected value of the variables in  $u$ . This is a simple Monte Carlo estimation, which results in the partial dependence function (PD Function) for the variables in  $u$  (Hooker, 2004). The PD Function can then be used to estimate the empirical projection of interest. He states that his method works well for functions that truly have nearly additive structure and purely additive functions are exactly recoverable with this approach. However, the approach suffers from extrapolation issues or artefacts when the true function involves interactions and inputs are dependent.

Therefore, in Hooker (2007) a new estimation scheme is proposed for his version of the generalized fANOVA decomposition (see section 3). Hooker rewrites his proposed system of equations as restricted weighted least squares problem and solves it via Lagrange multiplier for the exact solution of the simultaneously defined generalized components. The function is evaluated at a grid of points to reduce computational costs. Because of the parallel to weighted least squares, it is also possible to compute a weighted standard ANOVA with existing software; however, like so it is difficult to incorporate the system constraints and one might obtain components that are not hierarchical orthogonal.

None of these estimation approaches has a standard software implementation or published code. Some existing unfinished implementations are numerically instable or yield illogical results. This underpins the need for a more robust estimation scheme with stable software implementation.

## 5 Conclusion

The fANOVA decomposition is a foundational method that has been studied from many perspectives, with recent interest driven by its potential in model interpretability. Its key strength lies in producing terms that represent the unique contribution of each variable or interaction, cleanly separated from one another without mixing effects.

Despite being established in the literature, we found a lack of unified formalization and notational clarity, which we aimed to address in this work.

We began by revisiting the classical fANOVA decomposition, bridging Sobol’s initial formulation with more recent developments. We filled small gaps in mathematical proofs and illustrated the parallel to the Hoeffding decomposition under independent and zero-centered inputs. Further, we established a direct connection to conditional expectations and orthogonal projections, which we argue is key to unifying different existing notations and formal approaches. Alongside the functional decomposition, we presented the variance decomposition, which underpins Sobol indices.

Next, we extended the decomposition to dependent input variables. Here, multiple approaches exist; we focused on the frameworks of Hooker (2007) and Rahman (2014). Using an illustrative example, we demonstrated that obtaining interpretable fANOVA components under dependence requires the careful construction of specific constraints. We also noted that closed-form solutions for the generalized components are difficult to obtain in practice and remain an open problem. Nevertheless, the associated variance decomposition still holds, allowing for the construction of generalized Sobol indices.

To complement the theoretical work, we visualized fANOVA components for simple polynomial functions with Gaussian inputs. These visualizations illustrated how fANOVA separates effects, but were limited to simple functions and Gaussian inputs.

This work did not present a full treatment of Sobol indices, and only touched on variance decomposition as their foundation. Our empirical demonstrations were restricted to toy examples; applying fANOVA in practice will require estimation methods for trained models on real data. We also briefly mentioned the ability of fANOVA to purify interaction effects (Lengerich et al., 2020) and its relation to other interpretability techniques (e.g., PD, Shapley (Fumagalli et al., 2025)), but did not explore these connections in depth.

Future work could extend Rahman’s polynomial expansion approach to more complex polynomials and non-Gaussian distributions, and further investigate mathematical connections to other interpretability frameworks. Additionally, a lack of standard software for performing fANOVA remains a barrier to its widespread adoption, so developing robust, open-source implementations would be an important step toward enabling its use in practical model interpretability.

## A Appendix

### Proof of classical fANOVA decomposition

Here we show the proof of Theorem 1 in Sobol (1993).

**Theorem A.1.** *Any function  $y$ , which is integrable over the unit hypercube  $[0, 1]^k$ , has a unique fANOVA expansion of the form:*

$$y(\mathbf{X}) = \sum_{u \subseteq \{1, \dots, N\}} y_u(\mathbf{X}_u),$$

*subject to the zero-mean constraint Equation 3.*

Sobol proofs existence and uniqueness of the fANOVA decomposition by showing how the summands of the desired decomposition look and showing that they satisfy the desired zero-mean property.

*Proof.* Assume that  $\mathbf{X}$  is an  $N$ -dimensional vector of independent random variables and that the zero-mean property holds for the - still abstract - fANOVA terms. He defines the integral w.r.t. all variables except for the ones with indices in  $v$ :

$$g_v(\mathbf{x}_v) = \int_{[0,1]^{N-|v|}} y(\mathbf{x}) f_{\mathbf{X}}(\mathbf{x}) d\nu(\mathbf{x}_{-v})$$

He then builds the fANOVA terms subsequently and shows that they indeed satisfy the desired properties.

The very first term in the decomposition is the integral of  $y$  with respect to all variables:

$$y_0 = \int_{\mathbb{R}^N} y(\mathbf{x}) f_{\mathbf{X}}(\mathbf{x}) d\nu(\mathbf{x})$$

This integral exists because  $y \in \mathcal{L}^2(\mathbb{R}^N, f_{\mathbf{X}}(\mathbf{x})d\nu)$ , and the product measure is finite on the domain.

Next, Sobol derives the one-dimensional fANOVA terms. For this, he takes the integral

of Equation 1 w.r.t. all variables except for the one with index  $i$ , so  $v_1 = \{i\}$ :

$$\begin{aligned} \int_{\mathbb{R}^{N-1}} y(\mathbf{x}) f_{\mathbf{X}}(\mathbf{x}) d\nu(\mathbf{x}_{-v_1}) &= \int_{\mathbb{R}^{N-1}} \sum_{u \subseteq \{1, \dots, N\}} y_u(\mathbf{x}_u) f_{\mathbf{X}}(\mathbf{x}) d\nu(\mathbf{x}_{-v_1}) \\ &= \sum_{u \subseteq \{1, \dots, N\}} \int_{\mathbb{R}^{N-1}} y_u(\mathbf{x}_u) f_{\mathbf{X}}(\mathbf{x}) d\nu(\mathbf{x}_{-v_1}) \\ &= \sum_{u \subseteq \{1, \dots, N\}} \int_{\mathbb{R}^{N-1}} y_u(\mathbf{x}_u) \left( \prod_{j=1}^N f_{\{j\}}(x_j) \right) d\nu(\mathbf{x}_{-v_1}) \end{aligned}$$

For every summand  $y_u(\mathbf{x}_u)$  with  $u \not\ni i$ , the integrand does not depend on  $x_i$ , and thus vanished due to the zero-mean constraint. Similarly, for any term  $y_u(\mathbf{x}_u)$  with  $i \in u$  and  $|u| > 1$ , the integration will include at least one other variable in  $u$ , again causing the integral to vanish. In the end, only the constant term  $y_\emptyset$  and the one-dimensional term  $y_{\{i\}}(x_i)$  remain, which depend only on  $x_i$  and are not integrated. Therefore, we can derive the simplified expression:

$$\int_{\mathbb{R}^{N-1}} y(\mathbf{x}) f_{\mathbf{X}}(\mathbf{x}) d\nu(\mathbf{x}_{-v_1}) = y_\emptyset + y_{\{i\}}(x_i).$$

This equation allows to define the one-dimensional term  $y_{\{i\}}$  explicitly as:

$$y_{\{i\}}(x_i) = \int_{\mathbb{R}^{N-1}} y(\mathbf{x}) f_{\mathbf{X}}(\mathbf{x}) d\nu(\mathbf{x}_{-v_1}) - y_\emptyset.$$

Next, he considers  $v_2 = \{i, j\}$ . The ANOVA decomposition is integrated over all variables except  $x_i$  and  $x_j$ :

$$\begin{aligned} \int_{\mathbb{R}^{N-2}} y(\mathbf{x}) f_{\mathbf{X}}(\mathbf{x}) d\nu(\mathbf{x}_{-\{i,j\}}) &= \int_{\mathbb{R}^{N-2}} \sum_{u \subseteq \{1, \dots, N\}} y_u(\mathbf{x}_u) f_{\mathbf{X}}(\mathbf{x}) d\nu(\mathbf{x}_{-\{i,j\}}) \\ &= \sum_{u \subseteq \{1, \dots, N\}} \int_{\mathbb{R}^{N-2}} y_u(\mathbf{x}_u) \left( \prod_{j=1}^N f_{\{j\}}(x_j) \right) d\nu(\mathbf{x}_{-\{i,j\}}) \\ &= y_\emptyset + y_{\{i\}}(x_i) + y_{\{j\}}(x_j) + y_{\{i,j\}}(x_i, x_j) \end{aligned}$$

Hence, the two-dimensional components are given by:

$$y_{\{i,j\}}(x_i, x_j) = \int_{\mathbb{R}^{N-2}} y(\mathbf{x}) f_{\mathbf{X}}(\mathbf{x}) d\nu(\mathbf{x}_{-\{i,j\}}) - y_\emptyset - y_{\{i\}}(x_i) - y_{\{j\}}(x_j)$$



One can continue this process for all combinations of indices  $v \subseteq \{1, \dots, N\}$  to derive the corresponding fANOVA terms  $y_v(\mathbf{x}_v)$ .

Now let  $v \subseteq \{1, \dots, N\}$ . The general expression for the component  $y_v(\mathbf{x}_v)$  is given by:

$$y_v(\mathbf{x}_v) = \int_{\mathbb{R}^{N-|v|}} y(\mathbf{x}) f_{\mathbf{X}}(\mathbf{x}) d\nu(\mathbf{x}_{-v}) - \sum_{u \subsetneq v} y_u(\mathbf{x}_u)$$

The last term is the decomposition integrated with respect to no variables, i.e., the function itself:

$$y_{\{1, \dots, N\}}(\mathbf{x}) = y(\mathbf{x}) - \sum_{u \subsetneq \{1, \dots, N\}} y_u(\mathbf{x}_u)$$

Finally, it remains to verify that the constructed component functions satisfy the zero-mean constraint. Let  $v \subseteq \{1, \dots, N\}$ , and let  $i \in v$ . Then:

$$\begin{aligned} \int y_v(\mathbf{x}_v) f_{X_i}(x_i) d\nu(x_i) &= \int \left( \int y(\mathbf{x}) f_{\mathbf{X}}(\mathbf{x}) d\nu(\mathbf{x}_{-v}) - \sum_{u \subsetneq v} y_u(\mathbf{x}_u) \right) f_{\{i\}}(x_i) d\nu(x_i) \\ &= \int \left( \int y(\mathbf{x}) f_{\mathbf{X}}(\mathbf{x}) d\nu(\mathbf{x}_{-v}) \right) f_{\{i\}}(x_i) d\nu(x_i) - \sum_{u \subsetneq v} \int y_u(\mathbf{x}_u) f_{\{i\}}(x_i) d\nu(x_i) \\ &= \int y(\mathbf{x}) f_{\mathbf{X}}(\mathbf{x}) d\nu(\mathbf{x}_{-v}) d\nu(x_i) - \sum_{u \subsetneq v} \int y_u(\mathbf{x}_u) f_{\mathbf{X}_u}(\mathbf{x}_u) d\nu(\mathbf{x}_u) \end{aligned}$$

The first term integrates out all of  $\mathbf{x}_v$ , leaving  $y_0$ . Each term in the sum vanishes by the zero-mean property of lower-order components:

$$\int y_v(\mathbf{x}_v) f_{\{i\}}(x_i) d\nu(x_i) = y_0 - y_0 = 0$$

Thus, every component  $y_v(\mathbf{x}_v)$  satisfies:

$$\int y_v(\mathbf{x}_v) f_{\{i\}}(x_i) d\nu(x_i) = 0, \quad \text{for all } i \in v.$$

□

## B Electronic appendix

Data, code and figures are provided in electronic form at: [https://github.com/juliet-fleischer/fANOVA\\_decomposition](https://github.com/juliet-fleischer/fANOVA_decomposition)

## References

- Borgonovo, E., Li, G., Barr, J., Plischke, E. and Rabitz, H. (2022). Global Sensitivity Analysis with Mixtures: A Generalized Functional ANOVA Approach, *Risk Analysis* **42**(2): 304–333.  
**URL:** <https://onlinelibrary.wiley.com/doi/10.1111/risa.13763>
- Chastaing, G., Gamboa, F. and Prieur, C. (2012). Generalized Hoeffding-Sobol decomposition for dependent variables - application to sensitivity analysis, *Electronic Journal of Statistics* **6**: 2420–2448.
- Choi, Y., Park, S., Park, C., Kim, D. and Kim, Y. (2025). Meta-anova: screening interactions for interpretable machine learning, *Journal of the Korean Statistical Society* .  
**URL:** <https://link.springer.com/10.1007/s42952-024-00302-2>
- Fumagalli, F., Muschalik, M., Hüllermeier, E., Hammer, B. and Herbinger, J. (2025). Unifying Feature-Based Explanations with Functional ANOVA and Cooperative Game Theory. arXiv:2412.17152 [cs].  
**URL:** <http://arxiv.org/abs/2412.17152>
- Gu, C. (2013). *Smoothing Spline ANOVA Models*, Vol. 297 of *Springer Series in Statistics*, Springer New York, New York, NY.  
**URL:** <https://link.springer.com/10.1007/978-1-4614-5369-7>
- Hoeffding, W. (1948). A Class of Statistics with Asymptotically Normal Distribution, *The Annals of Mathematical Statistics* **19**(3): 293–325. Publisher: Institute of Mathematical Statistics.  
**URL:** <https://www.jstor.org/stable/2235637>
- Hooker, G. (2004). Discovering additive structure in black box functions, *Proceedings of the tenth ACM SIGKDD international conference on Knowledge discovery and data mining*, ACM, Seattle WA USA, pp. 575–580.  
**URL:** <https://dl.acm.org/doi/10.1145/1014052.1014122>
- Hooker, G. (2007). Generalized Functional ANOVA Diagnostics for High-Dimensional Functions of Dependent Variables, *Journal of Computational and Graphical Statistics* **16**(3): 709–732.  
**URL:** <http://www.tandfonline.com/doi/abs/10.1198/106186007X237892>

- Hu, L., Nair, V. N., Sudjianto, A., Zhang, A., Chen, J. and Yang, Z. (2025). Interpretable Machine Learning Based on Functional ANOVA Framework: Algorithms and Comparisons, *Applied Stochastic Models in Business and Industry* **41**(1): e2916.  
**URL:** <https://onlinelibrary.wiley.com/doi/10.1002/asmb.2916>
- Il Idrissi, M., Bousquet, N., Gamboa, F., Iooss, B. and Loubes, J.-M. (2025). Hoeffding decomposition of functions of random dependent variables, *Journal of Multivariate Analysis* **208**: 105444.  
**URL:** <https://linkinghub.elsevier.com/retrieve/pii/S0047259X25000399>
- König, G., Günther, E. and Luxburg, U. v. (2024). Disentangling Interactions and Dependencies in Feature Attribution. arXiv:2410.23772.  
**URL:** <http://arxiv.org/abs/2410.23772>
- Lengerich, B., Tan, S., Chang, C.-H., Hooker, G. and Caruana, R. (2020). Purifying Interaction Effects with the Functional ANOVA: An Efficient Algorithm for Recovering Identifiable Additive Models, *Proceedings of the Twenty Third International Conference on Artificial Intelligence and Statistics*, PMLR, pp. 2402–2412. ISSN: 2640-3498.  
**URL:** <https://proceedings.mlr.press/v108/lengerich20a.html>
- Liu, R. and Owen, A. B. (2006). Estimating Mean Dimensionality of Analysis of Variance Decompositions, *Journal of the American Statistical Association* **101**(474): 712–721.  
**URL:** <https://www.tandfonline.com/doi/full/10.1198/016214505000001410>
- Molnar, C. (2025). *Interpretable Machine Learning*, 3 edn.  
**URL:** <https://christophm.github.io/interpretable-ml-book>
- Muehlenstaedt, T., Roustant, O., Carraro, L. and Kuhnt, S. (2012). Data-driven Kriging models based on FANOVA-decomposition, *Statistics and Computing* **22**(3): 723–738.  
**URL:** <http://link.springer.com/10.1007/s11222-011-9259-7>
- Nagler, T. (2024a). Mathematical statistics, Lecture Script. <https://tnagler.github.io/mathstat-lmu-2024.pdf>.  
**URL:** <https://tnagler.github.io/mathstat-lmu-2024.pdf>
- Nagler, T. (2024b). Methoden der Linearen Algebra in der Statistik – Vorlesungsskript, <https://tnagler.github.io/linalg-2024.pdf>. Version Sommersemester 2024.

- Owen, A. B. (2013). Variance Components and Generalized Sobol' Indices, *SIAM/ASA Journal on Uncertainty Quantification* **1**(1): 19–41. tex.eprint: <https://doi.org/10.1137/120876782>.  
**URL:** <https://doi.org/10.1137/120876782>
- Owen, A. B. (2014). Sobol' Indices and Shapley Value, *SIAM/ASA Journal on Uncertainty Quantification* **2**(1): 245–251.
- Rahman, S. (2014). A Generalized ANOVA Dimensional Decomposition for Dependent Probability Measures, *SIAM/ASA Journal on Uncertainty Quantification* **2**(1): 670–697.  
**URL:** <https://doi.org/10.1137/120904378>
- Sobol, I. (2001). Global sensitivity indices for nonlinear mathematical models and their Monte Carlo estimates, *Mathematics and Computers in Simulation* **55**(1-3): 271–280.  
**URL:** <https://linkinghub.elsevier.com/retrieve/pii/S0378475400002706>
- Sobol, I. M. (1993). Sensitivity Estimates for Nonlinear Mathematical Models, *Mathematical Modelling and Computational Experiments* **1**: 407–414.
- Stone, C. J. (1994). The Use of Polynomial Splines and Their Tensor Products in Multivariate Function Estimation, *The Annals of Statistics* **22**(1): 118–171. Publisher: Institute of Mathematical Statistics.  
**URL:** <https://www.jstor.org/stable/2242446>
- Stone, C. J., Hansen, M. H., Kooperberg, C. and Truong, Y. K. (1997). Polynomial Splines and their Tensor Products in Extended Linear Modeling, *The Annals of Statistics* **25**(4): 1371–1425. Publisher: Institute of Mathematical Statistics.  
**URL:** <https://www.jstor.org/stable/2959054>
- Takemura, A. (1983). Tensor Analysis of ANOVA Decomposition, *Journal of the American Statistical Association* **78**(384): 894–900. Publisher: [American Statistical Association, Taylor & Francis, Ltd.].  
**URL:** <https://www.jstor.org/stable/2288201>
- Van der Vaart, A. W. (1998). *Asymptotic Statistics*, Cambridge Series in Statistical and Probabilistic Mathematics, Cambridge University Press.

## Declaration of authorship

I hereby declare that the report submitted is my own unaided work. All direct or indirect sources used are acknowledged as references. I am aware that the Thesis in digital form can be examined for the use of unauthorized aid and in order to determine whether the report as a whole or parts incorporated in it may be deemed as plagiarism. For the comparison of my work with existing sources I agree that it shall be entered in a database where it shall also remain after examination, to enable comparison with future Theses submitted. Further rights of reproduction and usage, however, are not granted here. This paper was not previously presented to another examination board and has not been published.

Location, date

---

Name