

Project Description: Multigraded Homological Algebra and Geometry

This proposal involves research in algebraic geometry and commutative algebra with several connections to computation and combinatorics, as well as a wide array of broader impacts.

- § 1 Results of Prior NSF Support.
- § 2 The Geometry of Multigraded Syzygies on Stacks
- § 3 Castelnuovo–Mumford Regularity: Resolutions, Bounds, and Asymptotics
- § 4 Uniform Homological Algebra on Toric Varieties
- § 5 Broader Impacts.

1. Results of Prior NSF Support

During 2020–2022 the PI was supported by an NSF Postdoctoral Research Fellowship (NSF Grant No. MSPRF DMS-2002239, \$150,000) titled *Asymptotic Syzygies in Algebraic Geometry*. In this period the PI made major contributions to extending our understanding of homological algebra on toric varieties (see Section 1.2) and the cohomology of moduli spaces (see Section 1.4). These works resulted in 5 new papers being posted to the arXiv [BCE⁺22, BHS24, BHS22, BBC23, BBC⁺24].

During 2015–2020 the PI was supported by an NSF Graduate Research Fellowship (NSF Grant No. DGE-1256259, \$150,000) title *Syzygies in Algebraic Geometry*. In this period the PI made substantial contributions to commutative algebra and algebraic geometry, including furthering our understanding of syzygies on higher dimensional varieties. This resulted in the PI posting 8 new papers to the arXiv [ABLS20, BBB⁺17, BE19, BL20, BEGY20, BEGY21, Bru24, Bru22a].

Of the 12 new papers the PI posted to the arXiv during these periods of support 9 of these above-mentioned papers were accepted for publication, including in such journals as *Algebra & Number Theory* [BE19], *Geometry & Topology* [BBC⁺24], *Journal of Algebra* [BCE⁺22], and *Experimental Mathematics* [BEGY20]. The PI contributed to 4 open-source software packages, one public-facing database, and two articles for the *Notices of the AMS* [BBB21, Bru22b]. A more detailed summary of these results and their intellectual merits continues in the following sections.

1.1 Asymptotic Syzygies Broadly speaking, asymptotic syzygies is the study of the graded Betti numbers (i.e. the syzygies) of a projective variety as the positivity of the embedding grows. For a smooth projective variety $X \subset \mathbb{P}^{n_d}$ embedded by a very ample line bundle L_d , following [?], set,

$$\rho_q(X, L_d) := \frac{\#\{p \in \mathbb{N} \mid \beta_{p,p+q}(X, L_d) \neq 0\}}{n_d},$$

which is the percentage of degrees in which non-zero syzygies appear [?]*Theorem 1.1. The asymptotic perspective asks how $\rho_q(X; L_d)$ behaves along the sequence of line bundles $(L_d)_{d \in \mathbb{N}}$. In this language, Green’s foundational work on the syzygies of high degree curves can be phrased in the following way.

Theorem 1.1. [?] *Let $X \subset \mathbb{P}^n$ be a smooth projective curve. If $(L_d)_{d \in \mathbb{N}}$ is a sequence of very ample line bundles on X such that $\deg L_d = d$ then $\lim_{d \rightarrow \infty} \rho_2(X; L_d) = 0$.*

Put differently, asymptotically the syzygies of curves are as simple as possible, occurring in the lowest possible degree. Ein and Lazarsfeld established the opposite asymptotic behavior for higher dimensional varieties under the analogous positivity condition.

Theorem 1.2. [?]*Theorem C *Let $X \subset \mathbb{P}^n$ be a smooth projective variety, $\dim X \geq 2$, and fix an index $1 \leq q \leq \dim X$. If $(L_d)_{d \in \mathbb{N}}$ is a sequence of very ample line bundles such that $L_{d+1} - L_d$ is constant and ample then $\lim_{d \rightarrow \infty} \rho_q(X; L_d) = 1$.*

My work weakens the ample-growth hypothesis to semi-ample growth, recall a line bundle L is *semi-ample* if $|kL|$ is base point free for $k \gg 0$ (e.g. $\mathcal{O}(1,0)$ on $\mathbb{P}^n \times \mathbb{P}^m$). I established the following asymptotic nonvanishing result for $PP^n \times \mathbb{P}^m$ embedded by $\mathcal{O}(d_1, d_2)$.

Theorem 1.3. [?] **Corollary B Let $X = \mathbb{P}^n \times \mathbb{P}^m$ and fix an index $1 \leq q \leq n + m$. There exist constants $C_{i,j}$ and $D_{i,j}$ such that*

$$\rho_q(X; \mathcal{O}(d_1, d_2)) \geq 1 - \sum_{\substack{i+j=q \\ i \leq n, j \leq m}} \left(\frac{C_{i,j}}{d_1^i d_2^j} + \frac{D_{i,j}}{d_1^{n-i} d_2^{m-j}} \right) - O\left(\frac{\text{lower ord.}}{\text{terms}}\right).$$

If both $d_1 \rightarrow \infty$ and $d_2 \rightarrow \infty$ then $\rho_q(\mathbb{P}^n \times \mathbb{P}^m; \mathcal{O}(d_1, d_2)) \rightarrow 1$, recovering the results of Ein and Lazarsfeld for $\mathbb{P}^n \times \mathbb{P}^m$. However, if d_1 is fixed and $d_2 \rightarrow \infty$ (i.e. semi-ample growth) my results bound the asymptotic percentage of non-zero syzygies away from zero. In fact I conjecture that, unlike in previously studied cases, in the semi-ample setting $\rho_q(\mathbb{P}^n \times \mathbb{P}^m; \mathcal{O}(d_1, d_2))$ does not approach 1. Proving this would require a vanishing result for asymptotic syzygies, which is open even in the ample case [?, Conjectures 7.1, 7.5].

1.2 Eisenbud-Goto on Products of Projective Space Introduced by Mumford, the Castelnuovo–Mumford Regularity of a projective variety $X \subset \mathbb{P}^r$ is a measure of the complexity of X given in terms of the vanishing of certain cohomology groups of X . Roughly speaking one should think about Castelnuovo–Mumford regularity as being a numerical measure of geometric complexity. Mumford was interested in such a measure as it plays a key role in constructing Hilbert and Quot schemes. In particular, being d -regular implies that $\mathcal{F}(d)$ is globally generated. However, Eisenbud and Goto showed that regularity is also closely connected to interesting homological properties.

Theorem 1.4. [EG84] *Let $\mathcal{F}$ be a coherent sheaf on $\mathbb{P}^r$ and $M = \bigoplus_{e \in \mathbb{Z}} H^0(\mathbb{P}^r, \mathcal{F}(e))$ the corresponding section ring. The following are equivalent: (1) M is d -regular; (2) $\beta_{p,q}(M) = 0$ for all $p \geq 0$ and $q > d + i$; (3) $M_{\geq d}$ has a linear resolution.*

MacLagan and Smith introduced multigraded Castelnuovo–Mumford regularity, where $\mathbb{P}^r$ can be replaced by any toric variety. Similarly to the definition in the classical setting multigraded Castelnuovo–Mumford regularity is defined in terms of the vanishing of cohomology groups, however, the multigraded Castelnuovo–Mumford regularity of a subvariety or module is not a single number, but an infinite subset of $\mathbb{Z}^r$. As an example, let us consider the case of products of projective spaces. Fixing a dimension vector $\mathbf{n} = (n_1, n_2, \dots, n_r) \in \mathbb{N}^r$ we let $\mathbb{P}^{\mathbf{n}} := \mathbb{P}^{n_1} \times \mathbb{P}^{n_2} \times \dots \times \mathbb{P}^{n_r}$ and $S = \mathbb{K}[x_{i,j} \mid 1 \leq i \leq r, 0 \leq j \leq n_i]$ be the Cox ring of $\mathbb{P}^{\mathbf{n}}$ with the $\text{Pic}(X) \cong \mathbb{Z}^r$ -grading given by $\deg x_{i,j} = \mathbf{e}_i \in \mathbb{Z}^r$, where $\mathbf{e}_i$ is the i -th standard basis vector in $\mathbb{Z}^r$. Fixing some notation given $\mathbf{d} \in \mathbb{Z}^r$ and $i \in \mathbb{Z}_{\geq 0}$ we let $L_i(\mathbf{d}) := \bigcup_{\mathbf{v} \in \mathbb{N}_{\setminus i}, |\mathbf{v}|=i} (\mathbf{d} - \mathbf{v}) + \mathbb{N}^r$. Note when $r = 2$ the region $L_i(\mathbf{d})$ looks like a staircase with $(i + 1)$ -corners. Roughly speaking we define regularity by requiring the i -th cohomology of certain twists of $\mathcal{F}$ to vanish on L_i .

Definition 1.5. [MS04, Definition 6.1] *A coherent sheaf $\mathcal{F}$ on $\mathbb{P}^{\mathbf{n}}$ is $\mathbf{d}$ -regular if and only if $H^i(\mathbb{P}^{\mathbf{n}}, \mathcal{F}(\mathbf{e})) = 0$ for all $\mathbf{e} \in L_i(\mathbf{d})$. The multigraded Castelnuovo–Mumford regularity of $\mathcal{F}$ is then the set:*

$$\text{reg}(\mathcal{F}) := \{\mathbf{d} \in \mathbb{Z}^r \mid \mathcal{F} \text{ is } \mathbf{d}\text{-regular}\} \subset \mathbb{Z}^r.$$

The obvious approaches to generalize Theorem 1.4 to a product of projective spaces turn out not to work. For example, the multigraded Betti numbers do not determine multigraded Castelnuovo–Mumford regularity [BHS24, Example 5.1] Despite this we show that part (3) of Theorem 1.4 can be generalized. To do so we introduce the following generalization of linear resolutions.

Definition 1.6. *A complex $F_{\bullet}$ of $\mathbb{Z}^r$ -graded free S -modules is $\mathbf{d}$ -quasilinear if and only if F_0 is generated in degree $\mathbf{d}$ and each twist of F_i is contained in $L_{i-1}(\mathbf{d} - \mathbf{1})$.*

Theorem 1.7. [BHS24, Theorem A] *Let M be a (saturated) finitely generated $\mathbb{Z}^r$ -graded S -module: M is $\mathbf{d}$ -regular $\iff M_{\geq \mathbf{d}}$ has a $\mathbf{d}$ -quasilinear resolution.*

The proof of Theorem 1.7 is based in part on a spectral sequence argument that relates the Betti numbers of $M_{\geq \mathbf{d}}$ to the Fourier–Mukai transform of $\widetilde{M}$ with Beilinson’s resolution of the diagonal as the kernel. Recent breakthroughs [HHL24, BE24] understanding resolutions of the diagonal on arbitrary toric varieties mean that there is hope one may be able to generalize the above argument to arbitrary toric varieties. With this in mind, I am interested in pursuing the following question

1.3 Asymptotic Multigraded Regularity of Ideals Building upon our work discussed above – and inspired by previous work on $\mathbb{P}^r$ [BEL91, CHT99, Kod00] – my collaborators and I also studied the asymptotic regularity of powers of ideals on arbitrary toric varieties. In particular, Definition 1.5 can be extended to all toric varieties by letting S be Cox ring of the toric variety X , replacing $\mathbb{Z}^r$ with the Picard group of X , and replacing $\mathbb{N}^r$ with the nef cone of X . My collaborators and I show that the multigraded regularity of powers of ideals is bounded and translates in a predictable way. In particular, the regularity of I^t essentially translates within $\text{Nef } X$ in fixed directions at a linear rate.

Theorem 1.8. [BHS22, Theorem 4.1] *There exists a degree $\mathbf{a} \in \text{Pic } X$, depending only on I , such that for each integer $t > 0$ and each pair of degrees $\mathbf{q}_1, \mathbf{q}_2 \in \text{Pic } X$ satisfying $\mathbf{q}_1 \geq \deg f_i \geq \mathbf{q}_2$ for all generators f_i of I , we have*

$$t\mathbf{q}_1 + \mathbf{a} + \text{reg } S \subseteq \text{reg}(I^t) \subseteq t\mathbf{q}_2 + \text{Nef } X.$$

1.3.1 Syzygies via Highly Distributed Computing It is quite difficult to compute examples of syzygies. For example, until recently the syzygies of the projective plane embedded by the d -uple Veronese embedding were only known for $d \leq 5$. My co-authors and I exploited recent advances in numerical linear algebra and high-throughput high-performance computing to generate a number of new examples of Veronese syzygies. A follow-up project used similar computational approaches to compute the syzygies of $\mathbb{P}^1 \times \mathbb{P}^1$ in over 200 new examples. This data provided support for several existing conjectures, as well as led us to make a number of new conjectures [?, BCE⁺22]. The resulting data is publicly available via the website SyzygyData and a package for Macaulay2 [BE19, ?].

1.4 Top-Weight Cohomology of $\mathcal{A}_g$ The moduli space of (principally polarized) abelian varieties of dimension g , is a smooth variety $\mathcal{A}_g$ (truthfully a smooth Deligne–Mumford stack) whose points are in one to one correspondence with isomorphism classes of principally polarized abelian varieties of dimension g . Concretely, we may view it as the quotient $[\mathbb{H}_g/\text{Sp}(2g, \mathbb{Z})]$ where $\mathbb{H}_g$ is the Siegel upper half-space. Similar to the moduli space of curves $\mathcal{A}_g$ has long been studied, but much remains unknown about its geometry. For example, the (singular) cohomology of $\mathcal{A}_g$ is only fully known for $g \leq 4$ [?, ?, ?, ?]. Further, Grushevsky [?] posed the question whether $H^{2k+1}(\mathcal{A}_g; \mathbb{Q}) \neq 0$ for some g and k . Building upon the work of Chan, Galatius, and Payne, my co-authors and I developed new methods for understanding a certain canonical quotient of the cohomology of $\mathcal{A}_g$. In particular, our results construct non-trivial cohomology classes in $H^k(\mathcal{A}_g; \mathbb{Q})$ in a number of new cases and answer Grushevsky’s question.

Theorem 1.9. [BBC⁺24, Theorem A] *The rational cohomology $H^k(\mathcal{A}_g; \mathbb{Q}) \neq 0$ for:*

$$(g, k) = (5, 15), (5, 20), (6, 30), (7, 28), (7, 33), (7, 37), \text{ and } (7, 42).$$

For broader context, since $\mathcal{A}_g$ is a rational classifying space for $\text{Sp}_{2g}(\mathbb{Z})$ there is natural isomorphism $H^*(\mathcal{A}_g; \mathbb{Q}) \cong H^*(\text{Sp}_{2g}(\mathbb{Z}); \mathbb{Q})$. In particular, the above results provide new non-vanishing results for $H^*(\text{Sp}_{2g}(\mathbb{Z}); \mathbb{Q})$. However, my work takes advantage of the fact that since $\mathcal{A}_g$ is a smooth

and separated Deligne Mumford stack with a coarse moduli space which is an algebraic variety, permitting Deligne’s mixed Hodge theory to be applied to study the rational cohomology of these groups. In this way we deduce Theorem 1.9 above as a corollary to computing the top-weight cohomology of $\mathcal{A}_g$ for all $g \leq 7$.

1.5 Broader Impacts from Prior NSF Support

1.5.1 Organizing I organized 11 conferences: *Math Careers Beyond Academia* (50 participants), *M2@UW* (45 participants), *Geometry & Arithmetic of Surfaces* (40 participants), *CAZoom* (70 participants), *WAGS* (100 participants), *Gems of Combinatorics I & II* (40 & 30 participants), *Spec($\mathbb{Q}$)* (50 participants), *BATMOBILE (2022)* (20 participants), *Gems of Commutative Algebra I & II* (35 & 40 participants). In this time I organized three special sessions at AMS Sectional Meetings and the Joint Math Meetings. Many of these workshops, namely the *Gems of* series, focused on supporting the next generation of researchers by building community and networks.

1.5.2 Math Circles From 2015 through 2018 I was heavily involved in the Madison Math Circle, including 2 years as the lead organizer. As an organizer, I worked to build stronger connections between the Madison Math Circle, local schools, teachers, and other outreach organizations. This led to weekly attendance to more than double. I also created a new outreach arm of the circle, which visits high schools around the state of Wisconsin to better serve students. While a post-doc at UC Berkeley I volunteered as a speaker with the Berkeley Math Circle.

1.5.3 Mentoring I have actively sought out ways to mentor undergraduate and graduate students, especially those from generally underrepresented groups. While a graduate student, I led reading courses with three undergraduates. One of these students, an undergraduate woman, worked with me for over a year, during which time I helped her apply for summer research projects. Working with *Girls’ Math Night Out* I lead two girls in high school through a project exploring cryptography. During 2018-2019, I mentored 6 first-year graduate students (all women or non-binary students). I also mentored two undergraduate women via the AWM’s Mentoring Network.

I advised two summer research projects for undergraduate students. The first of these projects ran virtually during Summer 2021 when 6 undergraduates worked on combinatorial questions related to my work in [BBC⁺24]. In Summer 2022 I advised an undergraduate student on a research project related to my work on syzygies discussed in Section 1.2. This student is now a graduate student in mathematics and received an NSF GRF. As a postdoc, I began research projects with three graduate students (a majority of whom identify with a generally underrepresented group). These projects have resulted in two pre-prints, with additional projects still ongoing. Throughout the Spring and Summer of 2022, I did a reading course with a first-year graduate woman on algebraic geometry.

1.5.4 Virtual Mathematics In response to the COVID-19 pandemic and the shift of many mathematical activities to virtual formats, I worked to find ways for these online activities to reach those often at the periphery. During Summer and Fall 2020, I helped with Ravi Vakil’s *Algebraic Geometry in the Time of Covid* project. This massive online open-access course in algebraic geometry brought together $\sim 2,000$ participants from around the world. In Spring 2021, I organized an 8-week virtual reading course for undergraduates in algebraic geometry and commutative algebra.

2. The Geometry of Multigraded Syzygies on Stacks

2.1 N_p -Conditions for Stacky Curves

Research Goal 2.1. *Understand the syzygies of stacky curves.*

Research Problem 2.2. Let $\mathcal{X}$ be a stacky curve of genus 0 and 1. If $\mathcal{L}$ is a line bundle on $\mathcal{X}$ and $\deg(\mathcal{X}) \gg 0$ what N_p conditions does $R(\mathcal{X} : \mathcal{L})$ satisfy?

2.2 Green's Conjecture for Canonical Stacky Curves

Research Goal 2.3. Find and prove an appropriate generalization of Green's conjecture describing the vanishing of the syzygies of the canonical rings of stacky curves.

Theorem 2.4. [?]*Theorem 8.4.1 Let $\mathcal{X}$ be stacky curve whose coarse space has genus ≥ 1 and let e be the maximum order of the stabilizer groups of the stacky points; then the canonical ring $R(\mathcal{X}; K_{\mathcal{X}})$ is generated by elements in degree at most $3e$ with relations in degree at most $6e$.

Research Problem 2.5. Can Theorem 2.4 be refined by considering the existence of certain stacky or weighted linear series on $\mathcal{X}$?

Research Problem 2.6. If $\mathcal{X}$ is a general stacky curve is the stacky Brill-Noether locus $\mathcal{W}_d^r(\mathcal{X})$ of expected dimension, smooth away from $\mathcal{W}_d^{r+1}(\mathcal{X})$, and irreducible when of positive dimension?

Lemma 2.7. If $\mathcal{X}$ is a stacky curve then the canonical ring $R(\mathcal{X}; K_{\mathcal{X}})$ is Gorenstein.

Research Problem 2.8. Let $\mathcal{X}$ be a stacky whose coarse space has genus g . For what k does the canonical ring $R(\mathcal{X}; K_{\mathcal{X}})$ satisfy begin k -Koszul in the sense above.

The proof of Theorem 2.4 builds upon the ideas of Schreyer [?] to show the existence of a Gröbner basis for the canonical ring of a stacky curve whose elements have certain degrees. However, Voight and Zureick-Brown do not explicitly construct such Gröbner bases. It should be possible to make their arguments explicit for simple stacky curves. With this in mind, one approach toward Problem ?? is to make Voight and Zureick-Brown's arguments explicit for a large number of examples, and then use a computer algebra system like Macaulay2 to compute the minimal graded free resolutions for these examples.

Research Problem 2.9. Compute the minimal graded free resolution of the canonical ring $R(\mathcal{X}, K_{\mathcal{X}})$ for stacky curves $\mathcal{X}$ where:

- the coarse space of $\mathcal{X}$ has genus ≤ 4 ,
- there are at most 4 stacky points, and
- the stabilizers of the stacky points have order at most 5.

Building upon code generously shared by Voight and Zureick-Brown, I have carried out Problem 2.9 for most of the genus zero examples. In many of these computed cases, and as noted in [?]*Appendix, the canonical ring is a (weighted) complete intersection. However, there are examples showing that even for small examples the canonical ring of a stacky curve is quite complicated.

Example 2.10. Let $\mathcal{X}$ be a stacky curve with three stacky points whose stabilizers are $\mathbb{Z}/4\mathbb{Z}$, $\mathbb{Z}/4\mathbb{Z}$, and $\mathbb{Z}/5\mathbb{Z}$ whose coarse space has genus zero. The canonical ring $R(\mathcal{X}; K_{\mathcal{X}})$ can be presented as S/I where $S = \mathbb{C}[x_1, x_2, x_3, x_4, x_5]$ where $\deg(x_1) = 3$, $\deg(x_2) = \deg(x_3) = 4$, and $\deg(x_4) = \deg(x_5) = 5$ and I is an ideal generated by 7 homogenous polynomials of degrees 8, 8, 9, 9, 10, 13, 13. The minimal graded free resolution of $R(\mathcal{X}; K_{\mathcal{X}})$ is shown below (with the degrees suppressed for brevity):

$$0 \longleftarrow R(\mathcal{X}; K_{\mathcal{X}}) \longleftarrow S \longleftarrow S^{\oplus 7} \longleftarrow S^{\oplus 20} \longleftarrow S^{\oplus 30} \longleftarrow S^{\oplus 22} \longleftarrow S^{\oplus 6} \longleftarrow 0.$$

The approach of using large-scale computations to find conjectures concerning syzygies is very much in the same spirit as my prior work on the syzygies of surfaces [?, BCE⁺22]. While the precise techniques need to address Problem 2.9 will likely be different much of the general framework of organizing, distributing, and publicizing such computations will likely be similar.

2.3 Asymptotic Syzygies of Weighted Projective Stacks

Research Problem 2.11. Let $\mathcal{X}$ be a smooth DM stack, $\dim \mathcal{X} \geq 2$, and fix an index $1 \leq q \leq \dim \mathcal{X}$. If $(L_d)_{d \in \mathbb{N}}$ is a sequence of very ample line bundles such that $L_{d+1} - L_d$ is constant and ample compute the limit (if it exists):

$$\lim_{d \rightarrow \infty} \rho_q(\mathcal{X}; L_d) = \lim_{d \rightarrow \infty} \frac{\#\{p \in \mathbb{N} \mid \beta_{p,p+q}(X, L_d) \neq 0\}}{n_d}.$$

Research Problem 2.12. Fix a positive integer $a > 1$. For what values of p does the weighted projective stacky plane $\mathcal{P}(1, 1, 1, a)$ embedded by $\mathcal{O}(d)$ satisfy N_p .

2.4 Computations with Toric Stacks

Research Problem 2.13. Develop a robust software package for Macaulay2 allowing for computations with toric stacks.

Research Problem 2.14. Develop a robust software package for Macaulay2 allowing for computations with toric stacks.

Research Problem 2.15. Compute the graded Betti tables of $\mathcal{P}(1, 1, 1, a)$ embedded by $\mathcal{O}(d)$ for all $1 < a \leq 15$ and all $2 \leq d \leq 6$ by adapting the high-performance computing approaches developed in  Juliette: [cite]

3. Castelnuovo–Mumford Regularity: Resolutions, Bounds, and Asymptotics

3.1 Asymptotic Regularity of Powers of Ideals and Ideal Sheaves

Research Goal 3.1. Let X be a smooth projective toric variety with S its $\text{Pic}(X)$ -graded Cox ring. Understand the asymptotic behavior of $\text{reg}(I^p)$ and $\text{reg}(\mathcal{I}^p)$ where $I \subset S$ is an $\mathbb{Z}^r$ -graded ideal and $\mathcal{I} \subset \mathcal{O}_X$ is an ideal sheaf.

Conjecture 3.2. Let X be a smooth projective toric variety and $\mathcal{I} \subset \mathcal{O}_X$ an ideal sheaf. With the notation as above:

$$\lim_{p \rightarrow \infty} \frac{\text{reg}(\mathcal{I}^p)_{\mathbb{R}}}{p} = \left\{ L \in \text{Pic}(X)_{\mathbb{R}} \mid \pi^*(L) - E \text{ is nef on } \tilde{X} \right\}.$$

Research Problem 3.3. Prove Conjecture 3.2.

Research Problem 3.4. Is the Seshadri regions defined above be determined by the Newton–Okunokov body of some flag on X .

3.2 Bounds on Multigraded Regularity

Research Problem 3.5. Let X be a smooth projective toric variety with S its $\text{Pic}(X)$ -graded Cox ring. If $I \subset S$ is a $\mathbb{Z}^r$ -graded ideal bound the number of minimal generators of $\text{reg}(I^p)$ as a function of p in terms of invariants of I ?

Research Problem 3.6. Let X be a smooth projective toric variety and $Z \subset X$ a smooth subvariety. Can one bound the multigraded regularity of I_Z in terms of the codimension of $Z \subset X$ and the degrees of minimal generators of I_Z ?

Research Problem 3.7. Compute the multigraded regularity of torus invariant subvarieties of X in terms of the combinatorics of the cone defining them.

Research Problem 3.8. Compute the multigraded Castelunovo–Mumford Regularity of $\overline{\mathcal{M}}_{0,n} \subset ??$

Research Problem 3.9. Compute the minimal graded free resultion of the defining ideal I of $\overline{\mathcal{M}}_{0,n} \subset ??$

3.3 A Generalized Eisenbud–Goto Theorem for Toric Varieties

Research Problem 3.10. *Let X be a smooth projective toric variety with S its $\text{Pic}(X)$ -graded Cox ring and B its irrelevant ideal. Let M be a $\text{Pic}(X)$ -graded S -module. Find the correction generalization of d -quasi-linear resolution such that a $\text{Pic}(X)$ -graded S -module M is d -regular if and only if the truncation $M_{\geq d}$ has a d -quasilinear resolution.*

Conjecture 3.11. *Let S be the $\mathbb{Z}^2$ -graded Cox ring of the Hirzebruch surface H_t . If M is a $\mathbb{Z}^2$ -graded S -module then $M_{\geq d}$ has a d -linear resolution if and only if $d + \text{reg}(S) \subset \text{reg}(M)$.*

Research Problem 3.12. *Prove Conjecture 3.11*

Research Problem 3.13. *Let X be a smooth projective toric variety with S its $\text{Pic}(X)$ -graded Cox ring and B its irrelevant ideal. Let M be a $\text{Pic}(X)$ -graded S -module. Find a characterization for when the truncation $M_{\geq d}$ for $d \in \text{Pic}(X)$ has a linear resolution in terms of vanishing of the local cohomology groups $H_B^i(M)_t$.*

Research Problem 3.14. *Find an algorithm to compute a minimal generating set for $\text{reg}(M)$.*

Research Problem 3.15. *Let X be a smooth projective toric variety with S its $\text{Pic}(X)$ -graded Cox ring and B its irrelevant ideal. Let M be a $\text{Pic}(X)$ -graded S -module. Does the set of (multigraded) Betti numbers of all virtual resolutions of M determine the multigraded regularity of M ?*

4. Uniform Homological Algebra on Toric Varieties

4.1 Regularity on the D_{Cox} Category

Research Problem 4.1. *Develop a theory of Castelunovo–Mumford regularity on the category D_{Cox} the restrict to the classical notion of regularity X_i .*

Research Problem 4.2. *Let $I \subset S$ be a $\mathbb{Z}^r$ -graded homogenous ideal. Let X and X' be toric varieties corresponding to two chambers $\Gamma, \Gamma' \subset \Sigma_{GKZ}$ that share a wall, with $B, B' \subset S$ the respective irrelevant ideals. Prove there exists a “small” subset $F \subset \mathbb{Z}^r$ depending only on X and X' such that:*

$$\text{reg}_X((J :_S B^\infty)) + F = \text{reg}_{X'}((J :_S B'^\infty)) + F.$$

Research Problem 4.3. *With the set up as above prove the minimal graded free S -resolutions of $(J :_S B^\infty)$ and $(J :_S B'^\infty)$ agree in small degree.*

5. Broader Impacts

5.1 Organizing As discussed previously, I have organized 10+ national/international conferences, and I plan to continue organizing conferences at roughly this pace. Currently, I am organizing two research conferences in algebraic geometry *Algebraic Geometry Northeastern Series (AGNES)* and *BATMOBILE* both occurring in late 2024. I am also in the early stages of organizing an international workshop with Diane Maclagan on multigraded homological algebra and toric geometry, which I hope will occur in 2025. Tyler Kelly, Mike Hill, and I are also in the planning stages of organizing a satellite conference for LGBTQ+ mathematicians for the 2026 ICM in Philadelphia.

The “GEMS” workshop series that I co-founded sought to build a diverse community of mathematicians to address gender equity in the mathematical community from new perspectives. In particular, these conferences are indented to supplement the affinity group model of the *Women in...* conference series by promoting broad conversations and networks. Going forward I am interested in expanding these “GEMS” workshops to other areas of mathematics and creating cross-field discussions that broaden the standard notion of gender equity in mathematics. In particular, there is a planned follow-up to *GEMS in Commutative Algebra* scheduled for Fall 2025, and I am tentatively

planning a version of *GEMS in Algebraic Geometry* for 2026. As more of these conferences occur, I have plans to try to work with mathematicians in other fields to organize their own version of *GEMS in...* conferences in their fields.

When organizing these conferences I paid particular attention to making them as inclusive of women and non-binary researchers as possible. For example, I designed the registration form to be thoughtful of the concerns of transgender researchers and highlighted the locations of single occupancy and ADA-compliant restrooms. The importance of such efforts was highlighted by the following comment I received from a participant, “I just wanted to thank you for making this workshop inclusive for people with all gender identifications. ... I have always felt out of place when I participated in conferences/workshops for women when they do not specify that non-binary people are welcome ... I really appreciate those questions you put in the registration form. It means a lot to me.” I would like to work with various professional societies (AMS, AWM, NAM, Spectra, etc.) and research institutes (SLMath, ICERM, IAS, etc.) to produce a guide of best practices for organizing welcoming and inclusive mathematics conferences. By doing this I hope to make it easier for others to organize welcoming events.

5.2 National & International Advocacy. I have worked with the Executive Committee of the *Association for Women in Mathematics* to consider ways they could expand their support of women and non-binary mathematicians. I hope to continue this work and find further ways the AWM can support both women and non-binary mathematicians. Since Winter 2023 I joined SLMath’s (formerly MSRI’s) *Committee on Women in Mathematics*. I plan to remain on this committee until 2028 when my term is set to expire. In this role, I am hoping to organize a yearly retreat at SLMath that brings together small groups of 5-8 women (and underrepresented individuals more generally) who have been implementing innovative ideas promoting equity in mathematics and brainstorm ways to scale their efforts and effect radical positive change for women in mathematics.

5.3 A More Inclusive Learning Community. Going forward I am excited to continue my work supporting LGBTQ+ students, and would love to continue building organizations to do so. In particular, given the amazing successes of programs like MSRI-UP and the EDGE Program, I would love to organize a summer REU program specifically aimed at supporting and promoting LGBTQ+ mathematicians. Further, I am in the early stages of planning a mentorship program to help guide LGBTQ+ undergraduates through the process of applying to graduate programs in mathematics and helping young LGBTQ+ graduate students establish themselves. The plan would be to break participants into groups with each group having LGBTQ+ mathematicians at various career stages, thus allowing participants to exchange advice, find support, and build mentoring networks.

5.4 Mentoring. As a new faculty member at Dartmouth College I hope to continue and expand the mentoring I do with students, continuing a focus on mentoring students from underrepresented groups. Since arriving on campus in the late Summer I have begun mentoring one graduate student and one undergraduate woman. While still early it seems likely that I will end up co-advising the graduate student and advising the undergraduate’s senior thesis. I am also working with the college’s AWM Student Chapter and the Out in STEM Student Chapter.

References

- [ABLS20] Ayah Almousa, Juliette Bruce, Michael Loper, and Mahrud Sayrafi. The virtual resolutions package for Macaulay2. *J. Softw. Algebra Geom.*, 10(1):51–60, 2020.
- [BBB⁺17] Madeline Brandt, Juliette Bruce, Taylor Brysiewicz, Robert Krone, and Elina Robeva. The degree of $\mathrm{SO}(n, \mathbb{C})$. In *Combinatorial algebraic geometry*, volume 80 of *Fields Inst. Commun.*, pages 229–246. Fields Inst. Res. Math. Sci., Toronto, ON, 2017.
- [BBB21] Anthony Bonato, Juliette Bruce, and Ron Buckmire. Spaces for all: the rise of LGBTQ+ mathematics conferences. *Notices Amer. Math. Soc.*, 68(6):998–1003, 2021.
- [BBC23] Madeline Brandt, Juliette Bruce, and Daniel Corey. The virtual euler characteristic for binary matroids, 2023.
- [BBC⁺24] Madeline Brandt, Juliette Bruce, Melody Chan, Margarida Melo, Gwyneth Moreland, and Corey Wolfe. On the top-weight rational cohomology of $\mathcal{A}_g$. *Geom. Topol.*, 28(2):497–538, 2024.
- [BCE⁺22] Juliette Bruce, Daniel Corey, Daniel Erman, Steve Goldstein, Robert P. Laudone, and Jay Yang. Syzygies of $\mathbb{P}^1 \times \mathbb{P}^1$: data and conjectures. *J. Algebra*, 593:589–621, 2022.
- [BE19] Juliette Bruce and Daniel Erman. A probabilistic approach to systems of parameters and Noether normalization. *Algebra Number Theory*, 13(9):2081–2102, 2019.
- [BE24] Michael K. Brown and Daniel Erman. A short proof of the Hanlon-Hicks-Lazarev theorem. *Forum Math. Sigma*, 12:Paper No. e56, 6, 2024.
- [BEGY20] Juliette Bruce, Daniel Erman, Steve Goldstein, and Jay Yang. Conjectures and computations about Veronese syzygies. *Exp. Math.*, 29(4):398–413, 2020.
- [BEGY21] Juliette Bruce, Daniel Erman, Steve Goldstein, and Jay Yang. The Schur-Veronese package in Macaulay2. *J. Softw. Algebra Geom.*, 11(1):83–87, 2021.
- [BEL91] Aaron Bertram, Lawrence Ein, and Robert Lazarsfeld. Vanishing theorems, a theorem of Severi, and the equations defining projective varieties. *J. Amer. Math. Soc.*, 4(3):587–602, 1991.
- [BHS22] Juliette Bruce, Lauren Cranton Heller, and Mahrud Sayrafi. Bounds on multigraded regularity, 2022.
- [BHS24] Juliette Bruce, Lauren Cranton Heller, and Mahrud Sayrafi. Characterizing multigraded regularity and virtual resolutions on products of projective spaces, 2024.
- [BL20] Juliette Bruce and Wanlin Li. Effective bounds on the dimensions of Jacobians covering abelian varieties. *Proc. Amer. Math. Soc.*, 148(2):535–551, 2020.
- [Bru22a] Juliette Bruce. The quantitative behavior of asymptotic syzygies for Hirzebruch surfaces. *J. Commut. Algebra*, 14(1):19–26, 2022.
- [Bru22b] Juliette Bruce. A word from... juliette bruce, inaugural president of spectra. *Notices Amer. Math. Soc.*, 69(6):898–899, 2022.
- [Bru24] Juliette Bruce. Asymptotic syzygies for products of projective spaces. *J. Algebra*, 649:347–391, 2024.
- [CHT99] S. Dale Cutkosky, Jürgen Herzog, and Ngô Việt Trung. Asymptotic behaviour of the Castelnuovo-Mumford regularity. *Compositio Math.*, 118(3):243–261, 1999.
- [EG84] David Eisenbud and Shiro Goto. Linear free resolutions and minimal multiplicity. *J. Algebra*, 88(1):89–133, 1984.
- [HHL24] Andrew Hanlon, Jeff Hicks, and Oleg Lazarev. Resolutions of toric subvarieties by line bundles and applications, 2024.
- [Kod00] Vijay Kodiyalam. Asymptotic behaviour of Castelnuovo-Mumford regularity. *Proc. Amer. Math. Soc.*, 128(2):407–411, 2000.
- [MS04] Diane Maclagan and Gregory G. Smith. Multigraded Castelnuovo-Mumford regularity. *J. Reine Angew. Math.*, 571:179–212, 2004.