

Classificação de psicofármacos

Antidepressivos Tricíclicos (ADT)

- Imipramina (Tofranil®)
- Clomipramina (Anafranil®, Clo®)
- Amitriptilina (Trytanol®, Amytril®, Limbitrol®, Protanol®)
- Nortriptilina (Pamelor®)
- Maprotilina (Ludiomil®)

Inibidores Seletivos da Recaptação de Serotonina (ISRS)

- Fluoxetina (Prozac®, Daforin®, Verotina®, Fluxene®, Prozen®)
- Sertralina (Zoloft®, Serenata®, Assert®, Tolrest®, Dieloft®)
- Paroxetina (Aropax®, Pondera®, Paxil®, Celebrin®, Paxan®, Paxtrat®)
- Citalopram (Cipramil®, Procimax®, Citta®, Denyl®, Maxapran®)
- Escitalopram (Lexapro®, Reconter®, Exodus®, Espran®, Esc®, Sedopan®, Unitram®)
- Fluvoxamina (Luvox®, Revoc®)

Inibidores Seletivos da Recaptação de Serotonina e Noradrenalina (ISRSN)

- Venlafaxina (Efexor®, Venlift®, Venlaxin®, Alenthus®)
- Desvenlafaxina (Pristiq®, Desve®, Desduo®, Desventag®, Deller®, Elifore®, Aviv®, Vellana®, Andes®, Imense®)
- Duloxetina (Cymbalta®, Velija®, Neulox®)

Inibidores da Recaptação de Noradrenalina e Dopamina

- Bupropiona (Welbutrin®, Zyban®, Bup®, Zetron®, Noradop®, Bupium®)

Inibidores Seletivos da Recaptação de Noradrenalina (ISRN)

- Reboxetina (Prolift®)
- Atomoxetina (Atentah®)

Inibidores da Monoaminoxidase (IMAO)

- Tranilcipromina (Parnate®)
- Moclobemida (Aurorix®)

Antidepressivos atípicos

- Trazodona (Donaren®, Sonic®)
- Mirtazapina (Remeron®, Menelat®, Razapina®)
- Agomelatina (Valdoxan®)
- Vortioxetina (Voextor®, Brintellix®, Virtuoso®)
- Vilazodona
- Esketamina / Escetamina

Antipsicóticos típicos

- Clorpromazina (Amplictil®, Longactil®)
- Levomepromazina (Neozine®, Levozine®, Meprozin®)
- Flufenazina (Anatensoll®, Flufenan®)
- Haloperidol (Haldol®)

- Tioridazina (Melleril®, Unitidazin®)
- Pimozida (Orap®)
- Periciazina (Neuleptil®)

Antipsicóticos atípicos

- Risperidona (Risperdal®, Riss®, Rispidon®, Zargus®, Risperidon®)
- Olanzapina (Zyprexa®, Axoniun®, Neupine®, Zap®, Zopix®)
- Quetiapina (Seroquel®, Neutiapim®, Quetiapin®, Quetros®, Queropax®)
- Ziprasidona (Geodon®)
- Aripiprazol (Abilify®, Aristab®, Arpejo®)
- Sulpirida (Equilid®, Dogmatil®, Sulpan®)
- Amisulprida (Socian®)
- Asenapina (Saphris®)
- Clozapina (Leponex®, Pinazan®)
- Lurasidona (Latuda®)
- Brexpiprazol (Rexulti®)

Estabilizadores de Humor

- Lítio (Carbolitium®, Literata®)
- Ácido Vapóico / Valproato / Divalproato de sódio (Divalcon®, Valpi®, Depakene®, Depakine®/Depakote®, Torval CR®)
- Carbamazepina (Tegretol®)
- Oxcarbazepina (Trileptal®, Oxcarb®, Oleptal®)
- Lamotrigina (Lamictal®, Lamitor®)
- Topiramato (Topamax®, Amato®, Égide®)

Benzodiazepínicos

- Diazepam (Valium®, Somaplast®, Ansilive®, Dienpax®)
- Clonazepam (Rivotril®, Clonotril®, Clonam®, Epileptil®)
- Alprazolam (Frontal®, Apraz®, Tranquinal®, Altrox®)
- Bromazepam (Lexotan®, Somalium®, Brozepax®, Nervium®, Novazepam®)
- Lorazepam (Lorax®, Lorium®, Ansirax®, Lorapan®)
- Clobazam (Frisium®, Urbanil®)
- Cloxazolam (Olcadil®, Elum®, Anoxolan®)
- Nitrazepam (Sonebon®, Nitrazepol®, Nitrapan®)
- Flunitrazepam (Rohypnol®, Rohydorm®)
- Flurazepam (Dalmadorm®)
- Midazolam (Dormonid®, Dormire®, Dormium®, Hipnazolam®)
- Clordiazepóxido (Psicosedin®)

Antagonistas opióides

- Naloxona
- Naltrexona (Uninaltrex®)

Psicoestimulantes

- Metilfenidato (Ritalina®, Medato®, Concerta®, Ragione®, FOQ XR®)
- Lisdexanfetamina (Venvanse®, Juneve®)
- Modafinila/Modafinil (Stavigile®)

Anticonvulsivantes

- Pregabalina (Lyrica®, Konduz®, Dorene®)
- Lacosamida
- Gabapentina
- Levetiracetam

Azapironas

- Buspirona (Ansitec®)

Hipnóticos Não Benzodiazepínicos

- Zolpidem (Prysmá®, Stilnox®)
- Eszopiclona
- Ramelteona (Rozerem®)

Agradecemos o psiquiatra Danilo Silveira Seabra, que revisou e colaborou neste trabalho.

Referências

BAES, Cristiane von Werne; JURUENA, Mário Francisco. Psicofarmacoterapia para o clínico geral. *Revista de Medicina*, Ribeirão Preto, v. 50, supl. 1, p. 22-36, 2017. Disponível em: <https://doi.org/10.11606/issn.2176-7262.v50isupl1.p22-36>.

GOZZANI, Judymara Lauzi. Opióides e antagonistas. *Revista Brasileira de Anestesiologia*, v. 44, n. 1, p. 65-73, 1994.