

Individual Assingment 5 JM

June McNally

2026-05-10

Set Up

Packages

```
# geberal and cleaning
library(tidyverse)
library(here)
library(janitor)
library(snakecase)

# wrapping axis labels
library(scales)

# reading in .xlsx files
library(readxl)

# visualizing missing data
library(naniar)

# arranging plots
library(patchwork)
```

Data

```
# taxonomic information
taxon_list <- read_csv(here("data", "taxon_list.csv"))
```

```

# aquatic invertebrates
aq_ins <- read_xlsx(here("data", "Aquatic Sampling Data-2026-03-10.xlsx"),
                    sheet = "Aquatic Insects")

# site information
sites <- read_xlsx(here("data", "Aquatic Sampling Data-2026-03-10.xlsx"),
                   sheet = "sites")

# water quality information
water_quality <- read_xlsx(here("data", "Aquatic Sampling Data-2026-03-10.xlsx"),
                           sheet = "Water Quality",
                           na = c("", "n/a", "N/A", "over", "173+"))

```

Class code

```

# creating new object from sites
ncos_sites <- sites |>
  # filter to only include NCOS sites sampled four times a year
  filter(sector == "NCOS" & sampling_frequency == "Quarterly")

```

```

# creating new clean object from taxon_list
taxon_list_clean <- taxon_list |>
  # converting taxon name to snake case (no spaces or capital letters,
  # only underscores)
  mutate(verbatim_name = to_any_case(verbatim_name, case = "snake"))

```

```

# creating new clean object from water quality
water_quality_clean <- water_quality |>
  # clean column names
  clean_names() |>
  # filtering join: only include sites in ncos_sites
  semi_join(ncos_sites, by = "site") |>
  # create new column to join with later data frames
  unite("date_site", date, site, remove = TRUE) |>
  # group by date_site column
  group_by(date_site) |>
  # calculate median pH, salinity, DO
  summarize(
    med_ph = median(p_h, na.rm = TRUE),
    med_sal = median(salinity_ppt, na.rm = TRUE),
    med_do = median(dissolved_oxygen_mg_l, na.rm = TRUE)) |>

```

```

# ungroup data frame
ungroup() |>
# filter out salinity outliers
filter(date_site != "2022-08-12_NVBR")

# creating new clean object from aquatic inverts
aq_ins_clean <- aq_ins |>
# clean column names
clean_names() |>
# filtering join: only include sites occurring in ncos_sites
semi_join(ncos_sites, by = c("site")) |>
# renaming columns
rename(date = date_on_vial) |>
# select columns of interest
select(date, sample_type, site,
        ostracod,
        copepod,
        hemiptera_corixidae_boatman,
        cladocera,
        nematode,
        diptera_ceratopogonidae,
        annelida_oligochaete,
        diptera_chironomid,
        annelida_polychaete) |>
# filter to only include "filtered beaker" samples
filter(sample_type %in% c("FB 250", "FB250")) |>
# convert the data frame to long format
pivot_longer(cols = ostracod:annelida_polychaete,
              names_to = "taxon",
              values_to = "abundance") |>
# group by date, site, taxon
group_by(date, site, taxon) |>
# calculate average abundance per liter (sum abundance divided by 7.5 liters)
summarize(ave_lit = sum(abundance, na.rm = TRUE)/7.5) |>
# ungroup data frame
ungroup() |>
# create a new column called `date_site` to join with water quality
unite("date_site", date, site, remove = FALSE) |>
# join with water quality
left_join(water_quality_clean, by = c("date_site")) |>
# join with taxon list
left_join(taxon_list_clean, by = c("taxon" = "verbatim_name")) |>

```

```

# add full names of sites
mutate(site_full = case_when(
  site == "NVBR" ~ "North of Venoco Bridge",
  site == "NEC" ~ "South of Venoco Bridge",
  site == "NMC" ~ "Main Channel",
  site == "NPB" ~ "South of Phelps Bridge",
  site == "NPB1" ~ "North of Phelps Bridge",
  site == "NPB2" ~ "Phelps Road",
  site == "NWP" ~ "West Pond",
  site == "NDC" ~ "Devereux Creek"
)) |>
# setting factor levels for site
mutate(site_full = fct_reorder(site_full, med_sal, .fun = "median", .na_rm = TRUE),
  site_full = fct_rev(site_full))

```