

## INTRODUCTION

In cylindrical coordinates, we have

$$\begin{pmatrix} \eta_r \\ \eta_\phi \end{pmatrix} = \begin{pmatrix} \cos \phi & \sin \phi \\ -\sin \phi & \cos \phi \end{pmatrix} \begin{pmatrix} \eta_x \\ \eta_y \end{pmatrix} \quad (1)$$

and

$$\begin{pmatrix} \partial_r \\ \frac{1}{r}\partial_\phi \end{pmatrix} = \begin{pmatrix} \cos \phi & \sin \phi \\ -\sin \phi & \cos \phi \end{pmatrix} \begin{pmatrix} \partial_x \\ \partial_y \end{pmatrix} \quad (2)$$

In bulk, for the two time-reversed ground states,

$$(\eta_x^{\text{bulk}}, \eta_y^{\text{bulk}}) = \eta_0(1, \pm i) \quad (3)$$

$$(\eta_r^{\text{bulk}}, \eta_\phi^{\text{bulk}}) = e^{\pm i\phi} \eta_0(1, \pm i) \quad (4)$$

In GL theory, the order parameter is determined by minimizing the free energy functional

$$F[\boldsymbol{\eta}] = \int d^3r (f_{\text{bulk}} + f_{\text{grad}}) \quad (5)$$

where

$$f_{\text{bulk}}[\boldsymbol{\eta}] = \alpha \boldsymbol{\eta} \cdot \boldsymbol{\eta}^* + \beta_1 (\boldsymbol{\eta} \cdot \boldsymbol{\eta}^*)^2 + \beta_2 |\boldsymbol{\eta} \cdot \boldsymbol{\eta}|^2, \quad (6)$$

and

$$f_{\text{grad}}^{\text{Car}}[\boldsymbol{\eta}] = \kappa_1 (\partial_i \eta_j) (\partial_i \eta_j)^* + \kappa_2 (\partial_i \eta_i) (\partial_j \eta_j)^* + \kappa_3 (\partial_i \eta_j) (\partial_j \eta_i)^* \quad (7)$$

In cylindrical basis, we have  $\partial_i^{\text{Car}} = R_{ik} \partial_k$  and  $\eta_j^{\text{Car}} = R_{jl} \eta_l$ , where  $R_{ij}$  is the 2d rotational matrix. The form of the bulk free energy term doesn't change. The gradient term becomes

$$\begin{aligned} f_{\text{grad}}^{\text{Cyl}}[\boldsymbol{\eta}] = & \kappa_1 (\partial_i \eta_j) (\partial_i \eta_j)^* + \kappa_2 (\partial_i \eta_i) (\partial_j \eta_j)^* + \kappa_3 (\partial_i \eta_j) (\partial_j \eta_i)^* \\ & + \frac{2}{r} \text{Re} \left\{ \kappa_1 \left( \eta_r^* \frac{\partial_\phi}{r} \eta_\phi - \eta_\phi^* \frac{\partial_\phi}{r} \eta_r \right) + \kappa_2 \eta_r^* \partial_j \eta_j + \kappa_3 \left( \eta_r^* \frac{\partial_\phi}{r} \eta_\phi - \eta_\phi^* \partial_r \eta_\phi \right) \right\} \\ & + \frac{1}{r^2} [\kappa_1 \eta_j^* \eta_j + (\kappa_2 + \kappa_3) \eta_r^* \eta_r]. \end{aligned} \quad (8)$$

Take the  $\kappa_1$  term for example, note that  $R_{ij}R_{ik} = R_{ji}^T R_{ik} = \delta_{jk}$

$$(\partial_i^{\text{Car}} \eta_j^{\text{Car}})(\partial_i^{\text{Car}} \eta_j^{\text{Car}})^* \quad (9)$$

$$= \left[ R_{ik} \partial_k (R_{jl} \eta_l) \right] \left[ R_{im} \partial_m (R_{jn} \eta_n^*) \right] \quad (10)$$

$$= \left[ \partial_k (R_{jl} \eta_l) \right] \left[ \partial_k (R_{jn} \eta_n^*) \right] \quad (11)$$

$$= \left[ (\partial_k R_{jl}) \eta_l + R_{jl} (\partial_k \eta_l) \right] \left[ (\partial_k R_{jn}) \eta_n^* + R_{jn} (\partial_k \eta_n^*) \right] \quad (12)$$

$$= (\partial_k R_{jl}) \eta_l (\partial_k R_{jn}) \eta_n^* + (\partial_k R_{jl}) \eta_l R_{jn} (\partial_k \eta_n^*) + R_{jl} (\partial_k \eta_l) (\partial_k R_{jn}) \eta_n^* + R_{jl} R_{jn} (\partial_k \eta_l) (\partial_k \eta_n^*)$$

Note that  $\partial_r R(\phi) = 0$ ,  $\partial_\phi R(\phi) = R(\phi) R(\pi/2) = R(\pi/2) R(\phi)$ . Finally we have

$$(\partial_i^{\text{Car}} \eta_j^{\text{Car}})(\partial_i^{\text{Car}} \eta_j^{\text{Car}})^* \quad (13)$$

$$= \frac{1}{r^2} \eta_l \eta_l^* + \frac{1}{r} R_{nl}^{\pi/2} \eta_l \left( \frac{\partial_\phi}{r} \eta_n^* \right) + \frac{1}{r} R_{ln}^{\pi/2} \eta_n^* \left( \frac{\partial_\phi}{r} \eta_l \right) + (\partial_k \eta_l) (\partial_k \eta_l)^* \quad (14)$$

$$= \frac{1}{r^2} \eta_l \eta_l^* + \frac{2}{r} \text{Re} \left\{ R_{ln}^{\pi/2} \eta_n^* \left( \frac{\partial_\phi}{r} \eta_l \right) \right\} + (\partial_k \eta_l) (\partial_k \eta_l)^* \quad (15)$$

$$= \frac{1}{r^2} \eta_l \eta_l^* + \frac{2}{r} \text{Re} \left\{ \eta_r^* \frac{\partial_\phi}{r} \eta_\phi - \eta_\phi^* \frac{\partial_\phi}{r} \eta_r \right\} + (\partial_k \eta_l) (\partial_k \eta_l)^* \quad (16)$$

Similarly, for the  $\kappa_2$  terms, we have

$$(\partial_i^{\text{Car}} \eta_i^{\text{Car}})(\partial_j^{\text{Car}} \eta_j^{\text{Car}})^* \quad (17)$$

$$= \left[ R_{ik} \partial_k (R_{il} \eta_l) \right] \left[ R_{jm} \partial_m (R_{jn} \eta_n^*) \right] \quad (18)$$

$$= \left[ R_{ik} (\partial_k R_{il}) \eta_l + R_{ik} R_{il} (\partial_k \eta_l) \right] \left[ R_{jm} (\partial_m R_{jn}) \eta_n^* + R_{jm} R_{jn} (\partial_m \eta_n^*) \right] \quad (19)$$

$$= \left[ \frac{1}{r} \eta_r + \partial_k \eta_k \right] \left[ \frac{1}{r} \eta_r^* + \partial_m \eta_m^* \right] \quad (20)$$

$$= \frac{1}{r^2} \eta_r \eta_r^* + \frac{2}{r} \text{Re} \{ \eta_r^* \partial_k \eta_k \} + (\partial_k \eta_k) (\partial_m \eta_m)^* \quad (21)$$

For the  $\kappa_3$  terms, we have

$$(\partial_i^{\text{Car}} \eta_j^{\text{Car}})(\partial_j^{\text{Car}} \eta_i^{\text{Car}})^* \quad (22)$$

$$= \left[ R_{ik} \partial_k (R_{jl} \eta_l) \right] \left[ R_{jm} \partial_m (R_{in} \eta_n^*) \right] \quad (23)$$

$$= \left[ R_{ik} (\partial_k R_{jl}) \eta_l + R_{ik} R_{jl} (\partial_k \eta_l) \right] \left[ R_{jm} (\partial_m R_{in}) \eta_n^* + R_{jm} R_{in} (\partial_m \eta_n^*) \right] \quad (24)$$

$$= R_{ik} (\partial_k R_{jl}) \eta_l R_{jm} (\partial_m R_{in}) \eta_n^* + R_{ik} (\partial_k R_{jl}) \eta_l R_{jm} R_{in} (\partial_m \eta_n^*) \\ + R_{ik} R_{jl} (\partial_k \eta_l) R_{jm} (\partial_m R_{in}) \eta_n^* + R_{ik} R_{jl} (\partial_k \eta_l) R_{jm} R_{in} (\partial_m \eta_n^*) \quad (25)$$

$$= \frac{1}{r^2} R_{\phi l}^{\pi/2} \eta_l R_{\phi n}^{\pi/2} \eta_n^* + \frac{1}{r} R_{ml}^{\pi/2} \eta_l (\partial_m \eta_\phi^*) + \frac{1}{r} R_{kn}^{\pi/2} \eta_n^* (\partial_k \eta_\phi) + (\partial_k \eta_l) (\partial_l \eta_k)^* \quad (26)$$

$$= \frac{1}{r^2} \eta_r \eta_r^* + \frac{2}{r} \text{Re} \left\{ R_{kn}^{\pi/2} \eta_n^* (\partial_k \eta_\phi) \right\} + (\partial_k \eta_l) (\partial_l \eta_k)^* \quad (27)$$

$$= \frac{1}{r^2} \eta_r \eta_r^* + \frac{2}{r} \text{Re} \left\{ \eta_r^* \frac{\partial_\phi}{r} \eta_\phi - \eta_\phi^* \partial_r \eta_\phi \right\} + (\partial_k \eta_l) (\partial_l \eta_k)^* \quad (28)$$

So the free energy density is

$$f_{\text{bulk}}[\boldsymbol{\eta}] = \alpha \boldsymbol{\eta} \cdot \boldsymbol{\eta}^* + \beta_1 (\boldsymbol{\eta} \cdot \boldsymbol{\eta}^*)^2 + \beta_2 |\boldsymbol{\eta} \cdot \boldsymbol{\eta}|^2, \quad (29)$$

and

$$f_{\text{grad}}^{\text{Cyl}}[\boldsymbol{\eta}] = \kappa_1 (\partial_i \eta_j) (\partial_i \eta_j)^* + \kappa_2 (\partial_i \eta_i) (\partial_j \eta_j)^* + \kappa_3 (\partial_i \eta_j) (\partial_j \eta_i)^* \\ + \frac{2}{r} \text{Re} \left\{ \kappa_1 \left( \eta_r^* \frac{\partial_\phi}{r} \eta_\phi - \eta_\phi^* \frac{\partial_\phi}{r} \eta_r \right) + \kappa_2 \eta_r^* \partial_j \eta_j + \kappa_3 \left( \eta_r^* \frac{\partial_\phi}{r} \eta_\phi - \eta_\phi^* \partial_r \eta_\phi \right) \right\} \\ + \frac{1}{r^2} [\kappa_1 \eta_j^* \eta_j + (\kappa_2 + \kappa_3) \eta_r^* \eta_r]. \quad (30)$$

## EULER-LAGRANGE EQUATIONS

The GL differential equations  $\delta F / \delta \eta_k^* = 0$ .

If we do the functional variation of the free energy in cylindrical basis, we will get

$$\alpha \eta_k + 2(\beta_1 \eta_i \eta_i^* \eta_k + \beta_2 \eta_i^2 \eta_k^*) - (\kappa_1 \partial_i^2 \eta_k + \kappa_2 \partial_k \partial_i \eta_i + \kappa_3 \partial_j \partial_k \eta_j) \\ - \frac{1}{r} \left[ (2\kappa_1 + \kappa_3) \frac{\partial_\phi}{r} (\eta_r \delta_{k\phi} - \eta_\phi \delta_{kr}) + \kappa_1 \partial_r \eta_k + (\kappa_2 + \kappa_3) \partial_k \eta_r \right] \\ + \frac{1}{r^2} [\kappa_1 \eta_k + (\kappa_2 + \kappa_3) \eta_r \delta_{kr}] = 0. \quad (31)$$

For the  $\kappa_1$  term in first line of Eq. (30)

$$F \sim \iint (\partial_i \eta_j)(\partial_i \eta_j)^* r d\phi dr = \text{surface term} - \iint \eta_j^* \partial_i (r \partial_i \eta_j) d\phi dr \quad (32)$$

$$= \text{surface term} - \iint \eta_j^* \frac{1}{r} \partial_i (r \partial_i \eta_j) r d\phi dr \quad (33)$$

the corresponding functional variation is

$$\delta F / \delta \eta_k^* \sim -\frac{1}{r} \partial_i (r \partial_i \eta_k) = -\frac{1}{r} (\partial_i r) (\partial_i \eta_k) - \partial_i^2 \eta_k = -\frac{1}{r} \partial_r \eta_k - \partial_i^2 \eta_k \quad (34)$$

For the  $\kappa_2$  term in first line of Eq. (30)

$$\delta F / \delta \eta_k^* \sim -\frac{1}{r} \partial_k (r \partial_i \eta_i) = -\frac{1}{r} (\partial_k r) (\partial_i \eta_i) - \partial_k \partial_i \eta_i = -\delta_{kr} \frac{1}{r} \partial_i \eta_i - \partial_k \partial_i \eta_i \quad (35)$$

For the  $\kappa_3$  term in first line of Eq. (30)

$$\delta F / \delta \eta_k^* \sim -\frac{1}{r} \partial_i (r \partial_k \eta_i) = -\frac{1}{r} (\partial_i r) (\partial_k \eta_i) - \partial_i \partial_k \eta_i = -\frac{1}{r} \partial_k \eta_r - \partial_i \partial_k \eta_i \quad (36)$$

For the  $\kappa_1$  term in second line of Eq. (30)

$$F \sim \iint \frac{1}{r} \left( \eta_r^* \frac{\partial \phi}{r} \eta_\phi - \eta_\phi^* \frac{\partial \phi}{r} \eta_r + \eta_r \frac{\partial \phi}{r} \eta_\phi^* - \eta_\phi \frac{\partial \phi}{r} \eta_r^* \right) r d\phi dr \quad (37)$$

the corresponding functional variation is

$$\delta F / \delta \eta_k^* \sim \frac{1}{r} \left( 2\delta_{kr} \frac{\partial \phi}{r} \eta_\phi - 2\delta_{k\phi} \frac{\partial \phi}{r} \eta_r \right) \quad (38)$$

The  $\kappa_2$  term in second line of Eq. (30) is

$$F \sim \iint \frac{1}{r} (\eta_r^* \partial_j \eta_j + \eta_r \partial_j \eta_j^*) r d\phi dr \quad (39)$$

the corresponding functional variation is

$$\delta F / \delta \eta_k^* \sim \frac{1}{r} (\delta_{kr} \partial_j \eta_j - \partial_k \eta_r) \quad (40)$$

The  $\kappa_3$  term in second line of Eq. (30) is

$$F \sim \iint \frac{1}{r} \left( \eta_r^* \frac{\partial \phi}{r} \eta_\phi - \eta_\phi^* \partial_r \eta_\phi + \eta_r \frac{\partial \phi}{r} \eta_\phi^* - \eta_\phi \partial_r \eta_\phi^* \right) r d\phi dr \quad (41)$$

the corresponding functional variation is

$$\delta F / \delta \eta_k^* \sim \frac{1}{r} \left( \delta_{kr} \frac{\partial \phi}{r} \eta_\phi - \cancel{\delta_{k\phi} \partial_r \eta_\phi} - \delta_{k\phi} \frac{\partial \phi}{r} \eta_r + \cancel{\delta_{kr} \partial_r \eta_\phi} \right) \quad (42)$$

Adding all these contributions together, we can get the GL equations in cylindrical basis.

In cartesian basis the GL equations are

$$\alpha\eta_k + 2(\beta_1\eta_i\eta_i^*\eta_k + \beta_2\eta_i^2\eta_k^*) - (\kappa_1\partial_i^2\eta_k + \kappa_2\partial_k\partial_i\eta_i + \kappa_3\partial_j\partial_k\eta_j) = 0 \quad (43)$$

If we transform it to cylindrical basis, we can also get

$$\begin{aligned} \alpha\eta_k + 2(\beta_1\eta_i\eta_i^*\eta_k + \beta_2\eta_i^2\eta_k^*) - (\kappa_1\partial_i^2\eta_k + \kappa_2\partial_k\partial_i\eta_i + \kappa_3\partial_j\partial_k\eta_j) \\ - \frac{1}{r} \left[ (2\kappa_1 + \kappa_3) \frac{\partial_\phi}{r} (\eta_r\delta_{k\phi} - \eta_\phi\delta_{kr}) + \kappa_1\partial_r\eta_k + (\kappa_2 + \kappa_3)\partial_k\eta_r \right] \\ + \frac{1}{r^2} [\kappa_1\eta_k + (\kappa_2 + \kappa_3)\eta_r\delta_{kr}] = 0. \end{aligned} \quad (44)$$

The  $\kappa_1$  term is just a Laplacian term

$$\partial_i^{\text{Car}} \partial_i^{\text{Car}} \eta_k^{\text{Car}} = R_{il} \partial_l (R_{im} \partial_m (R_{kn} \eta_n)) \quad (45)$$

$$= R_{il} (\partial_l R_{im}) \partial_m (R_{kn} \eta_n) + R_{il} R_{im} \partial_l \partial_m (R_{kn} \eta_n) \quad (46)$$

$$= \frac{1}{r} \partial_r (R_{kn} \eta_n) + \partial_l^2 (R_{kn} \eta_n) \quad (47)$$

$$= R_{kn} \frac{1}{r} \partial_r \eta_n + R_{kn} \partial_l^2 \eta_n + 2(\partial_l R_{kn}) (\partial_l \eta_n) + \eta_n \partial_l^2 R_{kn} \quad (48)$$

$$= R_{kn} \left( \frac{1}{r} \partial_r \eta_n + \partial_l^2 \eta_n + \frac{2}{r} R_{nm}^{\pi/2} \frac{\partial_\phi}{r} \eta_m - \frac{1}{r^2} \eta_n \right) \quad (49)$$

$$= R_{kn} \left( \frac{1}{r} \partial_r \eta_n + \partial_l^2 \eta_n + \frac{2}{r} \delta_{n\phi} \frac{\partial_\phi}{r} \eta_r - \frac{2}{r} \delta_{nr} \frac{\partial_\phi}{r} \eta_\phi - \frac{1}{r^2} \eta_n \right) \quad (50)$$

The  $\kappa_2$  term is

$$\partial_k^{\text{Car}} \partial_i^{\text{Car}} \eta_i^{\text{Car}} = R_{kl} \partial_l (R_{im} \partial_m (R_{in} \eta_n)) \quad (51)$$

$$= R_{kl} \partial_l (R_{im} (\partial_m R_{in}) \eta_n + R_{im} R_{in} (\partial_m \eta_n)) \quad (52)$$

$$= R_{kl} \partial_l \left( \frac{1}{r} \eta_r + \partial_m \eta_m \right) \quad (53)$$

$$= R_{kl} \left( \frac{1}{r} \partial_l \eta_r - \delta_{lr} \frac{1}{r^2} \eta_r + \partial_l \partial_m \eta_m \right) \quad (54)$$

The  $\kappa_3$  term is

$$\partial_j^{\text{Car}} \partial_k^{\text{Car}} \eta_j^{\text{Car}} = R_{jl} \partial_l (R_{km} \partial_m (R_{jn} \eta_n)) \quad (55)$$

$$= R_{jl} (\partial_l R_{km}) \partial_m (R_{jn} \eta_n) + R_{jl} R_{km} \partial_l \partial_m (R_{jn} \eta_n) \quad (56)$$

$$= R_{j\phi} \frac{1}{r} R_{ki} R_{im}^{\pi/2} \partial_m (R_{jn} \eta_n) + R_{jl} R_{ki} \partial_l \partial_i (R_{jn} \eta_n) \quad (57)$$

$$= R_{ki} \left[ R_{j\phi} \frac{1}{r} R_{im}^{\pi/2} \partial_m (R_{jn} \eta_n) + R_{jl} \partial_l \partial_i (R_{jn} \eta_n) \right] \quad (58)$$

$$= R_{ki} \left[ R_{j\phi} \frac{1}{r} R_{im}^{\pi/2} (\partial_m R_{jn}) \eta_n + R_{j\phi} \frac{1}{r} R_{im}^{\pi/2} R_{jn} (\partial_m \eta_n) \right] \quad (59)$$

$$\begin{aligned} & + R_{jl} (\partial_l \partial_i R_{jn}) \eta_n + R_{jl} (\partial_l R_{jn}) (\partial_i \eta_n) + R_{jl} (\partial_i R_{jn}) (\partial_l \eta_n) + R_{jl} R_{jn} (\partial_l \partial_i \eta_n) \\ & = R_{ki} \left[ \frac{1}{r^2} R_{i\phi}^{\pi/2} \eta_r + \frac{1}{r} R_{im}^{\pi/2} \partial_m \eta_\phi \right. \\ & \quad \left. + \delta_{i\phi} R_{jl} \eta_n \partial_l \left( \frac{1}{r} R_{jn} \right) + \frac{1}{r} \partial_i \eta_r + \delta_{i\phi} \frac{1}{r} R_{ln}^{\pi/2} \partial_l \eta_n + \partial_l \partial_i \eta_l \right] \\ & = R_{ki} \left[ -\frac{1}{r^2} \delta_{ir} \eta_r + \cancel{\frac{1}{r} \delta_{i\phi} \partial_r \eta_\phi} - \frac{1}{r^2} \delta_{ir} \partial_\phi \eta_\phi \right. \\ & \quad \left. + \delta_{i\phi} R_{jl} \eta_n \left( \partial_l \frac{1}{r} \right) R_{jn} + \delta_{i\phi} R_{jl} \eta_n \frac{1}{r} (\partial_l R_{jn}) + \frac{1}{r} \partial_i \eta_r + \delta_{i\phi} \frac{1}{r^2} \partial_\phi \eta_r - \cancel{\delta_{i\phi} \frac{1}{r} \partial_r \eta_\phi} + \partial_l \partial_i \eta_l \right] \\ & = R_{ki} \left[ -\frac{1}{r^2} \delta_{ir} \eta_r - \frac{1}{r^2} \delta_{ir} \partial_\phi \eta_\phi \right. \\ & \quad \left. - \cancel{\delta_{i\phi} \eta_r \frac{1}{r^2} + \delta_{i\phi} \eta_r \frac{1}{r}} + \frac{1}{r} \partial_i \eta_r + \delta_{i\phi} \frac{1}{r^2} \partial_\phi \eta_r + \partial_l \partial_i \eta_l \right] \\ & = R_{ki} \left[ -\frac{1}{r^2} \delta_{ir} \eta_r - \frac{1}{r^2} \delta_{ir} \partial_\phi \eta_\phi + \frac{1}{r} \partial_i \eta_r + \delta_{i\phi} \frac{1}{r^2} \partial_\phi \eta_r + \partial_l \partial_i \eta_l \right] \end{aligned} \quad (60)$$

## PHYSICAL OBSERVABLES

In Cartesian basis we denote  $\mathbf{r} = (x, y)$ . A Galilean boost with velocity  $\mathbf{u}$  introduce a local gauge transformation  $\eta_i(\mathbf{r}) \xrightarrow{\mathbf{u}} \eta_i(\mathbf{r}) e^{-iM\mathbf{u}\cdot\mathbf{r}/\hbar}$ , where  $M$  is the mass of a pair of Helium atoms. The phase gradient correspond to a velocity field  $\mathbf{v} = \frac{\hbar}{M} \nabla \theta$  which transform as  $\mathbf{v} \xrightarrow{\mathbf{u}} \mathbf{v} - \mathbf{u}$  under the Galilean boost. We also have  $\partial_i \xrightarrow{\mathbf{u}} \partial_i - iM u_i / \hbar$ . The GL free energy density also transforms as  $f \xrightarrow{\mathbf{u}} f - \mathbf{j} \cdot \mathbf{u} + \mathcal{O}(u^2)$ , where  $\mathbf{j}_k = \frac{2M}{\hbar} \text{Im} \{ \kappa_1 \eta_j^* \partial_k \eta_j + \kappa_2 \eta_k^* \partial_j \eta_j + \kappa_3 \eta_j^* \partial_j \eta_k \}$  is the superfluid mass current density or the momentum density. We transform it to cylindrical

basis and get

$$\mathbf{j}_k = \frac{2M}{\hbar} \left[ \text{Im} \{ \kappa_1 \eta_j^* \partial_k \eta_j + \kappa_2 \eta_k^* \partial_j \eta_j + \kappa_3 \eta_j^* \partial_j \eta_k \} \right. \\ \left. + \text{Im} \{ (2\kappa_1 + \kappa_2 + \kappa_3) \delta_{k\phi} \eta_\phi^* \eta_r / r \} \right], \quad (61)$$

In weak-coupling limit, where  $\beta_1 = 2\beta_2$  and  $\kappa_1 = \kappa_2 = \kappa_3$ , the angular momentum density  $l_z = r j_\phi$  becomes

$$l_z = \frac{2M\kappa_1}{\hbar} \left[ \text{Im} \{ 3\eta_\phi^* \partial_\phi \eta_\phi + \eta_r^* \partial_\phi \eta_r + 4\eta_\phi^* \eta_r \} \right. \\ \left. + \text{Im} \{ r \eta_\phi^* \partial_r \eta_r + r \eta_r^* \partial_r \eta_\phi \} \right]. \quad (62)$$

## ROTATING FRAME

We can stabilize these low-flow states by rotating the annulus at certain angular velocities  $\Omega_m$ . Transform into the rotating frame with angular velocity  $\mathbf{\Omega} = \hat{\mathbf{z}}\Omega$ , and the free energy becomes

$$F' = F - \mathbf{L} \cdot \mathbf{\Omega}. \quad (63)$$

The critical angular velocity  $\Omega_m$  which increase the winding number from  $m-1$  to  $m$  should satisfy  $F'(m, \Omega_m) = F'(m-1, \Omega_m)$ , which gives

$$\Omega_m^\pm = \frac{F_\pm(m) - F_\pm(m-1)}{L_\pm(m) - L_\pm(m-1)}. \quad (64)$$

## UNIFORM-FLOW APPROXIMATION

We can have superflow in the annulus  $\mathbf{v}(\mathbf{r}) = \frac{\hbar}{M} \nabla \theta$  where  $M$  is the mass of a pair of Helium atoms.

When we assume a uniform flow field along the azimuthal direction in the annulus,  $\mathbf{v} = v \hat{\phi}$  and  $v(r) = \frac{\hbar}{Mr} \partial_\phi \theta$ , the order parameter in the annulus simply gains an extra phase factor along the azimuthal direction.

$$\boldsymbol{\eta}(r, \phi, v_s) = \boldsymbol{\eta}(r, v_s) e^{iMvr\phi/\hbar} = \boldsymbol{\eta}(r, v_s) e^{i(\partial_\phi \theta)\phi} \quad (65)$$

When the flow field is small, we denote

$$\boldsymbol{\eta}(r, \phi, v_s) = \boldsymbol{\eta}(r, v_s) e^{i(\partial_\phi \theta)\phi} = e^{\pm i\phi} \left( \eta_r^{(m)}(r), \pm i \eta_\phi^{(m)}(r) \right) e^{im\phi} \quad (66)$$

where  $m$  is an integer as the order parameter should be single-valued after  $2\pi$  winding.

## LOW-FLOW APPROXIMATION

For low flow states in an annulus, we further assume the radial profile won't be affected by the small flow field

$$\eta_i^{(m)}(r) = \eta_i^{(0)}(r) \quad (67)$$

For the denominator in Eq. (64), we have

$$L_{\pm}(m) - L_{\pm}(m-1) = \frac{2M\kappa_1}{\hbar} \int d^3r \left( 3\eta_{\phi}^{(0)}\eta_{\phi}^{(0)} + \eta_r^{(0)}\eta_r^{(0)} \right) \quad (68)$$

For the numerator in Eq. (64), we have to pick out terms associated with  $\partial_{\phi}$ , which are

$$\begin{aligned} f_{\text{grad}} \sim & \frac{\kappa_1}{r^2} \left[ 3(\partial_{\phi}\eta_{\phi})(\partial_{\phi}\eta_{\phi})^* + (\partial_{\phi}\eta_r)(\partial_{\phi}\eta_r)^* \right] \\ & + \frac{\kappa_1}{r} \left[ (\partial_{\phi}\eta_{\phi})(\partial_r\eta_r)^* + (\partial_r\eta_r)(\partial_{\phi}\eta_{\phi})^* + (\partial_{\phi}\eta_r)(\partial_r\eta_{\phi})^* + (\partial_r\eta_{\phi})(\partial_{\phi}\eta_r)^* \right] \\ & + \frac{2\kappa_1}{r^2} \text{Re}\{3\eta_r^*\partial_{\phi}\eta_{\phi} - \eta_{\phi}^*\partial_{\phi}\eta_r\} \end{aligned} \quad (69)$$

Substitute in Eq. (66)

$$\begin{aligned} f_{\text{grad}}(m) \sim & \frac{\kappa_1}{r^2} (m \pm 1)^2 \left[ 3 \left( \eta_{\phi}^{(m)} \right)^2 + \left( \eta_r^{(m)} \right)^2 \right] \\ & + \frac{2\kappa_1}{r} (1 \pm m) \left[ \eta_r^{(m)} \partial_r \eta_{\phi}^{(m)} - \eta_{\phi}^{(m)} \partial_r \eta_r^{(m)} \right] \\ & - \frac{8\kappa_1}{r^2} (1 \pm m) \eta_r^{(m)} \eta_{\phi}^{(m)} \end{aligned} \quad (70)$$

For the second line, the corresponding total free energy is zero as long as  $\eta_i^{(m)}(r)$  is an even function with respect to  $r = R + D/2$ ,

$$2\pi \int_R^{R+D} r dr \rightarrow \int_R^{R+D} \eta_i^{(m)} \partial_r \eta_j^{(m)} dr = 0 \quad (71)$$

Then we will have

$$\begin{aligned} F_{\pm}(m) - F_{\pm}(m-1) = & \int d^3r \frac{\kappa_1}{r^2} \left\{ (2m-1) \left[ 3 \left( \eta_{\phi}^{(0)} \right)^2 + \left( \eta_r^{(0)} \right)^2 \right] \right. \\ & \left. \pm 2 \left[ 3 \left( \eta_{\phi}^{(0)} \right)^2 + \left( \eta_r^{(0)} \right)^2 - 4\eta_r^{(0)}\eta_{\phi}^{(0)} \right] \right\} \end{aligned} \quad (72)$$

We denote the volume integral as  $\langle \dots \rangle = \int d^3r (\dots)$ . Then we have

$$\Omega_m^{\pm} = \frac{\hbar}{M} \frac{(m - \frac{1}{2}) \left\langle \frac{1}{r^2} \left[ 3 \left( \eta_{\phi}^{(0)} \right)^2 + \left( \eta_r^{(0)} \right)^2 \right] \right\rangle \pm \left\langle \frac{1}{r^2} \left[ 3 \left( \eta_{\phi}^{(0)} \right)^2 + \left( \eta_r^{(0)} \right)^2 - 4\eta_r^{(0)}\eta_{\phi}^{(0)} \right] \right\rangle}{\left\langle 3 \left( \eta_{\phi}^{(0)} \right)^2 + \left( \eta_r^{(0)} \right)^2 \right\rangle}$$



## LONDON APPROXIMATION

We assume  $\eta_i^{(m)}(r) = \eta_0$ . The angular momentum increase is

$$L_{\pm}(m) - L_{\pm}(m-1) = \frac{8M}{\hbar} \kappa_1 \eta_0^2 \pi \left[ (R+D)^2 - R^2 \right] h \quad (73)$$

The free energy increase is

$$F_{\pm}(m) - F_{\pm}(m-1) = 8\pi \kappa_1 \eta_0^2 \ln \left( \frac{R+D}{R} \right) h (2m-1) \quad (74)$$

where  $h$  is the thickness of the annulus. The critical angular velocity is

$$\Omega_m^{\pm} = \Omega_c \frac{\xi}{D} \frac{\ln(1+D/R)}{2+D/R} (2m-1), \quad (75)$$

where  $\Omega_c = v_c/R$ ,  $v_c = \hbar/M\xi$ ,  $\xi^2 = \kappa_1/|\alpha|$ . Actually, all of the GL parameters drop out

$$\Omega_m^{\pm} = \frac{\hbar}{MRD} \frac{\ln(1+D/R)}{2+D/R} (2m-1), \quad (76)$$

When  $R \gg D$ , we have

$$\Omega_m^{\pm} = \frac{\hbar}{MR^2} \left( m - \frac{1}{2} \right) \quad (77)$$

i.e.

$$\Omega_1 = \frac{\hbar}{MR^2} \frac{1}{2} \quad \text{for } m=0 \rightarrow m=1 \quad (78)$$

$$\Omega_2 = \frac{\hbar}{MR^2} \frac{3}{2} \quad \text{for } m=1 \rightarrow m=2 \quad (79)$$

$$\Omega_3 = \frac{\hbar}{MR^2} \frac{5}{2} \quad \text{for } m=2 \rightarrow m=3 \quad (80)$$

$$\dots \quad (81)$$

## NUMERICAL SOLUTION

From the bulk free energy term we can get the bulk order parameter  $\eta_0 = \frac{1}{2} \sqrt{\frac{|\alpha|}{\beta_1}}$ . We also define the coherence length  $\xi = \sqrt{\kappa_1/|\alpha|}$ .

In weak-coupling limit, the GL equations in cylindrical basis are

$$\begin{aligned} \alpha \eta_k + 2\beta_1 \eta_i \eta_i^* \eta_k + \beta_1 \eta_i^2 \eta_k^* - \kappa_1 (\partial_i^2 \eta_k + \partial_k \partial_i \eta_i + \partial_j \partial_k \eta_j) \\ - \frac{\kappa_1}{r} \left[ 3 \frac{\partial \phi}{r} (\eta_r \delta_{k\phi} - \eta_{\phi} \delta_{kr}) + \partial_r \eta_k + 2 \partial_k \eta_r \right] \\ + \frac{\kappa_1}{r^2} [\eta_k + 2\eta_r \delta_{kr}] = 0. \end{aligned} \quad (82)$$

The dimensionless GL equations in cylindrical basis are

$$\begin{aligned}
& -\eta_k + \frac{1}{2}\eta_i\eta_i^*\eta_k + \frac{1}{4}\eta_i^2\eta_k^* - (\partial_i^2\eta_k + \partial_k\partial_i\eta_i + \partial_j\partial_k\eta_j) \\
& - \frac{1}{r} \left[ 3\frac{\partial_\phi}{r}(\eta_r\delta_{k\phi} - \eta_\phi\delta_{kr}) + \partial_r\eta_k + 2\partial_k\eta_r \right] \\
& + \frac{1}{r^2}[\eta_k + 2\eta_r\delta_{kr}] = 0.
\end{aligned} \tag{83}$$

The  $r$  component of the GL equations is

$$\begin{aligned}
& -\eta_r + \frac{1}{2}\eta_i\eta_i^*\eta_r + \frac{1}{4}\eta_i^2\eta_r^* - (\partial_i^2\eta_r + \partial_r\partial_i\eta_i + \partial_j\partial_r\eta_j) - \frac{1}{r} \left[ -3\frac{\partial_\phi}{r}\eta_\phi + 3\partial_r\eta_r \right] + \frac{3}{r^2}\eta_r = 0 \\
& -\eta_r + \frac{1}{2}\eta_i\eta_i^*\eta_r + \frac{1}{4}\eta_i^2\eta_r^* - (\partial_i^2\eta_r + \partial_r\partial_i\eta_i + \partial_j\partial_r\eta_j) + 3 \left[ \frac{1}{r^2}\partial_\phi\eta_\phi - \frac{1}{r}\partial_r\eta_r + \frac{\eta_r}{r^2} \right] = 0 \\
& -\eta_r + \frac{3}{4}\eta_r^2\eta_r^* + \frac{1}{2}\eta_\phi\eta_\phi^*\eta_r + \frac{1}{4}\eta_\phi^2\eta_r^* - 3\partial_r^2\eta_r - \frac{1}{r^2}\partial_\phi^2\eta_r - \frac{2}{r}\partial_r\partial_\phi\eta_\phi + \frac{4}{r^2}\partial_\phi\eta_\phi - \frac{3}{r}\partial_r\eta_r + \frac{3}{r^2}\eta_r = 0
\end{aligned}$$

The  $\phi$  component of the GL equations is

$$\begin{aligned}
& -\eta_\phi + \frac{1}{2}\eta_i\eta_i^*\eta_\phi + \frac{1}{4}\eta_i^2\eta_\phi^* - \left( \partial_i^2\eta_\phi + \frac{\partial_\phi}{r}\partial_i\eta_i + \partial_j\frac{\partial_\phi}{r}\eta_j \right) - \frac{1}{r} \left[ 5\frac{\partial_\phi}{r}\eta_r + \partial_r\eta_\phi \right] + \frac{1}{r^2}\eta_\phi = 0 \\
& -\eta_\phi + \frac{3}{4}\eta_\phi^2\eta_\phi^* + \frac{1}{2}\eta_r\eta_r^*\eta_\phi + \frac{1}{4}\eta_r^2\eta_\phi^* - 3\frac{1}{r^2}\partial_\phi^2\eta_\phi - \partial_r^2\eta_\phi - \frac{2}{r}\partial_r\partial_\phi\eta_r - \frac{4}{r^2}\partial_\phi\eta_r - \frac{1}{r}\partial_r\eta_\phi + \frac{1}{r^2}\eta_\phi = 0
\end{aligned}$$

For the uniform flow approximation,

$$\boldsymbol{\eta}^{(n)}(r, \phi) = \left( \eta_r^{(n)}(r), \eta_\phi^{(n)}(r) \right) e^{in\phi} \tag{84}$$

which means that  $\partial_\phi \rightarrow in$ . The solutions of  $n = 1$  may corresponds to the  $(p_x + ip_y, m = 0)$  state, or the  $(p_x - ip_y, m = 2)$  state.

## LARGE FLOW

When the flow field is large, we denote the superfluid flow field at the inner radius as  $v_s = \frac{\hbar}{MR}\partial_\phi\theta$ , and

$$\boldsymbol{\eta}(r, \phi, v_s) = \boldsymbol{\eta}^{(n)}(r)e^{in\phi} = \boldsymbol{\eta}(r, v_s)e^{i(\partial_\phi\theta)\phi} = \boldsymbol{\eta}(r, v_s)e^{i\frac{v_s R}{v_c \xi}\phi} \tag{85}$$

where the critical flow field  $v_c = \hbar/M\xi$  and  $\xi^2 = \kappa_1/|\alpha|$ . We also have  $n = \frac{v_s R}{v_c \xi}$ .

## CARTESIAN BASIS

In Cartesian basis, weak-coupling limit dimensionless GL equations are

$$-\eta_k + \frac{1}{2}\eta_i\eta_i^*\eta_k + \frac{1}{4}\eta_i^2\eta_k^* - (\partial_i^2\eta_k + \partial_k\partial_i\eta_i + \partial_j\partial_k\eta_j) = 0 \quad (86)$$

The  $x$  component of the GL equations is

$$-\eta_x + \frac{3}{4}\eta_x^2\eta_x^* + \frac{1}{2}\eta_y\eta_y^*\eta_x + \frac{1}{4}\eta_y^2\eta_x^* - 3\partial_x^2\eta_x - \partial_y^2\eta_x - 2\partial_y\partial_x\eta_y = 0 \quad (87)$$

The  $y$  component of the GL equations is

$$-\eta_y + \frac{3}{4}\eta_y^2\eta_y^* + \frac{1}{2}\eta_x\eta_x^*\eta_y + \frac{1}{4}\eta_x^2\eta_y^* - 3\partial_y^2\eta_y - \partial_x^2\eta_y - 2\partial_x\partial_y\eta_x = 0 \quad (88)$$