

Midterm Exam

1. (25 points) Write T for a true statement and F for a false statement. You DO NOT need to justify your answers. (다음 중 참인 것과 거짓인 것을 구분하십시오. 답만 적으면 됨)

- (a) The sample space associated with a discrete random variable is always countable. (이 산확률변수와 연관된 표본공간은 항상 countable 이다)
- (b) For discrete random variables X, Y and Z , if $p_{X,Y,Z}(x, y, z) = p_X(x)p_Y(y)p_Z(z), \forall x, y, z$, then $p_{X,Y}(x, y) = p_X(x)p_Y(y), \forall x, y$.
- (c) For the CDF $F_X(x)$ of a random variable X , $(F_X(x) - F_X(y))(x - y) \leq 0$. (확률변수 X 의 누적분포함수 $F_X(x)$ 에 대해 $(F_X(x) - F_X(y))(x - y) \leq 0$ 이 성립한다)
- (d) For a continuous random variable X , $\mathbb{P}(X = x) = 0$ for all real numbers x .
- (e) For a continuous random variable X , its PDF $f_X(x)$ is the probability that $X = x$.
- (f) If two events A and B are disjoint, $\mathbb{P}(A \cap B) = \mathbb{P}(A)\mathbb{P}(B)$.
- (g) If $\mathbb{P}(A) \leq \mathbb{P}(B)$, then $A \subseteq B$.
- (h) For two discrete random variables X and Y , $\text{var}(X+Y) = \text{var}(X) + \text{var}(Y)$ if $p_{X,Y}(x, y) = p_X(x)p_Y(y), \forall x, y$.
- (i) For a probabilistic model (Ω, \mathbb{P}) with countable sample space Ω , the probability of an event A is $\mathbb{P}(A) = \frac{|A|}{|\Omega|}$.
- (j) For the conditional PMF of X given $X \in A$ for some set A , we have $p_{X|X \in A}(x) = 0$ if $x \notin A$.

2. (20 points) Consider two random variables X and Y whose joint PMF $p_{X,Y}(x, y)$ is given as

	3	a	1/18	1/18
y	2	b	1/18	1/18
	1	c	1/18	1/18
		1	2	3
		x		

- (a) Determine the values of a, b and c so that X and Y are independent.
- (b) Find the marginal PMFs $p_X(x)$ and $p_Y(y)$.
- (c) Define two events $A = \{X \geq 2\}$ and $B = \{Y < 2\}$. Are A and B independent? Justify your answer.
- (d) Compute $\mathbb{E}[X|A]$ and $\mathbb{E}[Y|B]$.

3. **(20 points)** There are 6 coins and a fair 6-sided die. Let p_i be the probability of head of i th coin. Assume $p_i = \frac{i}{8}, i = 1, \dots, 6$. Consider the following experiment:

- 1) Roll the die (let i be the number that comes up)
- 2) Pick the i th coin and toss it three times independently.

- (a) Describe the sample space.
- (b) Calculate the probability that at least two heads come up. (you will get full score if the formula is correct; 식만 맞으면 됨)
- (c) Are B and A_i in (b) independent?
- (d) Given that three heads came up, what is the conditional probability that the roll is i .

4. **(20 points)** There are 4 coins and a fair 4-sided die. Let p_i be the probability of head of i th coin. Assume $p_i = \frac{i}{8}, i = 1, \dots, 4$. Consider the following experiment:

- 1) Roll the die (let i be the number that came up)
- 2) Pick the i th coin and toss it three times independently.

- (a) Describe the sample space.
 - (b) Calculate the probability that at least two heads come up. (you will get full score if the formula is correct; 식만 맞으면 됨)
 - (c) Are B and A_i in (b) independent?
 - (d) Given that three heads came up, what is the conditional probability that the roll is i .
-

5. **(20 points)** Consider the following PMF of a random variable X :

$$p_X(x) = \begin{cases} \frac{a+xy}{a^2}, & \text{if } x = 1, 2, 3, y = 1, 2, 3 \\ 0, & \text{otherwise} \end{cases}$$

where a is a constant.

- (a) Find the value of a .
- (b) Find the marginal PMF $p_X(x)$
- (c) Find the conditional PMF of X given $Y = 1$
- (d) Are X and Y independent? Justify your answer.
- (e) Calculate the expectation of X , given $Y = 1$.

6. **(10 points)** Let X be an exponential random variable with parameter $\lambda > 0$.

- (a) Calculate the probability $\mathbb{P}(X \leq a)$ where a is a constant.
- (b) Let X_1, \dots, X_n be independent and identically distributed (i.i.d.) exponential random variables with parameter λ . Define $Y = \max\{X_1, \dots, X_n\}$. Calculate the PDF of Y .

7. (10 points) Let X be an exponential random variable with parameter $\lambda > 0$.

- (a) Calculate the probability $\mathbb{P}(X \geq a)$ where a is a positive constant.
- (b) Let X_1, \dots, X_n be independent and identically distributed (i.i.d.) exponential random variables with parameter λ . Define $Y = \min\{X_1, \dots, X_n\}$. Calculate the PDF of Y .

8. (20 points) Consider investing your capital assets in the stock market. On day 1 (happy day), your stock goes up 20% with probability 0.9, and goes down 20% with probability 0.1. On day 2 (sad day), your stock goes up 20% with probability 0.1, and goes down 20% with probability 0.9. This is repeated afterwards. The following are notation and assumptions: (당신의 현금자산을 주식시장에 투자하는 것을 생각하자. 첫번째 날에는 당신의 주식이 0.9의 확률로 20% 오르고, 0.1의 확률로 20% 내린다. 두번째 날에는 당신의 주식이 0.1의 확률로 20% 오르고, 0.9의 확률로 20% 내린다. 그리고, 이것이 계속 반복된다. 다음은 이 문제 관련 기호와 가정)

- Y_0 : your original capital assets (원금)
- X_i : random variable representing the percentage of your stock price gain on the i th day. So for i odd, the PMF of X_i is (i 번째 날에 당신의 주식이 오른 퍼센티지를 나타내는 확률 변수. i 가 홀수라면 확률질량함수는 다음과 같음)

$$p_{X_i}(x) = \begin{cases} 0.9, & x = 0.2 \\ 0.1, & x = -0.2 \\ 0, & \text{otherwise} \end{cases}$$

For i even, the PMF of X_i is (i 가 짝수라면 확률질량함수는 다음과 같음)

$$p_{X_i}(x) = \begin{cases} 0.1, & x = 0.2 \\ 0.9, & x = -0.2 \\ 0, & \text{otherwise} \end{cases}$$

- X_i 's are independent
- Y_n : your total assets (market value) after n days (n 일 지난 후 총 자산)

(a) Calculate the expectation of X_1 and X_2 .

- (b) Find the expression for Y_n . (Y_n 의 식을 구하시오) (Hint. After the first day, it is $Y_1 = Y_0 + Y_0 \cdot X_1$).
- (c) Calculate your expected total assets after $2n$ days.
- (d) Based on the result in (c), calculate how much you will gain or lose after 30 days.

9. (20 points) There is a face recognition system at the entrance of a building. The system identifies each person (entering the building) correctly with probability p , and incorrectly with probability $1 - p$. The correctness of an identification is independent of all others and also the number of people entering. For each correct identification, the face recognition system company gets 1 dollar of incentive, whereas for incorrect identification, they lose 3 dollars as penalty. The number of people entering the building is a random variable N with the following PMF: (빌딩의 입구에 얼굴인식 시스템이 있다. 이 시스템은 빌딩에 들어오는 각 사람을 p 의 확률로 정확하게 인식하고, $1 - p$ 의 확률로 틀리게 인식한다. 각 사람별 인식 결과는 독립이다. 각각의 정확한 인식에 대해 얼굴인식 시스템 회사는 1달러의 인센티브를 받지만, 부정확하면 3달러의 벌금을 부여받는다. 빌딩에 들어오는 사람의 수는 확률변수 N 이고 다음의 확률질량함수를 갖는다)

$$p_N(n) = \begin{cases} 1/5, & n = 10 \\ 2/5, & n = 20 \\ 2/5, & n = 30 \\ 0, & \text{otherwise} \end{cases}$$

Let Y be the net gain based on the above incentive/penalty policy. (Y 는 상술한 인센티브/벌금 정책에 따른 순수 이득이라고 하자)

- (a) Given that the number of people who entered the building is n , what is the conditional expectation of Y ?
- (b) Based on the result in part (a), find the minimum value p_0 such that the expected net gain is positive for $p > p_0$.
- (c) Assume $p = p_0$ where p_0 is the value found in part (b). Calculate the probability that the net gain is no greater than zero, that is, $\mathbb{P}(Y \leq 0)$.