

# Measurement of electron-transfer cross sections for H(3l) in intermediate-energy H<sup>+</sup>-He collisions

R. A. Cline, W. B. Westerveld,\* and J. S. Risley

*Atomic Collisions Laboratory, Department of Physics, North Carolina State University, Raleigh, North Carolina 27695-8202*

(Received 24 September 1990)

Cross sections have been measured for the production of H(3l) atoms via electron transfer in proton-helium collisions from 25 to 100 keV. In the experiment, the intensity of Balmer- $\alpha$  radiation is measured as a function of proton energy. These measurements are combined with previously measured density matrices for the same collision to produce relative cross sections. Our relative electron-transfer results, accurate to 3–7%, are normalized to the absolute  $\sigma_{3s}$  cross-section measurements of Brower and Pipkin [Phys. Rev. A **39**, 3323 (1989)] at 60 keV. Comparison of our  $\sigma_{3l}$  cross-section measurements with other experimental measurements shows general agreement to within 20%, with larger differences for older results. The recent calculation of Shingal and Lin [J. Phys. B (to be published)] agrees to within 13% of our measurements of  $\sigma_{3s}$  above 30 keV, but predicts a low-energy oscillatory behavior of  $\sigma_{3p}$  and  $\sigma_{3d}$  not present in experimental results.

In ion-atom collisions, electron transfer dominates the intermediate-energy regime, where the ion velocity nearly matches the orbital-electron velocity of the target atom. For this reason, electron-transfer cross sections have been measured in simple collisions for many years.<sup>1–6</sup> Previously, we determined density matrices describing H( $n=3$ ) atoms formed in proton-helium collisions.<sup>7,8</sup> Our results were normalized to  $\sigma_{3s}$  at each energy. We have now measured  $\sigma_{3s}$  as a function of energy and can normalize our previous results to bring them onto an absolute scale. We compare our results to four experimental measurements and three theoretical calculations.

The apparatus was described previously.<sup>7</sup> A proton beam traverses a He gas cell in which there is an applied electric field. Balmer- $\alpha$  radiation is observed perpendicular to the beam from a section of the gas cell. The proton current varies with proton energy from 0.5  $\mu$ A at 25 keV to 3  $\mu$ A at 100 keV. The He gas pressure is 1 mTorr. A transverse electric field of 390 V/cm is directed perpendicular to both the proton beam and the direction of observation. At this field strength significant Stark mixing of the  $n=3$  manifold produces a shorter effective lifetime for H(3s) atoms, leading to a larger counting rate. Additional measurements using zero electric field, which have

a larger uncertainty, agree with the 390-V/cm results.

The intensity of light polarized parallel to the beam  $I_{\parallel}$  is measured as a function of proton energy. Each density-matrix element contributes to  $I_{\parallel}$ :

$$I_{\parallel} = \sigma_{3s} \sum_{\substack{l,m \\ l',m'}} \rho_{3lm;3l'm'} \left[ f_{0;lm;l'm'}^{(t)} + f_{1;lm;l'm'}^{(t)} + \frac{\sigma_{4s}}{\sigma_{3s}} (g_{0;lm;l'm'}^{(t)} + g_{1;lm;l'm'}^{(t)}) \right]. \quad (1)$$

The density-matrix elements  $\rho_{3lm;3l'm'}$ , relative to  $\sigma_{3s}$ , were measured previously.<sup>8</sup> The fitting functions  $f_{ijk}^{(t)}$  and  $g_{ijk}^{(t)}$  for the  $i$ th Stokes parameter for a transverse electric field for individual H( $n=3$ ) and H( $n=4$ ) density-matrix elements were numerically calculated.<sup>7</sup> The  $\sigma_{4s}/\sigma_{3s}$  ratio is determined from theoretical calculations.<sup>9</sup> By measuring  $I_{\parallel}$  and calculating the sum in Eq. (1) we determine  $\sigma_{3s}$  as a function of proton energy. The relative  $\sigma_{3s}$  are normalized at 60 keV to the result of Brower and Pipkin,<sup>6</sup> since the absolute detection efficiency of our optical

TABLE I. Electron-transfer cross sections  $\sigma_{3l}$  for proton-helium collisions.

Proton energy (keV)	$\sigma_{3s}$ ( $10^{-18}$ cm <sup>2</sup> )	$\sigma_{3p}$ ( $10^{-18}$ cm <sup>2</sup> )	$\sigma_{3d}$ ( $10^{-18}$ cm <sup>2</sup> )
25	0.686±0.022	1.209±0.042	0.2250±0.0081
30	1.215±0.039	1.194±0.042	0.1782±0.0070
35	1.652±0.049	1.118±0.037	0.1525±0.0058
40	1.951±0.054	0.944±0.029	0.1327±0.0049
50	2.141±0.065	0.659±0.024	0.1053±0.0047
60	2.000 <sup>a</sup> ±0.066	0.453±0.020	0.0838±0.0044
80	1.417±0.057	0.247±0.015	0.0507±0.0037
100	0.910±0.036	0.1453±0.0089	0.0300±0.0021

<sup>a</sup>Normalization value from Ref. 6.

system is unknown. We believe this absolute normalization is correct because of the general agreement at higher energies for  $\sigma_{3s}$  among our results, three other experimental measurements,<sup>5,5,6</sup> and one theoretical calculation,<sup>10,11</sup> in particular, the close agreement between Lenormand<sup>5</sup> and Brower and Pipkin<sup>6</sup> at 60 keV. Our relative cross-section measurements are put on an absolute scale using the normalized values of  $\sigma_{3s}$ .

Table I lists the absolute electron-transfer cross sections  $\sigma_{3l}$ . The relative uncertainties are (i) 0.3–0.7% from  $I_{\parallel}$ , (ii) 2–3.5% from previously determined density-matrix elements,<sup>8</sup> and (iii) 2% from the current measurement caused by an uncertainty in the amount of beam neutralization due to electron transfer. The results are repeatable within the total relative uncertainty of 3–7%. The uncertainty in the absolute normalization is 20%.<sup>6</sup> The results shown in Table I include corrections for cascade from  $H(n=4)$  which influence the 25-keV results by 5% and the 100-keV results by 1%.

Figures 1–3 compare our results to four experimental measurements<sup>1,3,5,6</sup> and three theoretical calculations.<sup>10–13</sup> Three experimental groups have used the exponential decay method: Hughes *et al.*,<sup>1</sup> Ford and Thomas,<sup>3</sup> and Lenormand.<sup>5</sup> The intensity of Balmer- $\alpha$  radiation is measured as a function of position following a gas cell. The signal is deconvoluted into three exponential curves characteristic of decay from 3s, 3p, and 3d states. No correction is made for cascade. Another experimental method is the microwave resonance optical detection technique of Brower and Pipkin.<sup>6</sup> A detector is placed downstream from a microwave cavity and a gas cell. Mi-




FIG. 1.  $\sigma_{3s}$  as a function of proton energy.  $\bullet$ , present results;  $\blacksquare$ , Brower and Pipkin (Ref. 6);  $\blacktriangle$ , Lenormand (Ref. 5);  $\times$ , Ford and Thomas, (Ref. 3);  $+$ , Hughes *et al.* (Ref. 1);  $\cdots$ , Jain, Lin, and Fritsch (Ref. 12);  $---$ , Shingal and Lin (Ref. 10);  $---$  (Ref. 13).




FIG. 2.  $\sigma_{3p}$  as a function of proton energy. Same symbols as Fig. 1.

crowave radiation drives transitions within the  $n=3$  manifold and Balmer- $\alpha$  intensity is measured as the microwaves are turned on and off. Corrections are made for cascade from  $H(n>3)$  atoms up to  $n=8$ . Systematic effects in these experiments have been discussed previously.<sup>2,14</sup> The uncertainty in the absolute value for all of the experiments is reported to be about 20%. Three calculations have been performed for this collision. Jain, Lin, and Fritsch<sup>12</sup> use the augmented atomic orbital (AO+) method, a modified two-center atomic-orbital expansion




FIG. 3.  $\sigma_{3d}$  as a function of proton energy. Same symbols as Fig. 1.

supplemented by pseudostates. Shingal and Lin<sup>10,11</sup> use the same method with a much larger set of basis states. The calculations by Dubé<sup>13</sup> use the continuum distorted-wave approximation with postcollision interaction<sup>15</sup> (CDW-PCI).

For  $\sigma_{3s}$  above a proton energy of 45 keV all of the experimental results agree to within 18%, except for Hughes *et al.*<sup>1</sup> For  $\sigma_{3p}$ , Lenormand,<sup>5</sup> Brower and Pipkin,<sup>6</sup> and the present results agree to within 23% at all energies. For  $\sigma_{3d}$  below 80 keV, our results agree to within 41% of both Lenormand<sup>5</sup> and Brower and Pipkin,<sup>6</sup> although their results differ by as much as 57%.

The CDW-PCI theory is a high-energy approximation. As the collision energy decreases, the predicted cross sections increase monotonically, overestimating  $\sigma_{3l}$  by a factor of 10 to 15 at 25 keV. At 100 keV, the calculations asymptotically approach the experimental measurements. For all  $\sigma_{3l}$ , CDW-PCI predicts smooth variations as a function of energy.

The calculations of Jain, Lin, and Fritsch<sup>12</sup> overestimate  $\sigma_{3s}$  by a factor of 2. The recent calculations by Shingal and Lin<sup>10,11</sup> agree to within 13% of our measurements of  $\sigma_{3s}$  above 30 keV. Both coupled-state calculations

predict oscillatory behavior for  $\sigma_{3p}$  and  $\sigma_{3d}$  not found in experimental measurements. Because the cross sections  $\sigma_{3l}$  are several orders of magnitude smaller than the  $H(n=1)$  cross section, coupled-state calculations must include large basis sets to be accurate.

In our previous research, considerable effort was made to eliminate systematic effects in the experiment, leading to self-consistent results.<sup>7</sup> Because of this we believe our relative results for  $\sigma_{3l}$  are the most accurate to date, and should therefore be used in comparison with future experimental and theoretical research.

The authors thank Ashok Jain, Louis Dubé, and Rajiv Shingal for providing us with their results prior to publication. The fitting functions used in Eq. (1) were calculated using the Cornell National Supercomputer Facility, a resource of the Cornell Theory Center, which is funded in part by the National Science Foundation, New York State, the IBM Corporation, and members of the Center's Corporate Research Institute. This work was supported in part by the Atomic, Molecular, and Plasma Physics Program of the National Science Foundation, under Grant No. PHY-88-09083.

\*Permanent address: Department of Physics, University of Utrecht, Utrecht, The Netherlands.

<sup>1</sup>R. H. Hughes, C. A. Stigers, B. M. Doughty, and E. D. Stokes, *Phys. Rev. A* **1**, 1424 (1970).

<sup>2</sup>E. W. Thomas, *Excitation in Heavy Particle Collisions* (Wiley, New York, 1972).

<sup>3</sup>J. C. Ford and E. W. Thomas, *Phys. Rev. A* **5**, 1694 (1972).

<sup>4</sup>J. E. Bayfield, in *Atomic Physics, Vol. 4*, edited by G. zu Putlitz, E. W. Weber, and A. Winnacker (Plenum, New York, 1975), p. 397.

<sup>5</sup>J. Lenormand, *J. Phys. (Paris)* **37**, 699 (1976).

<sup>6</sup>M. C. Brower and F. M. Pipkin, *Phys. Rev. A* **39**, 3323 (1989).

<sup>7</sup>J. R. Ashburn, R. A. Cline, C. D. Stone, P. J. M. van der

Burgt, W. B. Westerveld, and J. S. Risley, *Phys. Rev. A* **40**, 4885 (1989).

<sup>8</sup>J. R. Ashburn, R. A. Cline, P. J. M. van der Burgt, W. B. Westerveld, and J. S. Risley, *Phys. Rev. A* **41**, 2407 (1990).

<sup>9</sup>A. Jain (private communication).

<sup>10</sup>R. Shingal and C. D. Lin, *J. Phys. B* (to be published).

<sup>11</sup>R. Shingal (private communication).

<sup>12</sup>A. Jain, C. D. Lin, and W. Fritsch, *Phys. Rev. A* **36**, 2041 (1987).

<sup>13</sup>L. Dubé (private communication).

<sup>14</sup>B. Van Zyl, M. W. Gealy, and H. Neumann, *Phys. Rev. A* **33**, 2333 (1986).

<sup>15</sup>J. Burgdörfer and L. J. Dubé, *Phys. Rev. Lett.* **52**, 2225 (1984).