

Table 2. Parameter provenance

Panel A. Parameter identity

Symbol	Definition	Value or range	Unit	Status
g_SQ	isotropic doxorubicin semiquinone g value	2.0035	dimensionless	measured
deltaHpp_SQ	peak-to-peak aqueous one-line width	3.25	G	measured
deltaHpp_SQ_replicatio n	independent aqueous one-line width	3.2	G	measured
A_SQ	resolved doxorubicin/adriamycin semiquinone hyperfine values	numeric extraction unavailable	not available	measured source identified
lambda_0	gas-phase O2 spin-spin fine-structure constant	1.9848	cm ⁻¹	fitted
gamma_0	gas-phase O2 spin-rotation constant	-0.0084293	cm ⁻¹	fitted
D_O2_gas	axial D from D=2 lambda convention conversion	3.9695	cm ⁻¹	calculated
D_O2_solid_air	axial matrix O2 zero-field splitting	3.572(3)	cm ⁻¹	measured fitted
D_O2_N2_matrix	O2 zero-field splitting in solid nitrogen	3.497	cm ⁻¹	measured fitted
T1_O2_solution	bulk dissolved oxygen electron spin-lattice relaxation	approximately 7.5	ps	indirectly fitted
K_O2_T4L	O2 association constant in T4 lysozyme L99A cavities	48	M ⁻¹	measured fitted
tau_rot_O2_T4L	O2 rotational correlation components	0.164;1.41	ps	calculated fitted
g_D_E_O2_encounter	O2 g and ZFS tensors in transient doxorubicin-semiquinone/O2 encounter	unavailable	not available	unavailable
T1_T2_Tm_SQ	doxorubicin or close anthracycline semiquinone electron-spin relaxation/dephasing	unavailable	s	unavailable
k_O2_total_pH6	total bulk semiquinone oxidation by O2	3.500e+08	M ⁻¹ s ⁻¹	measured

Each parameter retains its stated conditions, evidence status, and permitted use. Presentation values are rounded. All 40 source rows and all 23 source columns remain in table02_parameter_provenance.csv; 7 provenance/support columns are source-only. 'Not available' is distinct from numeric zero.

Table 2. Parameter provenance

Panel A. Parameter identity (continued)

Symbol	Definition	Value or range	Unit	Status
k_O2_total_pH11_5	total bulk semiquinone oxidation by O2	1.700e+08	M ⁻¹ s ⁻¹	measured
k_O2_total_1981	earlier total bulk semiquinone oxidation by O2	4.400e+07	M ⁻¹ s ⁻¹	measured
t_half_SQ_anaerobic	semiquinone chemical half-life	50	us	measured
diffusion_distance_SQ	author-calculated bulk travel bounds	<0.6 anaerobic;<0.1 air-saturated	um	calculated
encounter_geometry_J_D_kassoc_ksep	separation; orientation; exchange; dipolar; association/separation; cage duration	unavailable	not available	unavailable
DQ_coherence	direct spin-correlated coherent D/Q encounter evidence	unavailable	not available	unavailable
k_D	doublet encounter electron-transfer rate	unavailable	s ⁻¹	unavailable
k_Q	quartet encounter electron-transfer rate	unavailable	s ⁻¹	unavailable
k_Q_over_k_D	quartet/doublet rate ratio	unavailable	dimensionless	unavailable
pKa_HO2	hydroperoxyl acid dissociation constant	4.88	dimensionless	measured
k_HH	elementary HO2 plus HO2 dismutation rate	7.600e+05	M ⁻¹ s ⁻¹	measured
k_HA	elementary HO2 plus O2 radical-anion dismutation rate	8.500e+07	M ⁻¹ s ⁻¹	measured
k_AA	direct superoxide plus superoxide disproportionation upper bound	<100	M ⁻¹ s ⁻¹	measured upper bound
k_SOD_oxidation	bovine CuZnSOD oxidized-enzyme half-reaction	1.200e+09	M ⁻¹ s ⁻¹	measured
k_SOD_reduction	bovine CuZnSOD reduced-enzyme half-reaction	2.200e+09	M ⁻¹ s ⁻¹	measured

Each parameter retains its stated conditions, evidence status, and permitted use. Presentation values are rounded. All 40 source rows and all 23 source columns remain in table02_parameter_provenance.csv; 7 provenance/support columns are source-only. 'Not available' is distinct from numeric zero.

Table 2. Parameter provenance

Panel A. Parameter identity (continued)

Symbol	Definition	Value or range	Unit	Status
k_SOD_overall	overall bovine CuZnSOD-superoxide rate	2.370e+09	M ⁻¹ s ⁻¹	measured
v_O2minus_heart_SMP	superoxide formation rate	control 1.6+/-0.2;doxorubicin 69.6+/-2.7	nmol min ⁻¹ mg ⁻¹	measured mean SE
v_H2O2_heart_SMP	hydrogen peroxide formation rate	control undetectable;d oxorubicin 2.2+/-0.3	nmol min ⁻¹ mg ⁻¹	measured mean SE
v_O2minus_Ehrlich_micr osomes	SOD-inhibitable superoxide proxy rate; control vs drug	0.51+/-0.26 n=3;14.71+/-1.43 n=6	nmol min ⁻¹ mg ⁻¹	measured mean SE
v_O2minus_Ehrlich_mito chondria	SOD-inhibitable superoxide proxy rate; control vs drug	1.12+/-0.15 n=7;7.29+/-0.61 n=10	nmol min ⁻¹ mg ⁻¹	measured mean SE
v_O2minus_Ehrlich_nucl ei	SOD-inhibitable superoxide proxy rate; control vs drug	0.36+/-0.05 n=7;3.29+/-0.33 n=12	nmol min ⁻¹ mg ⁻¹	measured mean SE
v_H2O2_Ehrlich	H2O2 formation in microsomes; mitochondria; intact cells	3.42+/-0.48 n=4;2.6+/-0.6 n=3;0.64+/-0.04 n=3	nmol min ⁻¹ mg ⁻¹ ; nmol min ⁻¹ per 1e7 cells	measured mean SE
H2O2_PC3	apparent intracellular steady-state hydrogen peroxide	basal 13+/-4;1 uM doxorubicin 51+/-13	pM	measured derived
MitoSOX_H9c2	relative mitochondrial oxidant fluorescence	1.40+/-0.04;1.80+/- 0.08;2.80+/-0.08	fold control	measured mean SD
negative_5_imino	quinone-disabled anthracycline negative control	doxorubicin 14.71+/- 1.43;5- iminodaunorubicin 0.45+/-0.15	nmol min ⁻¹ mg ⁻¹	measured mean SE

Each parameter retains its stated conditions, evidence status, and permitted use. Presentation values are rounded. All 40 source rows and all 23 source columns remain in table02_parameter_provenance.csv; 7 provenance/support columns are source-only. 'Not available' is distinct from numeric zero.

Table 2. Parameter provenance

Panel B. Chemical conditions

Symbol	Species	Charge state	Environment	Temperature	pH
g_SQ	doxorubicin semiquinone	protonation/formal charge not assigned	N2-purged 100 mM phosphate; 100 uM doxorubicin; 400 uM xanthine; 0.2 U xanthine oxidase in 2 mL	not reported	7.5
deltaHpp_SQ	doxorubicin semiquinone	protonation/formal charge not assigned	N2-purged 100 mM phosphate; 100 uM doxorubicin; 400 uM xanthine; 0.2 U xanthine oxidase in 2 mL	not reported	7.5
deltaHpp_SQ_replication	doxorubicin semiquinone	protonation/formal charge not assigned	hypoxic 50 mM Tris; 1 mM doxorubicin; 400 uM hypoxanthine; 0.1 U/mL xanthine oxidase	310 K	7.4
A_SQ	adriamycin semiquinone	not reported in accessible abstract	not reported in accessible abstract	not reported in accessible abstract	not reported in accessible abstract
lambda_0	16O2 X3Sigma_g- v=0	neutral	gas phase	not extracted	pH not applicable
gamma_0	16O2 X3Sigma_g- v=0	neutral	gas phase	not extracted	pH not applicable
D_O2_gas	16O2 X3Sigma_g- v=0	neutral	gas phase	not extracted	pH not applicable
D_O2_solid_air	ground-state O2	neutral	solid air condensed in waveguide	5 K	pH not applicable
D_O2_N2_matrix	ground-state O2	neutral	O2 diluted in solid N2	15 K	pH not applicable
T1_O2_solution	ground-state O2	neutral	water; 1:1 water/glycerol; cyclohexane; methanol; benzene; acetone; acetonitrile; DMSO	room temperature	not reported/not applicable

Each parameter retains its stated conditions, evidence status, and permitted use. Presentation values are rounded. All 40 source rows and all 23 source columns remain in table02_parameter_provenance.csv; 7 provenance/support columns are source-only. 'Not available' is distinct from numeric zero.

Table 2. Parameter provenance

Panel B. Chemical conditions (continued)

Symbol	Species	Charge state	Environment	Temperature	pH
K_O2_T4L	ground-state O2	neutral	T4 lysozyme L99A; 50 mM phosphate; 25 mM NaCl	298 K	5.5
tau_rot_O2_T4L	ground-state O2	neutral	T4 lysozyme L99A cavities; 50 mM phosphate; 25 mM NaCl	298 K	5.5
g_D_E_O2_encounter	ground-state O2	neutral	doxorubicin-semiquinone solution encounter	unavailable	unavailable
T1_T2_Tm_SQ	doxorubicin semiquinone	unresolved	chemically relevant solution	unavailable	unavailable
k_O2_total_pH6	adriamycin semiquinone plus ground-state O2	biological-pH semiquinone described as internally protonated/zwitterionic; O2 neutral	aqueous phosphate/formate system; O2 or O2/N2 purge at least 30 min	room temperature	6
k_O2_total_pH11_5	adriamycin semiquinone plus ground-state O2	deprotonated semiquinone form; O2 neutral	aqueous borate/OH- system; O2 or O2/N2 purge at least 30 min	room temperature	11.5
k_O2_total_1981	adriamycin semiquinone plus O2	not reported in accessible source	aqueous buffer	not reported in accessible source	not reported in accessible source
t_half_SQ_anaerobic	adriamycin semiquinone	not reported in accessible source	anaerobic aqueous medium	not reported in accessible source	not reported in accessible source
diffusion_distance_SQ	adriamycin semiquinone	not reported in accessible source	aqueous buffer	not reported in accessible source	not reported in accessible source
encounter_geometry_J_D_kassoc_ksep	doxorubicin-semiquinone/O2 encounter	unresolved	chemically relevant solution	unavailable	unavailable
DQ_coherence	doxorubicin-semiquinone/O2 encounter	unresolved	chemically relevant solution	unavailable	unavailable

Each parameter retains its stated conditions, evidence status, and permitted use. Presentation values are rounded. All 40 source rows and all 23 source columns remain in table02_parameter_provenance.csv; 7 provenance/support columns are source-only. 'Not available' is distinct from numeric zero.

Table 2. Parameter provenance

Panel B. Chemical conditions (continued)

Symbol	Species	Charge state	Environment	Temperature	pH
k _D	doublet doxorubicin-semiquinone/O ₂ encounter	unresolved	chemically relevant solution	unavailable	unavailable
k _Q	quartet doxorubicin-semiquinone/O ₂ encounter	unresolved	chemically relevant solution	unavailable	unavailable
k _Q /k _D	doxorubicin-semiquinone/O ₂ encounter	unresolved	chemically relevant solution	unavailable	unavailable
pK _a _{HO₂}	HO ₂ radical to O ₂ radical anion	neutral to -1	oxygen-saturated formic acid/sodium formate aqueous buffer	not reported	0-13 investigated
k _{HH}	HO ₂ radical plus HO ₂ radical	neutral plus neutral	oxygen-saturated formic acid/sodium formate aqueous buffer	not reported	0-13 mechanism study
k _{HA}	HO ₂ radical plus superoxide	neutral plus -1	oxygen-saturated formic acid/sodium formate aqueous buffer	not reported	0-13 mechanism study
k _{AA}	superoxide plus superoxide	-1 plus -1	oxygen-saturated formic acid/sodium formate aqueous buffer	not reported	0-13 mechanism study
k _{SOD_oxidation}	bovine CuZnSOD plus superoxide	enzyme state E0 plus -1	aqueous O ₂ ; EDTA; formate	not reported	not reported in accessible record
k _{SOD_reduction}	bovine CuZnSOD plus superoxide	enzyme state E- plus -1	aqueous O ₂ ; EDTA; formate	not reported	not reported in accessible record
k _{SOD_overall}	native bovine CuZnSOD plus superoxide	enzyme plus -1	aqueous	298 K	not reported in accessible abstract

Each parameter retains its stated conditions, evidence status, and permitted use. Presentation values are rounded. All 40 source rows and all 23 source columns remain in table02_parameter_provenance.csv; 7 provenance/support columns are source-only. 'Not available' is distinct from numeric zero.

Table 2. Parameter provenance

Panel B. Chemical conditions (continued)

Symbol	Species	Charge state	Environment	Temperature	pH
v_O2minus_heart_SMP	bovine-heart submitochondrial particles plus doxorubicin	doxorubicin HCl microspecies not reported	isolated submitochondrial particles	not reported in accessible abstract	not reported in accessible abstract
v_H2O2_heart_SMP	bovine-heart submitochondrial particles plus doxorubicin	doxorubicin HCl microspecies not reported	isolated submitochondrial particles	not reported in accessible abstract	not reported in accessible abstract
v_O2minus_Ehrlich_microsomes	Ehrlich-Lette tumor microsomes plus doxorubicin	doxorubicin HCl microspecies not reported	200 ug protein in 1 mL; 150 mM potassium phosphate; 100 uM EDTA; 1 mM NADPH	310 K	7.4
v_O2minus_Ehrlich_mitochondria	Ehrlich tumor mitochondria plus doxorubicin	doxorubicin HCl microspecies not reported	100 ug protein/mL; sucrose/20 mM HEPES/100 uM EDTA; 100 uM NADH; 4 uM rotenone; 5 min preincubation	310 K	8.2
v_O2minus_Ehrlich_nuclei	Ehrlich tumor nuclear fraction plus doxorubicin	doxorubicin HCl microspecies not reported	200 ug nuclear protein; sucrose/20 mM HEPES/100 uM EDTA; 1 mM NADPH	310 K	7.4
v_H2O2_Ehrlich	Ehrlich tumor fractions or intact cells plus doxorubicin	doxorubicin HCl microspecies not reported	fraction buffers as above; intact 1.5e7 cells/3 mL Chelex-treated PBS	310 K	7.4 microsomes; 8.2 mitochondria; cell PBS pH not reported
H2O2_PC3	PC3 human prostate cancer cells plus doxorubicin	doxorubicin microspecies not reported	cell culture	310 K	not reported for intracellular measurement
MitoSOX_H9c2	rat H9c2 cardiomyoblasts plus doxorubicin	doxorubicin microspecies not reported	HBSS with Ca/Mg and 1% BSA; 5 uM MitoSOX	310 K	not reported

Each parameter retains its stated conditions, evidence status, and permitted use. Presentation values are rounded. All 40 source rows and all 23 source columns remain in table02_parameter_provenance.csv; 7 provenance/support columns are source-only. 'Not available' is distinct from numeric zero.

Table 2. Parameter provenance

Panel B. Chemical conditions (continued)

Symbol	Species	Charge state	Environment	Temperature	pH
negative_5_imino	Ehrlich microsomes plus anthracycline	dissolved microspecies not reported	microsomal system above	310 K	7.4

Each parameter retains its stated conditions, evidence status, and permitted use. Presentation values are rounded. All 40 source rows and all 23 source columns remain in table02_parameter_provenance.csv; 7 provenance/support columns are source-only. 'Not available' is distinct from numeric zero.

Table 2. Parameter provenance

Panel C. Evidence and permitted use

Symbol	Method	Source	Uncertainty	Limitations	Permitted use	Code location
g_SQ	CW EPR; 1 G modulation; 40 G scan; 1 mW	https://doi.org/10.1016/0003-9861(91)90023-C	not reported	isotropic one-line signal; no g tensor	sensitivity and initialization	active_model.encounter.g_radical
deltaHpp_SQ	CW EPR	https://doi.org/10.1016/0003-9861(91)90023-C	not reported	unresolved hyperfine and homogeneous/inhomogeneous broadening; cannot uniquely convert to T2	validation only	reference_evidence.hyperfine.aqueous_linewidth_pH7_5
deltaHpp_SQ_replication	CW EPR	https://doi.org/10.1124/mol.115.098798	not reported	unresolved line; not a hyperfine tensor or T2 measurement	validation only	reference_evidence.hyperfine.aqueous_linewidth_replication
A_SQ	EPR; ENDOR; TRIPLE resonance; partial deuteration	https://doi.org/10.1002/mrc.1260260812	not reported in accessible abstract	resolved assignments were investigated; numeric table/tensors/signs/isotope mapping and conditions were not accessible; do not say not investigated	unavailable for parameterization	reference_evidence.hyperfine.exact_study
lambda_0	microwave/optical spectral fit	https://hitran.org/media/refs/HITRAN-1973.pdf	not available in accessible table	gas rovibronic Hamiltonian and convention; never transfer into solution encounter	context only	reference_evidence.oxygen.gas.lambda_0
gamma_0	microwave/optical spectral fit	https://hitran.org/media/refs/HITRAN-1973.pdf	not available in accessible table	gas rovibronic Hamiltonian; never transfer into solution encounter	context only	reference_evidence.oxygen.gas.gamma_0
D_O2_gas	derived convention conversion	https://hitran.org/media/refs/HITRAN-1973.pdf	not propagated	not a new measurement; sign and normalization depend on Hamiltonian convention	context only	reference_evidence.oxygen.gas.derived_D
D_O2_solid_air	94-550 GHz multifrequency EPR	https://doi.org/10.1006/jmre.2000.2175	0.003 cm ⁻¹	frozen solid only; prohibited as solution surrogate	context only	reference_evidence.oxygen.matrix_solid_air

Each parameter retains its stated conditions, evidence status, and permitted use. Presentation values are rounded. All 40 source rows and all 23 source columns remain in table02_parameter_provenance.csv; 7 provenance/support columns are source-only. 'Not available' is distinct from numeric zero.

Table 2. Parameter provenance

Panel C. Evidence and permitted use (continued)

Symbol	Method	Source	Uncertainty	Limitations	Permitted use	Code location
D_O2_N2_matrix	high-field high-frequency EPR	https://doi.org/10.1016/S0921-4526(00)00615-3	not reported	temperature dependent; signal disappears above 25 K; g=2/no rhombicity fit	context only	reference_evidence.oxygen.matrix_nitrogen
T1_O2_solution	proton magnetic relaxation dispersion	https://doi.org/10.1006/jmre.2000.2219	not reported in abstract	bulk fluid scale; no direct time-domain O2 T1; not encounter-specific and not a static g/D/E tensor	sensitivity context	reference_evidence.oxygen.bulk_solution
K_O2_T4L	gas-pressure NMR plus MD	https://doi.org/10.1038/srep20534	7 M ⁻¹	protein-cavity association; not doxorubicin/O2 and not a spin-Hamiltonian value	context only	reference_evidence.oxygen.protein_cavity.association
tau_rot_O2_T4L	molecular dynamics fit	https://doi.org/10.1038/srep20534	0.006;0.02 ps	rapid orientation averaging; not doxorubicin encounter dynamics	context only	reference_evidence.oxygen.protein_cavity.rotational_correlation
g_D_E_O2_encounter	targeted primary-literature search	NA	NA	not found; gas/matrix/liquid/protein values are not substitutes	sensitivity only	reference_evidence.oxygen.doxorubicin_semiquinone_encounter
T1_T2_Tm_SQ	pulse-EPR and linewidth search	NA	NA	only unresolved CW linewidths found; chemical half-life is not spin relaxation	sensitivity only	reference_evidence.hyperfine.spin_relaxation
k_O2_total_pH6	electron pulse radiolysis transient absorption	https://doi.org/10.1038/bjc.1985.74	0.4e8 M ⁻¹ s ⁻¹	total channel only; homogeneous solution; no encounter geometry or spin resolution	quantitative bulk validation	parameter_records.doxorubicin_semiquinone_plus_oxygen_pH6
k_O2_total_pH11_5	electron pulse radiolysis transient absorption	https://doi.org/10.1038/bjc.1985.74	0.2e8 M ⁻¹ s ⁻¹	total channel only; alkaline; no encounter geometry or spin resolution	quantitative bulk validation	parameter_records.doxorubicin_semiquinone_plus_oxygen_pH11_5

Each parameter retains its stated conditions, evidence status, and permitted use. Presentation values are rounded. All 40 source rows and all 23 source columns remain in table02_parameter_provenance.csv; 7 provenance/support columns are source-only. 'Not available' is distinct from numeric zero.

Table 2. Parameter provenance

Panel C. Evidence and permitted use (continued)

Symbol	Method	Source	Uncertainty	Limitations	Permitted use	Code location
k_O2_total_1981	pulse radiolysis	https://doi.org/10.1016/0003-9861(81)90263-0	not reported	differs from 1985 values; missing conditions prevent reconciliation; never k_D	validation only	parameter_records.d oxorubicin_semiquinone_plus_oxygen_1981
t_half_SQ_anaerobic	pulse radiolysis	https://doi.org/10.1016/0003-9861(81)90263-0	not reported	chemical loss not spin T1/T2 or encounter duration	context only	reference_evidence. encounter.chemical_lifetime_context
diffusion_distance_SQ	calculated from chemical lifetime/diffusion	https://doi.org/10.1016/0003-9861(81)90263-0	not reported	not pair separation or encounter radius; 8 us air-saturated chemical lifetime is not cage time	context only	reference_evidence. encounter.diffusion_distance_context
encounter_geometry_J_D_kassoc_ksep	targeted structure/EPR/kinetics search	NA	NA	not reported in inspected exact-system papers and not found elsewhere	sensitivity only	reference_evidence. encounter.geometry_exchange_dipolar_association_separation_duration
DQ_coherence	targeted transient-EPR; CIDEP/CIDNP; magnetic-field; product-branching search	NA	NA	retrieved exact-system studies did not investigate D/Q observables; no separate direct paper found	sensitivity only	reference_evidence. encounter.coherence
k_D	spin-resolved kinetics search	NA	NA	bulk M ⁻¹ s ⁻¹ total rates cannot be converted or decomposed	illustrative only	reference_evidence. encounter.k_D
k_Q	spin-resolved kinetics search	NA	NA	no state-resolved rate or channel yield	illustrative only	reference_evidence. encounter.k_Q
k_Q_over_k_D	spin-resolved kinetics search	NA	NA	no defensible physical range or ordering; 0/0.01/0.1/1 are inactive illustrative benchmarks only	illustrative only	reference_evidence. encounter.k_Q_over_k_D

Each parameter retains its stated conditions, evidence status, and permitted use. Presentation values are rounded. All 40 source rows and all 23 source columns remain in table02_parameter_provenance.csv; 7 provenance/support columns are source-only. 'Not available' is distinct from numeric zero.

Table 2. Parameter provenance

Panel C. Evidence and permitted use (continued)

Symbol	Method	Source	Uncertainty	Limitations	Permitted use	Code location
pKa_HO2	pulse radiolysis	https://doi.org/10.1021/j100711a009	0.1	buffer absence explains disagreement with earlier values; not a membrane/protein pKa	quantitative aqueous	reference_evidence.downstream.HO2_pKa
k_HH	pulse-radiolysis decay	https://doi.org/10.1021/j100711a009	not reported	species-resolved; do not call an O2- plus O2- rate	quantitative aqueous	reference_evidence.downstream.k_HO2_HO2
k_HA	pulse-radiolysis decay	https://doi.org/10.1021/j100711a009	not reported	requires explicit acid-base speciation	quantitative aqueous	reference_evidence.downstream.k_HO2_O2 minus
k_AA	pulse-radiolysis bound	https://doi.org/10.1021/j100711a009	not reported	2026 reevaluation gives tighter <0.3 M ⁻¹ s ⁻¹ ; common neutral-pH effective rates are not this elementary step	quantitative aqueous	reference_evidence.downstream.k_O2minus_s_O2minus
k_SOD_oxidation	pulse radiolysis optical 650/300 nm	https://doi.org/10.1021/ja00790a007	0.2e9 M ⁻¹ s ⁻¹	exact enzyme/preparation only; surrogate for human or mitochondrial SOD	sensitivity benchmark	reference_evidence.downstream.k_CuZnSO4_D_oxidation
k_SOD_reduction	pulse radiolysis optical 650/300 nm	https://doi.org/10.1021/ja00790a007	0.4e9 M ⁻¹ s ⁻¹	exact enzyme/preparation only; surrogate otherwise	sensitivity benchmark	reference_evidence.downstream.k_CuZnSO4_D_reduction
k_SOD_overall	pulse radiolysis plus optical/EPR	https://doi.org/10.1042/bj1390049	0.18e9 M ⁻¹ s ⁻¹	not a universal intracellular SOD1/SOD2 constant	sensitivity benchmark	reference_evidence.downstream.k_CuZnSO4_D_overall
v_O2minus_heart_SMP	SOD-inhibitable cytochrome-c reduction	https://doi.org/10.1016/S0021-9258(17)35747-2	0.2;2.7	90 uM dose; exact buffer oxygen and timing unavailable in accessible abstract	quantitative matched validation	validation_datasets.beef_heart_submitochondrial_superoxide
v_H2O2_heart_SMP	catalase-released oxygen	https://doi.org/10.1016/S0021-9258(17)35747-2	0.3	200 uM dose; detection limit and exact conditions not reported in accessible abstract	quantitative matched validation	validation_datasets.beef_heart_submitochondrial_H2O2

Each parameter retains its stated conditions, evidence status, and permitted use. Presentation values are rounded. All 40 source rows and all 23 source columns remain in table02_parameter_provenance.csv; 7 provenance/support columns are source-only. 'Not available' is distinct from numeric zero.

Table 2. Parameter provenance

Panel C. Evidence and permitted use (continued)

Symbol	Method	Source	Uncertainty	Limitations	Permitted use	Code location
v_O2minus_Ehrlich_microsomes	acetylated cytochrome-c reduction at 550 nm	https://doi.org/10.1155/2019/9474823	0.26;1.43	135 uM dose; initial linear rate; oxygenation not separately reported	quantitative matched validation	validation_datasets.Ehrlich_microsome_superoxide
v_O2minus_Ehrlich_mitochondria	acetylated cytochrome-c reduction at 550 nm	https://doi.org/10.1155/2019/9474823	0.15;0.61	135 uM dose; alkaline and rotenone-blocked; oxygenation not separately reported	quantitative matched validation	validation_datasets.Ehrlich_mitochondrial_superoxide
v_O2minus_Ehrlich_nuclei	acetylated cytochrome-c reduction at 550 nm	https://doi.org/10.1155/2019/9474823	0.05;0.33	135 uM dose; isolated fraction; oxygenation not separately reported	quantitative matched validation	validation_datasets.Ehrlich_nuclear_superoxide
v_H2O2_Ehrlich	catalase-released oxygen; linear 10-30 min interval	https://doi.org/10.1155/2019/9474823	0.48;0.6;0.04	135 uM fractions;400 uM cells; controls undetectable with detection limits not reported; PBS air-bubbled 30 min	quantitative matched validation	validation_datasets.Ehrlich_H2O2
H2O2_PC3	catalase-aminotriazole kinetic assay	https://doi.org/10.1016/j.abb.2005.06.015	4;13 pM	30 min exposure; oxygen fraction not reported; extracellular Amplex Red did not increase; catalase overexpression did not improve resistance	quantitative matched validation	validation_datasets.PC3_intracellular_H2O2
MitoSOX_H9c2	1 h fluorescence	https://doi.org/10.1016/j.bbrc.2007.04.106	0.04;0.08;0.08	10/20/50 uM doses; n=3; oxygen not reported; doxorubicin autofluorescence and nonspecific probe products prohibit absolute calibration	sensitivity only	validation_datasets.H9c2_MitoSOX
negative_5_imino	acetylated cytochrome-c reduction	https://doi.org/10.1155/2019/9474823	1.43;0.15	both 135 uM; supports quinone dependence but not D/Q selectivity	qualitative disconfirming	validation_datasets.quinone_disabled_negative_control

Each parameter retains its stated conditions, evidence status, and permitted use. Presentation values are rounded. All 40 source rows and all 23 source columns remain in table02_parameter_provenance.csv; 7 provenance/support columns are source-only. 'Not available' is distinct from numeric zero.