

19. ODEs III: Taylor methods and adaptivity – controlling the error

Summary of the previous class

- Trapezoid method: An implicit method
- Modified Euler method
- Runge–Kutta methods
- Stages as Euler steps

Goals for today

1 Taylor methods

- Taylor series solutions of ODEs

2 Adaptivity

- Controlling the error by varying the step size
- Embedded Runge–Kutta methods

Review: Runge–Kutta methods

- We want to solve

$$\dot{\mathbf{x}}(t) = \mathbf{f}(t, \mathbf{x}(t))$$

- To get approximations \mathbf{x}_n of solution $x(t_n)$ at time t_n

Review: Runge–Kutta methods

- We want to solve

$$\dot{\mathbf{x}}(t) = \mathbf{f}(t, \mathbf{x}(t))$$

- To get approximations \mathbf{x}_n of solution $x(t_n)$ at time t_n
- Runge–Kutta methods use several **stages**
- Each stage is an evaluation of f at some point
- Depending on previous evaluations

Runge–Kutta methods reproduce the Taylor series of the true solution

- Runge–Kutta methods calculate the Taylor series of the true solution $x(t)$ around the initial point x_n

Runge–Kutta methods reproduce the Taylor series of the true solution

- Runge–Kutta methods calculate the Taylor series of the true solution $x(t)$ around the initial point x_n
- Without actually calculating the Taylor series!

Runge–Kutta methods reproduce the Taylor series of the true solution

- Runge–Kutta methods calculate the Taylor series of the true solution $x(t)$ around the initial point x_n
- Without actually calculating the Taylor series!
- The amazing thing is that a given RK method works for *any* (smooth enough) f

Runge–Kutta methods reproduce the Taylor series of the true solution

- Runge–Kutta methods calculate the Taylor series of the true solution $x(t)$ around the initial point x_n
- Without actually calculating the Taylor series!
- The amazing thing is that a given RK method works for *any* (smooth enough) f
- However, for high accuracy we need high-order methods

Runge–Kutta methods reproduce the Taylor series of the true solution

- Runge–Kutta methods calculate the Taylor series of the true solution $x(t)$ around the initial point x_n
- Without actually calculating the Taylor series!
- The amazing thing is that a given RK method works for *any* (smooth enough) f
- However, for high accuracy we need high-order methods
- E.g. to simulate the solar system for millions of years

Runge–Kutta methods reproduce the Taylor series of the true solution

- Runge–Kutta methods calculate the Taylor series of the true solution $x(t)$ around the initial point x_n
- Without actually calculating the Taylor series!
- The amazing thing is that a given RK method works for *any* (smooth enough) f
- However, for high accuracy we need high-order methods
- E.g. to simulate the solar system for millions of years
- RK methods of high order are difficult to find

Taylor methods

Taylor series solutions

- **Taylor methods** use the Taylor series of the exact solution
- We have seen that they seem to require calculating high-order derivatives of f , such as f_t , f_x , f_{xx}

Taylor series solutions

- **Taylor methods** use the Taylor series of the exact solution
- We have seen that they seem to require calculating high-order derivatives of f , such as f_t , f_x , f_{xx}
- But it is possible to construct the Taylor series solution directly (up to a given order)!

Taylor series solutions

- **Taylor methods** use the Taylor series of the exact solution
- We have seen that they seem to require calculating high-order derivatives of f , such as f_t , f_x , f_{xx}
- But it is possible to construct the Taylor series solution directly (up to a given order)!
- The disadvantage is that we must construct a new Taylor series solution for each ODE

Taylor method

- Suppose we want to solve $\dot{x}(t) = f(x(t))$, with $x(t=0) = x_0$

Taylor method

- Suppose we want to solve $\dot{x}(t) = f(x(t))$, with $x(t=0) = x_0$
- Suppose f is **analytic**

Taylor method

- Suppose we want to solve $\dot{x}(t) = f(x(t))$, with $x(t=0) = x_0$
- Suppose f is **analytic**
- i.e. is equal to its Taylor expansion

$$f(x) = \tilde{f}_0 + \tilde{f}_1 x + \tilde{f}_2 x^2 + \dots$$

Taylor method

- Suppose we want to solve $\dot{x}(t) = f(x(t))$, with $x(t=0) = x_0$
- Suppose f is **analytic**
- i.e. is equal to its Taylor expansion

$$f(x) = \tilde{f}_0 + \tilde{f}_1 x + \tilde{f}_2 x^2 + \dots$$

- Then (theorem) the solution $x(t)$ exists and is analytic, so has a power series expansion in t :

$$x(t) = x_0 + x_1 t + x_2 t^2 + \dots$$

Taylor method II

■ We have

$$x(0) = x_0$$

$$\dot{x}(0) = x_1$$

$$\ddot{x}(0) = 2x_2$$

⋮

⋯

■ How can we calculate the x_i ?

Collaboration I

Taylor method

We are proposing a solution $x(t) = x_0 + x_1 t + x_2 t^2 + \dots$ with unknown coefficients x_i for the ODE $\dot{x} = f(x)$ with given initial condition $x(t) = x_0$.

- 1 How could we solve for the coefficients x_i ?
- 2 Try to do this for $f(x) = x$

Taylor method III

- Let's **substitute** the Taylor series for $x(t)$ into the ODE:
 - on the left-hand side we need $\dot{x}(t)$
 - on the right-hand side we need $f(x(t))$

Taylor method III

- Let's **substitute** the Taylor series for $x(t)$ into the ODE:
 - on the left-hand side we need $\dot{x}(t)$
 - on the right-hand side we need $f(x(t))$
- Both of these give *new Taylor series*:

$$\dot{x}(t) = x_1 + 2x_2t + 3x_3t^2 + \dots$$

- Substituting $x(t)$ into $f(x(t))$ gives a series *in* t :

$$f(x(t)) = f_0 + f_1t + f_2t^2 + \dots$$

Taylor method IV

- Since these two series are equal for all t
the coefficient of t^n must be equal for each n

Taylor method IV

- Since these two series are equal for all t
the coefficient of t^n must be equal for each n
- To prove this e.g. differentiate repeatedly

Taylor method IV

- Since these two series are equal for all t
the coefficient of t^n must be equal for each n
- To prove this e.g. differentiate repeatedly
- Equate coefficients of each power t^n :

$$\begin{aligned}x_1 &= f_0 \\2x_2 &= f_1 \\&\vdots \\nx_n &= f_{n-1}\end{aligned}$$

Taylor method IV

- Since these two series are equal for all t
the coefficient of t^n must be equal for each n
- To prove this e.g. differentiate repeatedly
- Equate coefficients of each power t^n :

$$\begin{aligned}x_1 &= f_0 \\2x_2 &= f_1 \\&\vdots \\nx_n &= f_{n-1}\end{aligned}$$

- We get a **recurrence relation**: $x_n = \frac{f_{n-1}}{n}$

Taylor method V

- To calculate f_{n-1} we insert the Taylor series for $x(t)$ into $f(x)$

Taylor method V

- To calculate f_{n-1}' we insert the Taylor series for $x(t)$ into $f(x)$
- f_{n-1}' is the coefficient of t^{n-1}

Taylor method V

- To calculate f_{n-1} we insert the Taylor series for $x(t)$ into $f(x)$
- f_{n-1} is the coefficient of t^{n-1}
- So f_{n-1} can depend only on x_0 up to x_{n-1}

Taylor method V

- To calculate f_{n-1} we insert the Taylor series for $x(t)$ into $f(x)$
- f_{n-1} is the coefficient of t^{n-1}
- So f_{n-1} can depend only on x_0 up to x_{n-1}
- Hence from the coefficients x_0, \dots, x_{n-1} we can obtain $x_n!$

Taylor method V

- To calculate f_{n-1} we insert the Taylor series for $x(t)$ into $f(x)$
- f_{n-1} is the coefficient of t^{n-1}
- So f_{n-1} can depend only on x_0 up to x_{n-1}
- Hence from the coefficients x_0, \dots, x_{n-1} we can obtain $x_n!$
- We generate all coefficients x_n *recursively*, one by one!

Example

- E.g. Solve $\dot{x} = x^2$ with $x_0 = 1$

Example

- E.g. Solve $\dot{x} = x^2$ with $x_0 = 1$
- Start with all coefficients unknown *except* x_0 :

$$x(t) = x_0 + \textcolor{red}{x}_1 t + \textcolor{red}{x}_2 t^2 + \dots$$

where **red** denotes coefficients we don't yet know

Example

- E.g. Solve $\dot{x} = x^2$ with $x_0 = 1$
- Start with all coefficients unknown *except* x_0 :

$$x(t) = x_0 + \textcolor{red}{x}_1 t + \textcolor{red}{x}_2 t^2 + \dots$$

where **red** denotes coefficients we don't yet know

- So

$$\begin{aligned} f(x(t)) &= [x(t)]^2 \\ &= (x_0 + \textcolor{red}{x}_1 t + \dots)^2 \\ &= x_0^2 + \mathcal{O}(t) \end{aligned}$$

- Hence $f_0 = x_0^2$

Example II

- So $x_1 = f_0/1 = x_0^2$

Example II

- So $x_1 = f_0/1 = x_0^2$
- Now we have $x(t) = x_0 + x_1 t + x_2 t^2 + \dots$

Example II

- So $x_1 = f_0/1 = x_0^2$
- Now we have $x(t) = x_0 + x_1t + x_2t^2 + \dots$
- Substitute into $f(x)$ again:

$$f(x(t)) = x_0^2 + t(x_0x_1 + x_1x_0) + \mathcal{O}(t^2)$$

Example II

- So $x_1 = f_0/1 = x_0^2$
- Now we have $x(t) = x_0 + x_1t + x_2t^2 + \dots$
- Substitute into $f(x)$ again:

$$f(x(t)) = x_0^2 + t(x_0x_1 + x_1x_0) + \mathcal{O}(t^2)$$

- This gives $f_1 = (\text{coefficient of } t) = 2x_0x_1$

Example II

- So $x_1 = f_0/1 = x_0^2$
- Now we have $x(t) = x_0 + x_1 t + x_2 t^2 + \dots$
- Substitute into $f(x)$ again:

$$f(x(t)) = x_0^2 + t(x_0 x_1 + x_1 x_0) + \mathcal{O}(t^2)$$

- This gives $f_1 = (\text{coefficient of } t) = 2x_0 x_1$
- Hence $x_2 = f_1/2 = x_0 x_1$

Example II

- So $x_1 = f_0/1 = x_0^2$
- Now we have $x(t) = x_0 + x_1 t + x_2 t^2 + \dots$
- Substitute into $f(x)$ again:

$$f(x(t)) = x_0^2 + t(x_0 x_1 + x_1 x_0) + \mathcal{O}(t^2)$$

- This gives $f_1 = (\text{coefficient of } t) = 2x_0 x_1$
- Hence $x_2 = f_1/2 = x_0 x_1$
- Repeat, including the new coefficient into $x(t)$
- Note: previous f_i are recalculated – this is inefficient

Alternative viewpoint: Integrals

- An alternative viewpoint is to use the integral formulation of the ODE:

$$x(t) = x_0 + \int_0^t f(x(s)) ds$$

- Define the n th order polynomial approximation $x^{(n)}(t) := x_0 + \dots + x_n t^n$
- We use **Picard iteration** to calculate $x^{(n)}$ recursively:

$$x^{(n+1)} = x_0 + \int \hat{f}^{(n)}(x^{(n)})$$

- Where $\hat{f}^{(n)}(x)$ means “ $f(x)$ truncated to order n ”

Example using integrals

- Let's return to our example:

$$\dot{x} = x^2 \quad \text{with } x^{(0)} := x_0$$

Example using integrals

- Let's return to our example:

$$\dot{x} = x^2 \quad \text{with } x^{(0)} := x_0$$

$$x^{(1)} = x_0 + \int (x^{(0)})^2 = x_0 + \int_0^t x_0^2 ds = x_0 + t x_0^2$$

Example using integrals

- Let's return to our example:

$$\dot{x} = x^2 \quad \text{with } x^{(0)} := x_0$$

$$x^{(1)} = x_0 + \int (x^{(0)})^2 = x_0 + \int_0^t x_0^2 ds = x_0 + t x_0^2$$

$$x^{(2)} = x_0 + \int (x^{(1)})^2$$

$$= x_0 + \int_0^t (x_0 + s x_0^2)^2 ds = x_0 + t x_0^2 + t^2 x_0^3 + \mathcal{O}(t^3)$$

Implementation

- How can we automate this in Julia?
- What operations do we need?

Implementation

- How can we automate this in Julia?
- What operations do we need?
- We are just manipulating polynomials!
- And truncating to a given degree
- So define operations like $*$ on polynomials of degree n that return polynomials of the *same* degree

Implementation

- How can we automate this in Julia?
- What operations do we need?
- We are just manipulating polynomials!
- And truncating to a given degree
- So define operations like $*$ on polynomials of degree n that return polynomials of the *same* degree
- These manipulations are done with *numeric* coefficients

Implementation

- How can we automate this in Julia?
- What operations do we need?
- We are just manipulating polynomials!
- And truncating to a given degree
- So define operations like $*$ on polynomials of degree n that return polynomials of the *same* degree
- These manipulations are done with *numeric* coefficients
- In the end, much of “symbolic” calculation manipulating numeric coefficients of polynomials

Adaptivity: Varying the step size and
controlling the error

Error checking

- Until now we have supposed that each step approximates the function well
- We have always taken the calculated step

Error checking

- Until now we have supposed that each step approximates the function well
- We have always taken the calculated step
- But we are stepping into the unknown, so we should **verify** if the step is “valid”

Error checking

- Until now we have supposed that each step approximates the function well
- We have always taken the calculated step
- But we are stepping into the unknown, so we should **verify** if the step is “valid”
- I.e. if the error is “small enough”

Error checking

- Until now we have supposed that each step approximates the function well
- We have always taken the calculated step
- But we are stepping into the unknown, so we should **verify** if the step is “valid”
- I.e. if the error is “small enough”
- But what can we check against?
- The exact solution is never available

Collaboration II

Checking Euler steps

Suppose we are using the Euler method and we take one step.

- 1 What could we check the step against if we don't have access to the true solution?
- 2 Can you think of another possibility (for a total of two)?
- 3 How much computational effort would that require?
- 4 How can you decide whether the error is small enough?

Same method, different step sizes

- One solution is to use the *same* numerical method, e.g. Euler

Same method, different step sizes

- One solution is to use the *same* numerical method, e.g. Euler
- But with *different* step sizes

Same method, different step sizes

- One solution is to use the *same* numerical method, e.g. Euler
- But with *different* step sizes
- y_1 := result after *one* Euler step of length h
- y_2 := result after *two* consecutive Euler steps, each of length $h/2$

Same method, different step sizes

- One solution is to use the *same* numerical method, e.g. Euler
- But with *different* step sizes
- $y_1 :=$ result after *one* Euler step of length h
- $y_2 :=$ result after *two* consecutive Euler steps, each of length $h/2$
- Local error = $\mathcal{O}(h^2)$ for *both*
- But the constant is different (but related)
- $\Delta y := |y_1 - y_2|$ *measures* the error

Same method, different step sizes

- One solution is to use the *same* numerical method, e.g. Euler
- But with *different* step sizes
- y_1 := result after *one* Euler step of length h
- y_2 := result after *two* consecutive Euler steps, each of length $h/2$
- Local error = $\mathcal{O}(h^2)$ for *both*
- But the constant is different (but related)
- $\Delta y := |y_1 - y_2|$ *measures* the error
- See the problem set

Alternative: Different methods with different order

- $y_1 :=$ result after a step with a method of order p
- $y_2 :=$ result after a step with a method of order $(p + 1)$

Alternative: Different methods with different order

- $y_1 :=$ result after a step with a method of order p
- $y_2 :=$ result after a step with a method of order $(p + 1)$
- How big will the error be?

Alternative: Different methods with different order

- $y_1 :=$ result after a step with a method of order p
- $y_2 :=$ result after a step with a method of order $(p + 1)$
- How big will the error be?
- If the step size is h then $\Delta y := |y_1 - y_2| = Ch^{p+1}$

Alternative: Different methods with different order

- $y_1 :=$ result after a step with a method of order p
- $y_2 :=$ result after a step with a method of order $(p + 1)$
- How big will the error be?
- If the step size is h then $\Delta y := |y_1 - y_2| = Ch^{p+1}$
- Note that C is related to a higher derivative, but is *unknown*
- This measures the approximate error in y_1 , since y_2 is presumably more accurate

Collaboration III

Choosing a step size

Suppose we want the error to be $\leq \epsilon$

- 1 How can we decide whether the step should be accepted?
- 2 If it is not accepted, what should we do?
- 3 How should we change things?

Varying the step size

- Suppose we want the error to be $\leq \epsilon$

Varying the step size

- Suppose we want the error to be $\leq \epsilon$
- Then we should *choose* the step size h' accordingly

Varying the step size

- Suppose we want the error to be $\leq \epsilon$
- Then we should *choose* the step size h' accordingly
- We need $C(h')^{p+1} \sim \epsilon$

Varying the step size

- Suppose we want the error to be $\leq \epsilon$
- Then we should *choose* the step size h' accordingly
- We need $C(h')^{p+1} \sim \epsilon$
- We can get rid of C by dividing!:

$$\left(\frac{h'}{h}\right)^{p+1} = \frac{\epsilon}{\Delta y}$$

- So we should take

$$h' = h \left(\frac{\epsilon}{\Delta y}\right)^{\frac{1}{p+1}}$$

- Alternative: “error per unit time step” should be ϵ

Variable step size algorithm

- We can use this to construct an algorithm with **variable step size**:
- 1 Propose a step and calculate the error Δy as above

Variable step size algorithm

- We can use this to construct an algorithm with **variable step size**:
 - 1 Propose a step and calculate the error Δy as above
 - 2 If the error is *too big*, $\Delta y > \epsilon$, then we **reject** the step:
we remain at the same place, but *decrease* the step size h

Variable step size algorithm

- We can use this to construct an algorithm with **variable step size**:
 - 1 Propose a step and calculate the error Δy as above
 - 2 If the error is *too big*, $\Delta y > \epsilon$, then we **reject** the step:
we remain at the same place, but *decrease* the step size h
 - 3 If error is *small enough*, $\Delta y \leq \epsilon$, then we **accept** the step:
we move with the *current* step size h , then *increase* h

Variable step size algorithm II

-
-
-
- 4 In both cases, the step size gets modified

Variable step size algorithm II

- 4 In both cases, the step size gets modified
 - In certain circumstances this can lead to wild increases of h
 - So we restrict to at most $h' = 2h$

Variable step size algorithm II

- 4 In both cases, the step size gets modified
 - In certain circumstances this can lead to wild increases of h
 - So we restrict to at most $h' = 2h$
 - This allows even Euler to “work”
 - But it can lead to taking many tiny steps

Embedded Runge–Kutta methods

- The above methods require too much computational work

Embedded Runge–Kutta methods

- The above methods require too much computational work
- E.g. Euler methods need *three* function evaluations for each step for an $\mathcal{O}(h)$ method

Embedded Runge–Kutta methods

- The above methods require too much computational work
- E.g. Euler methods need *three* function evaluations for each step for an $\mathcal{O}(h)$ method
- This becomes even worse with p - and $p + 1$ -order methods: we need at least $2p + 1$ function evaluations

Embedded Runge–Kutta methods

- The above methods require too much computational work
- E.g. Euler methods need *three* function evaluations for each step for an $\mathcal{O}(h)$ method
- This becomes even worse with p - and $p + 1$ -order methods: we need at least $2p + 1$ function evaluations
- Amazingly, with Runge–Kutta methods it is possible to find a better solution

Embedded Runge–Kutta methods II

- Suppose we use an order- $(p + 1)$ RK method with s stages

Embedded Runge–Kutta methods II

- Suppose we use an order- $(p + 1)$ RK method with s stages
- Then $x_{n+1} = x_n + h \sum_{i=1}^s b_i k_i$, where the k_i are the results of each stage

Embedded Runge–Kutta methods II

- Suppose we use an order- $(p + 1)$ RK method with s stages
- Then $x_{n+1} = x_n + h \sum_{i=1}^s b_i k_i$, where the k_i are the results of each stage
- Amazingly, for some RK methods there is a linear combination of the *same* k_i 's giving an order- p method

Embedded Runge–Kutta methods III

- Example: Bogacki–Shampine BS23:

0				
$\frac{1}{2}$	$\frac{1}{2}$			
$\frac{3}{4}$	0	$\frac{3}{4}$		
$\frac{1}{4}$	$\frac{2}{9}$	$\frac{1}{3}$	$\frac{4}{9}$	
1	$\frac{2}{9}$	$\frac{1}{3}$	$\frac{4}{9}$	0
	$\frac{7}{24}$	$\frac{1}{4}$	$\frac{1}{3}$	$\frac{1}{8}$

Embedded Runge–Kutta methods III

- Example: Bogacki–Shampine BS23:

0				
$\frac{1}{2}$	$\frac{1}{2}$			
$\frac{3}{4}$	0	$\frac{3}{4}$		
$\frac{1}{4}$	$\frac{2}{9}$	$\frac{1}{3}$	$\frac{4}{9}$	
1	$\frac{2}{9}$	$\frac{1}{3}$	$\frac{4}{9}$	0
	$\frac{7}{24}$	$\frac{1}{4}$	$\frac{1}{3}$	$\frac{1}{8}$

- The last two lines give 3rd- and 2nd-order methods, respectively

Embedded Runge–Kutta methods III

- Example: Bogacki–Shampine BS23:

0				
$\frac{1}{2}$	$\frac{1}{2}$			
$\frac{3}{4}$	0	$\frac{3}{4}$		
$\frac{1}{4}$	$\frac{2}{9}$	$\frac{1}{3}$	$\frac{4}{9}$	
1	$\frac{2}{9}$	$\frac{1}{3}$	$\frac{4}{9}$	0
	$\frac{7}{24}$	$\frac{1}{4}$	$\frac{1}{3}$	$\frac{1}{8}$

- The last two lines give 3rd- and 2nd-order methods, respectively
- We need about 3 function evaluations per step for this embedded 2nd / 3rd-order method

Embedded Runge–Kutta methods IV

- An additional useful property is **FSAL**: “First same as last”

Embedded Runge–Kutta methods IV

- An additional useful property is **FSAL**: “First same as last”
- If k_s is evaluated at $t_n + h$
- Then $(k_s \text{ from the previous step}) = (k_1 \text{ for the new step})$

Embedded Runge–Kutta methods IV

- An additional useful property is **FSAL**: “First same as last”
- If k_s is evaluated at $t_n + h$
- Then $(k_s \text{ from the previous step}) = (k_1 \text{ for the new step})$
- So there is no need to re-evaluate

Embedded Runge–Kutta methods IV

- An additional useful property is **FSAL**: “First same as last”
- If k_s is evaluated at $t_n + h$
- Then $(k_s$ from the previous step) = $(k_1$ for the new step)
- So there is no need to re-evaluate
- Modern version: Tsitouras 5/4 method (2011)
- Default in `DifferentialEquations.jl`

Summary

- We can generate Taylor methods of arbitrary order
 - By recursively calculating the coefficients
 - Using polynomial manipulation
-
- We can calculate the local error by running two different methods for the same step
 - We *choose* a variable step size to obtain the desired error
 - Embedded Runge–Kutta methods provide an efficient way to do this