

12. Linear algebra and data: Least squares and the Singular Value Decomposition (SVD)

Summary of last lecture

- Power iteration $\mathbf{x}_{n+1} = \mathbf{A} \mathbf{x}_n$
- Finds eigenvectors $\mathbf{A} \mathbf{x} = \lambda \mathbf{x}$
- Gram–Schmidt orthogonalisation
- QR factorisation
 - Orthogonal matrix \mathbf{Q} such that $\mathbf{Q}^T \mathbf{Q} = \mathbf{I}$
 - Upper-triangular matrix \mathbf{R}

Goals for today

- Fitting functions to data
- Optimization problems
- Linear least-squares problems
- Solution using linear algebra
- The Singular-Value Decomposition (SVD)

Least squares: Fitting functions to data

Parametric functions

- Suppose we have data points (x_i, y_i)

Parametric functions

- Suppose we have data points (x_i, y_i)
- We suppose they can be fitted by a **model** $y = f(x)$

Parametric functions

- Suppose we have data points (x_i, y_i)
- We suppose they can be fitted by a **model** $y = f(x)$
- I.e. we suspect that they are related as

$$y_i = f_{\mathbf{p}}(x_i) + \epsilon_i$$

- With **noise** (randomness) ϵ_i on observation i

Parametric functions

- Suppose we have data points (x_i, y_i)
- We suppose they can be fitted by a **model** $y = f(x)$
- I.e. we suspect that they are related as

$$y_i = f_{\mathbf{p}}(x_i) + \epsilon_i$$

- With **noise** (randomness) ϵ_i on observation i
- $f_{\mathbf{p}}$ is a **parametric function** with parameter vector \mathbf{p}

Parametric functions

- Suppose we have data points (x_i, y_i)
- We suppose they can be fitted by a **model** $y = f(x)$
- I.e. we suspect that they are related as

$$y_i = f_{\mathbf{p}}(x_i) + \epsilon_i$$

- With **noise** (randomness) ϵ_i on observation i
- $f_{\mathbf{p}}$ is a **parametric function** with parameter vector \mathbf{p}
- Simplest case: straight line $f_{\alpha,\beta}(x) = \alpha + \beta x$

Collaboration I

Finding the best-fitting straight line

Suppose we have data (x_i, y_i) and a model $f(x) = \alpha + \beta x$.

- 1 How can we write down the problem to find the straight line $y = f(x)$ that “fits” the data best?
- 2 What type of problem is it?
- 3 How could we solve it?

Model of best fit

- We want to find the model that **best fits** the data
- How can we formalize this?

Model of best fit

- We want to find the model that **best fits** the data
- How can we formalize this?
- We want to find the values of the parameters **p** that **minimize** the “distance” of the data from the function

Model of best fit

- We want to find the model that **best fits** the data
- How can we formalize this?
- We want to find the values of the parameters \mathbf{p} that **minimize** the “distance” of the data from the function
- To measure the distance we introduce a **loss function** or **cost function**

Model of best fit

- We want to find the model that **best fits** the data
- How can we formalize this?
- We want to find the values of the parameters \mathbf{p} that **minimize** the “distance” of the data from the function
- To measure the distance we introduce a **loss function** or **cost function**
- This will measure the total “distance” between *all* the data and the function

Least squares problems

■ **Loss** function:

$$\mathcal{L}(\mathbf{p}) = \sum_i [y_i - f_{\mathbf{p}}(t_i)]^2$$

Least squares problems

- **Loss** function:

$$\mathcal{L}(\mathbf{p}) = \sum_i [y_i - f_{\mathbf{p}}(t_i)]^2$$

- We want to minimise this over all vectors of parameters \mathbf{p}
- E.g. over all α and β

Least squares problems

- **Loss** function:

$$\mathcal{L}(\mathbf{p}) = \sum_i [y_i - f_{\mathbf{p}}(t_i)]^2$$

- We want to minimise this over all vectors of parameters \mathbf{p}
- E.g. over all α and β
- **Least-squares problem**

Optimization

- This is an example of an **optimization** problem

Optimization

- This is an example of an **optimization** problem
- Optimization problems are usually **very hard**; huge field

Optimization

- This is an example of an **optimization** problem
- Optimization problems are usually **very hard**; huge field
- There are special methods for problems with certain **structure**

Optimization

- This is an example of an **optimization** problem
- Optimization problems are usually **very hard**; huge field
- There are special methods for problems with certain **structure**

- There are many numerical methods for optimization
- Sometimes analytical solutions are possible – e.g. linear least squares

Matrix formulation of linear least squares

- We want to minimise $\mathcal{L}(\mathbf{p}) = \sum_i r_i^2$
- Where $r_i := y_i - f_{\mathbf{p}}(t_i)$ is the i th **residual**

Matrix formulation of linear least squares

- We want to minimise $\mathcal{L}(\mathbf{p}) = \sum_i r_i^2$
- Where $r_i := y_i - f_{\mathbf{p}}(t_i)$ is the i th **residual**
- Look for a matrix formulation: $\mathcal{L} = \sum_i r_i^2 = \mathbf{r}^\top \mathbf{r}$

Matrix formulation of linear least squares

- We want to minimise $\mathcal{L}(\mathbf{p}) = \sum_i r_i^2$
- Where $r_i := y_i - f_{\mathbf{p}}(t_i)$ is the i th **residual**
- Look for a matrix formulation: $\mathcal{L} = \sum_i r_i^2 = \mathbf{r}^\top \mathbf{r}$
- The vector \mathbf{r} of all residuals is given by

$$\mathbf{r} = \begin{bmatrix} r_1 \\ r_2 \\ \vdots \\ r_m \end{bmatrix} = \begin{bmatrix} 1 & x_1 \\ 1 & x_2 \\ \vdots & \vdots \\ 1 & x_m \end{bmatrix} \begin{bmatrix} \alpha \\ \beta \end{bmatrix} - \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_m \end{bmatrix} =: \mathbf{A}\mathbf{x} - \mathbf{b}$$

where $\mathbf{x} = \begin{bmatrix} \alpha \\ \beta \end{bmatrix}$ are the unknowns

Matrix formulation II

- Ideally we would like to solve $A\mathbf{x} = \mathbf{b}$

Matrix formulation II

- Ideally we would like to solve $A\mathbf{x} = \mathbf{b}$
- But we have an **overdetermined system**:
- There are more equations (n) than unknowns (2)

Matrix formulation II

- Ideally we would like to solve $A\mathbf{x} = \mathbf{b}$
- But we have an **overdetermined system**:
- There are more equations (n) than unknowns (2)
- What can we do?

Matrix formulation II

- Ideally we would like to solve $A\mathbf{x} = \mathbf{b}$
- But we have an **overdetermined system**:
- There are more equations (n) than unknowns (2)
- What can we do?
- The best we can do is to *minimize* $\mathbf{r}^T \mathbf{r}$

General linear least squares

- General case: want to “solve” $A\mathbf{x} = \mathbf{b}$
- For a **tall and narrow** matrix A , i.e. with $m > n$

General linear least squares

- General case: want to “solve” $A\mathbf{x} = \mathbf{b}$
- For a **tall and narrow** matrix A , i.e. with $m > n$
- We try to minimize

$$\|A\mathbf{x} - \mathbf{b}\|^2$$

General linear least squares

- General case: want to “solve” $A\mathbf{x} = \mathbf{b}$
- For a **tall and narrow** matrix A , i.e. with $m > n$
- We try to minimize

$$\|A\mathbf{x} - \mathbf{b}\|^2$$

- Example: fit a polynomial of degree $< n$ to $(n + 1)$ points

Geometrical interpretation

- Suppose \mathbf{x} ranges over all of \mathbb{R}^n
- Then $A\mathbf{x}$ ranges over the column space / range of A

Geometrical interpretation

- Suppose \mathbf{x} ranges over all of \mathbb{R}^n
- Then $A\mathbf{x}$ ranges over the column space / range of A
- The **column space** of A is the vector space spanned by the columns of A

Geometrical interpretation

- Suppose \mathbf{x} ranges over all of \mathbb{R}^n
- Then $A\mathbf{x}$ ranges over the column space / range of A
- The **column space** of A is the vector space spanned by the columns of A
- The column space is a hyperplane

Geometrical interpretation

- Suppose \mathbf{x} ranges over all of \mathbb{R}^n
- Then $A\mathbf{x}$ ranges over the column space / range of A
- The **column space** of A is the vector space spanned by the columns of A
- The column space is a hyperplane
- We are looking for an \mathbf{x} whose image $A\mathbf{x}$ is **closest** to \mathbf{b}

Collaboration II

Solving the least squares problem

We want \mathbf{x}^* to be the solution that minimises $\|\mathbf{Ax} - \mathbf{b}\|^2$.

- 1 Draw a sketch of this. Geometrically / intuitively, what condition should we satisfy?
- 2 How could we solve this problem?

Solving linear least squares

- We want to find \mathbf{x}^* minimising $\|\mathbf{Ax} - \mathbf{b}\|^2$

Solving linear least squares

- We want to find \mathbf{x}^* minimising $\|\mathbf{Ax} - \mathbf{b}\|^2$
- Intuitively this occurs when $\mathbf{r} := \mathbf{Ax}^* - \mathbf{b}$ is **perpendicular** to the column space

Solving linear least squares

- We want to find \mathbf{x}^* minimising $\|\mathbf{Ax} - \mathbf{b}\|^2$
- Intuitively this occurs when $\mathbf{r} := \mathbf{Ax}^* - \mathbf{b}$ is **perpendicular** to the column space
- Recall: $\|\mathbf{x}\|^2 = \mathbf{x}^\top \mathbf{x}$

Solving linear least squares

- We want to find \mathbf{x}^* minimising $\|\mathbf{A}\mathbf{x} - \mathbf{b}\|^2$
- Intuitively this occurs when $\mathbf{r} := \mathbf{A}\mathbf{x}^* - \mathbf{b}$ is **perpendicular** to the column space
- Recall: $\|\mathbf{x}\|^2 = \mathbf{x}^T \mathbf{x}$
- Let's look at other vectors in the subspace by taking any displacement \mathbf{y}
- Look at $\|\mathbf{A}(\mathbf{x}^* + \mathbf{y}) - \mathbf{b}\|^2$

Solving linear least squares II

- We have

$$\|A(\mathbf{x}^* + \mathbf{y}) - \mathbf{b}\|^2$$

$$= \|(\mathbf{Ax}^* - \mathbf{b}) + \mathbf{Ay}\|^2 = [(\mathbf{Ax}^* - \mathbf{b}) + \mathbf{Ay}]^T [(\mathbf{Ax}^* - \mathbf{b}) + \mathbf{Ay}]$$

Solving linear least squares II

- We have

$$\|A(\mathbf{x}^* + \mathbf{y}) - \mathbf{b}\|^2$$

$$= \|(\mathbf{Ax}^* - \mathbf{b}) + \mathbf{Ay}\|^2 = [(\mathbf{Ax}^* - \mathbf{b}) + \mathbf{Ay}]^T [(\mathbf{Ax}^* - \mathbf{b}) + \mathbf{Ay}]$$

$$= (\mathbf{Ax}^* - \mathbf{b})^T (\mathbf{Ax}^* - \mathbf{b}) + 2\mathbf{y}^T \mathbf{A}^T (\mathbf{Ax}^* - \mathbf{b}) + (\mathbf{Ay})^T (\mathbf{Ay})$$

Solving linear least squares II

- We have

$$\|A(\mathbf{x}^* + \mathbf{y}) - \mathbf{b}\|^2$$

$$= \|(\mathbf{Ax}^* - \mathbf{b}) + \mathbf{Ay}\|^2 = [(\mathbf{Ax}^* - \mathbf{b}) + \mathbf{Ay}]^T [(\mathbf{Ax}^* - \mathbf{b}) + \mathbf{Ay}]$$

$$= (\mathbf{Ax}^* - \mathbf{b})^T (\mathbf{Ax}^* - \mathbf{b}) + 2\mathbf{y}^T \mathbf{A}^T (\mathbf{Ax}^* - \mathbf{b}) + (\mathbf{Ay})^T (\mathbf{Ay})$$

- Here we used $\mathbf{y}^T \mathbf{z} = \mathbf{z}^T \mathbf{y}$, so $\mathbf{y}^T \mathbf{z} + \mathbf{z}^T \mathbf{y} = 2\mathbf{y}^T \mathbf{z}$

$$= \|\mathbf{Ax}^* - \mathbf{b}\|^2 + \|\mathbf{Ay}\|^2 + 2\mathbf{y}^T \mathbf{A}^T (\mathbf{Ax}^* - \mathbf{b})$$

Solving linear least squares II

- We have $\|A\mathbf{x}^* - \mathbf{b}\|^2 + \|A\mathbf{y}\|^2 + 2\mathbf{y}^T A^T (A\mathbf{x}^* - \mathbf{b})$

Solving linear least squares II

- We have $\|A\mathbf{x}^* - \mathbf{b}\|^2 + \|A\mathbf{y}\|^2 + 2\mathbf{y}^T A^T (A\mathbf{x}^* - \mathbf{b})$
- Take \mathbf{x}^* such that $A^T (A\mathbf{x}^* - \mathbf{b}) = \mathbf{0}$

Solving linear least squares II

- We have $\|A\mathbf{x}^* - \mathbf{b}\|^2 + \|A\mathbf{y}\|^2 + 2\mathbf{y}^T A^T (A\mathbf{x}^* - \mathbf{b})$
- Take \mathbf{x}^* such that $A^T (A\mathbf{x}^* - \mathbf{b}) = \mathbf{0}$
- Then for any \mathbf{y} we have

$$\|A(\mathbf{x}^* + \mathbf{y}) - \mathbf{b}\|^2 \geq \|A\mathbf{x}^* - \mathbf{b}\|^2$$

Solving linear least squares II

- We have $\|A\mathbf{x}^* - \mathbf{b}\|^2 + \|A\mathbf{y}\|^2 + 2\mathbf{y}^T A^T(A\mathbf{x}^* - \mathbf{b})$
- Take \mathbf{x}^* such that $A^T(A\mathbf{x}^* - \mathbf{b}) = \mathbf{0}$
- Then for any \mathbf{y} we have

$$\|A(\mathbf{x}^* + \mathbf{y}) - \mathbf{b}\|^2 \geq \|A\mathbf{x}^* - \mathbf{b}\|^2$$

- So \mathbf{x}^* is the minimizer !

Solving linear least squares II

- We have $\|A\mathbf{x}^* - \mathbf{b}\|^2 + \|A\mathbf{y}\|^2 + 2\mathbf{y}^T A^T (A\mathbf{x}^* - \mathbf{b})$
- Take \mathbf{x}^* such that $A^T (A\mathbf{x}^* - \mathbf{b}) = \mathbf{0}$
- Then for any \mathbf{y} we have

$$\|A(\mathbf{x}^* + \mathbf{y}) - \mathbf{b}\|^2 \geq \|A\mathbf{x}^* - \mathbf{b}\|^2$$

- So \mathbf{x}^* is the minimizer !
- Hence the unique solution of the least squares problem is given by the solution \mathbf{x} of

$$A^T A \mathbf{x} = A^T \mathbf{b}$$

Solving linear least squares III

- We have reduced the linear least squares problem to solving a linear system:

$$A^T A \mathbf{x} = A^T \mathbf{b}$$

Solving linear least squares III

- We have reduced the linear least squares problem to solving a linear system:

$$A^T A \mathbf{x} = A^T \mathbf{b}$$

- Called the **normal equations**

Solving linear least squares III

- We have reduced the linear least squares problem to solving a linear system:

$$A^T A \mathbf{x} = A^T \mathbf{b}$$

- Called the **normal equations**
- $A^T A$ is a *square* matrix

Solving linear least squares III

- We have reduced the linear least squares problem to solving a linear system:

$$A^T A \mathbf{x} = A^T \mathbf{b}$$

- Called the **normal equations**
- $A^T A$ is a *square* matrix
- This is solvable if A is **full rank** – columns span the space

Solving linear least squares III

- We have reduced the linear least squares problem to solving a linear system:

$$A^T A \mathbf{x} = A^T \mathbf{b}$$

- Called the **normal equations**
- $A^T A$ is a *square* matrix
- This is solvable if A is **full rank** – columns span the space
- Note that the condition $A^T(A\mathbf{x} - \mathbf{b}) = 0$ corresponds to $A\mathbf{x} - \mathbf{b}$ being perpendicular to each column of A !

Pseudoinverse

■ Mathematically: $\mathbf{x} = (\mathbf{A}^T \mathbf{A})^{-1} \mathbf{A}^T \mathbf{b}$

Pseudoinverse

- Mathematically: $\mathbf{x} = (\mathbf{A}^T \mathbf{A})^{-1} \mathbf{A}^T \mathbf{b}$
- Notation: $\mathbf{x} = \mathbf{A}^+ \mathbf{b}$
- $\mathbf{A}^+ := (\mathbf{A}^T \mathbf{A})^{-1} \mathbf{A}^T$ is called the **pseudo-inverse** of \mathbf{A}

Pseudoinverse

- Mathematically: $\mathbf{x} = (\mathbf{A}^T \mathbf{A})^{-1} \mathbf{A}^T \mathbf{b}$
- Notation: $\mathbf{x} = \mathbf{A}^+ \mathbf{b}$
- $\mathbf{A}^+ := (\mathbf{A}^T \mathbf{A})^{-1} \mathbf{A}^T$ is called the **pseudo-inverse** of \mathbf{A}
- $\mathbf{A}^T \mathbf{A}$ is invertible iff \mathbf{A} has full rank

Solving the normal equations

- It is possible to solve the normal equations using elimination, but to do so we need to construct $A^T A$

Solving the normal equations

- It is possible to solve the normal equations using elimination, but to do so we need to construct $A^T A$
- Slightly more efficient version of elimination for **symmetric** matrices: **Cholesky factorization**

Solving the normal equations

- It is possible to solve the normal equations using elimination, but to do so we need to construct $A^T A$
- Slightly more efficient version of elimination for **symmetric** matrices: **Cholesky factorization**
- Alternative: factorize $A = QR$

Solving the normal equations

- It is possible to solve the normal equations using elimination, but to do so we need to construct $A^T A$
- Slightly more efficient version of elimination for **symmetric** matrices: **Cholesky factorization**
- Alternative: factorize $A = QR$
- Need generalization of QR when A tall and narrow, $m \times n$

Solving the normal equations

- It is possible to solve the normal equations using elimination, but to do so we need to construct $A^T A$
- Slightly more efficient version of elimination for **symmetric** matrices: **Cholesky factorization**
- Alternative: factorize $A = QR$
- Need generalization of QR when A tall and narrow, $m \times n$
- **Thin** QR factorization: $A = QR$ where
 - Q is $(m \times n)$ with orthonormal columns
 - R is $(n \times n)$ upper triangular

Solving the normal equations

- It is possible to solve the normal equations using elimination, but to do so we need to construct $A^T A$
- Slightly more efficient version of elimination for **symmetric** matrices: **Cholesky factorization**
- Alternative: factorize $A = QR$
- Need generalization of QR when A tall and narrow, $m \times n$
- **Thin** QR factorization: $A = QR$ where
 - Q is $(m \times n)$ with orthonormal columns
 - R is $(n \times n)$ upper triangular
- If A is of full rank then R is non-singular

Solving normal equations II

- We want to solve $A^T A \mathbf{x} = A^T \mathbf{b}$
- Substitute $A = QR$

Solving normal equations II

- We want to solve $A^T A \mathbf{x} = A^T \mathbf{b}$
- Substitute $A = QR$
- So $R^T Q^T Q R \mathbf{x} = R^T Q^T \mathbf{b}$

Solving normal equations II

- We want to solve $A^T A \mathbf{x} = A^T \mathbf{b}$
- Substitute $A = QR$
- So $R^T Q^T Q R \mathbf{x} = R^T Q^T \mathbf{b}$
- So $R^T R \mathbf{x} = R^T Q^T \mathbf{b}$

Solving normal equations II

- We want to solve $A^T A \mathbf{x} = A^T \mathbf{b}$
- Substitute $A = QR$
- So $R^T Q^T Q R \mathbf{x} = R^T Q^T \mathbf{b}$
- So $R^T R \mathbf{x} = R^T Q^T \mathbf{b}$
- So $R \mathbf{x} = Q^T \mathbf{b}$

Solving normal equations II

- We want to solve $A^T A \mathbf{x} = A^T \mathbf{b}$
- Substitute $A = QR$
- So $R^T Q^T Q R \mathbf{x} = R^T Q^T \mathbf{b}$
- So $R^T R \mathbf{x} = R^T Q^T \mathbf{b}$
- So $R \mathbf{x} = Q^T \mathbf{b}$
- So $\mathbf{x} = R^{-1} Q^T \mathbf{b}$

Solving normal equations II

- We want to solve $A^T A \mathbf{x} = A^T \mathbf{b}$
- Substitute $A = QR$
- So $R^T Q^T Q R \mathbf{x} = R^T Q^T \mathbf{b}$
- So $R^T R \mathbf{x} = R^T Q^T \mathbf{b}$
- So $R \mathbf{x} = Q^T \mathbf{b}$
- So $\mathbf{x} = R^{-1} Q^T \mathbf{b}$
- Solve $R \mathbf{x} = Q^T \mathbf{b}$ by backsubstitution

Backslash for solving least squares

- Backslash in Julia is overloaded to give this least-squares solution when A is an $(m \times n)$ matrix (
- I.e. $A \setminus b$ gives $A^+ \mathbf{b} = (A^T A)^{-1} A^T \mathbf{b}$

Backslash for solving least squares

- Backslash in Julia is overloaded to give this least-squares solution when A is an $(m \times n)$ matrix (
- I.e. $A \setminus b$ gives $A^+ \mathbf{b} = (A^T A)^{-1} A^T \mathbf{b}$
- It is not necessarily clear that overloading \setminus in this way is a good idea!

Backslash for solving least squares

- Backslash in Julia is overloaded to give this least-squares solution when A is an $(m \times n)$ matrix (
- I.e. $A \setminus b$ gives $A^+ \mathbf{b} = (A^T A)^{-1} A^T \mathbf{b}$
- It is not necessarily clear that overloading \setminus in this way is a good idea!
- E.g. For a simple linear fit, use \setminus with above matrix

The Singular-Value Decomposition (SVD)

Action of a matrix

- Consider an arbitrary $(m \times n)$ matrix A

Action of a matrix

- Consider an arbitrary $(m \times n)$ matrix A
- Let's study its action on the unit sphere \mathbb{S}_n in \mathbb{R}^n
- $\mathbb{S}_n := \{\mathbf{x} \in \mathbb{R}^n : \|\mathbf{x}\|_2 = 1\}$

Action of a matrix

- Consider an arbitrary $(m \times n)$ matrix A
- Let's study its action on the unit sphere \mathbb{S}_n in \mathbb{R}^n
- $\mathbb{S}_n := \{\mathbf{x} \in \mathbb{R}^n : \|\mathbf{x}\|_2 = 1\}$
- A visualisation suggests that $A(\mathbb{S}_n)$ is an **ellipsoid** (hyper-ellipse)
- I.e. a sphere that is stretched and then rotated

Action of a matrix

- Consider an arbitrary $(m \times n)$ matrix A
- Let's study its action on the unit sphere \mathbb{S}_n in \mathbb{R}^n
- $\mathbb{S}_n := \{\mathbf{x} \in \mathbb{R}^n : \|\mathbf{x}\|_2 = 1\}$
- A visualisation suggests that $A(\mathbb{S}_n)$ is an **ellipsoid** (hyper-ellipse)
- I.e. a sphere that is stretched and then rotated
- Key quantities:
 - Lengths σ_i of semi-axes – stretches
 - Directions \mathbf{u}_i of stretches

Direction of maximum stretch

- Which is the direction with *maximal* stretch?
- This is a special direction associated to the matrix

Direction of maximum stretch

- Which is the direction with *maximal* stretch?
- This is a special direction associated to the matrix
- We need to solve

$$\operatorname{argmax}_{\mathbf{x}} \frac{\|\mathbf{Ax}\|}{\|\mathbf{x}\|}$$

Direction of maximum stretch

- Which is the direction with *maximal* stretch?
- This is a special direction associated to the matrix
- We need to solve

$$\operatorname{argmax}_{\mathbf{x}} \frac{\|\mathbf{Ax}\|}{\|\mathbf{x}\|}$$

- Another optimisation problem
- Also solvable using linear algebra

Action of a matrix II

- Let's think about the image of a sphere under a matrix

Action of a matrix II

- Let's think about the image of a sphere under a matrix
- Suppose $\mathbf{x} \in \mathbb{S}_n$; define $\mathbf{y} := A\mathbf{x}$ to be its image

Action of a matrix II

- Let's think about the image of a sphere under a matrix
- Suppose $\mathbf{x} \in \mathbb{S}_n$; define $\mathbf{y} := A\mathbf{x}$ to be its image
- Suppose that A is square and invertible for now

Action of a matrix II

- Let's think about the image of a sphere under a matrix
- Suppose $\mathbf{x} \in \mathbb{S}_n$; define $\mathbf{y} := A\mathbf{x}$ to be its image
- Suppose that A is square and invertible for now
- Then $\|A^{-1}\mathbf{y}\|^2 = 1$
- So $\mathbf{y}^T(A^{-1})^T A^{-1}\mathbf{y} = 1$

Action of a matrix II

- Let's think about the image of a sphere under a matrix
- Suppose $\mathbf{x} \in \mathbb{S}_n$; define $\mathbf{y} := A\mathbf{x}$ to be its image
- Suppose that A is square and invertible for now
- Then $\|A^{-1}\mathbf{y}\|^2 = 1$
- So $\mathbf{y}^T(A^{-1})^T A^{-1} \mathbf{y} = 1$
- Hence $\mathbf{y}^T S \mathbf{y} = 1$
- Where $S := (A^{-1})^T (A^{-1})$ is symmetric

Spectral theorem and eigen-decomposition

- The spectral theorem for symmetric matrices tells us that there is a basis of orthogonal eigenvectors \mathbf{v}_i for S
- Hence we have $SQ = QL$, where

$$Q := (\mathbf{v}_1 | \mathbf{v}_2 | \cdots | \mathbf{v}_n)$$

$$L := \begin{bmatrix} \lambda_1 & & & \\ & \lambda_2 & & \\ & & \ddots & \\ & & & \lambda_m \end{bmatrix}$$

Spectral theorem and eigen-decomposition

- The spectral theorem for symmetric matrices tells us that there is a basis of orthogonal eigenvectors \mathbf{v}_i for S
- Hence we have $SQ = QL$, where

$$Q := (\mathbf{v}_1 | \mathbf{v}_2 | \cdots | \mathbf{v}_n)$$

$$L := \begin{bmatrix} \lambda_1 & & & \\ & \lambda_2 & & \\ & & \ddots & \\ & & & \lambda_m \end{bmatrix}$$

- So $S = QLQ^T$

Action of a matrix III

- So $\mathbf{y}^T \mathbf{Q} \mathbf{L} \mathbf{Q}^T \mathbf{y} = 1$

Action of a matrix III

- So $\mathbf{y}^T \mathbf{Q} \mathbf{L} \mathbf{Q}^T \mathbf{y} = 1$
- Hence $\mathbf{z}^T \mathbf{L} \mathbf{z} = 1$
where $\mathbf{z} := \mathbf{Q}^T \mathbf{y}$

Action of a matrix III

- So $\mathbf{y}^T \mathbf{Q} \mathbf{L} \mathbf{Q}^T \mathbf{y} = 1$
- Hence $\mathbf{z}^T \mathbf{L} \mathbf{z} = 1$
where $\mathbf{z} := \mathbf{Q}^T \mathbf{y}$
- So $\lambda_1 z_1^2 + \cdots + \lambda_n z_n^2 = 1$

Action of a matrix III

- So $\mathbf{y}^T \mathbf{Q} \mathbf{L} \mathbf{Q}^T \mathbf{y} = 1$
- Hence $\mathbf{z}^T \mathbf{L} \mathbf{z} = 1$
where $\mathbf{z} := \mathbf{Q}^T \mathbf{y}$
- So $\lambda_1 z_1^2 + \cdots + \lambda_n z_n^2 = 1$
- Can show $(\mathbf{A}^T \mathbf{A})^{-1}$ has eigenvalues $\lambda_i > 0$ (exercise)

Action of a matrix III

- So $\mathbf{y}^T \mathbf{Q} \mathbf{L} \mathbf{Q}^T \mathbf{y} = 1$
- Hence $\mathbf{z}^T \mathbf{L} \mathbf{z} = 1$
 where $\mathbf{z} := \mathbf{Q}^T \mathbf{y}$
- So $\lambda_1 z_1^2 + \dots + \lambda_n z_n^2 = 1$
- Can show $(\mathbf{A}^T \mathbf{A})^{-1}$ has eigenvalues $\lambda_i > 0$ (exercise)
- Hence \mathbf{z} lies on an **ellipse**, with semi-axis lengths
 $\sigma_i := \sqrt{\lambda_i}$

Action of a matrix III

- So $\mathbf{y}^T \mathbf{Q} \mathbf{L} \mathbf{Q}^T \mathbf{y} = 1$
- Hence $\mathbf{z}^T \mathbf{L} \mathbf{z} = 1$
where $\mathbf{z} := \mathbf{Q}^T \mathbf{y}$
- So $\lambda_1 z_1^2 + \dots + \lambda_n z_n^2 = 1$
- Can show $(\mathbf{A}^T \mathbf{A})^{-1}$ has eigenvalues $\lambda_i > 0$ (exercise)
- Hence \mathbf{z} lies on an **ellipse**, with semi-axis lengths
 $\sigma_i := \sqrt{\lambda_i}$
- Thus \mathbf{y} lies on a rotated ellipse !

Singular values and singular vectors

- σ_i are **singular values** of A – this is a terrible name!

Singular values and singular vectors

- σ_i are **singular values** of A – this is a terrible name!
- The \mathbf{u}_i , the columns of Q , are the **left singular vectors**
- I.e. the directions in which the ellipse is stretched

Singular values and singular vectors

- σ_i are **singular values** of A – this is a terrible name!
- The \mathbf{u}_i , the columns of Q , are the **left singular vectors**
- I.e. the directions in which the ellipse is stretched
- \mathbf{v}_i such that $A\mathbf{v}_i = \sigma_i\mathbf{u}_i$ are the **right singular vectors**

Singular-value decomposition (SVD)

- Any $(m \times n)$ matrix A has an SVD:

$$A = U\Sigma V^T$$

- the columns of U are \mathbf{u}_i
- the columns of V are \mathbf{v}_i
- Σ is diagonal matrix with σ_i on diagonal

Singular-value decomposition (SVD)

- Any $(m \times n)$ matrix A has an SVD:

$$A = U\Sigma V^T$$

- the columns of U are \mathbf{u}_i
 - the columns of V are \mathbf{v}_i
 - Σ is diagonal matrix with σ_i on diagonal
-
- Note: only square, diagonalisable matrices have eigen-decompositions
 - But *any* matrix has an SVD

Singular-value decomposition (SVD)

- Any $(m \times n)$ matrix A has an SVD:

$$A = U\Sigma V^T$$

- the columns of U are \mathbf{u}_i
 - the columns of V are \mathbf{v}_i
 - Σ is diagonal matrix with σ_i on diagonal
-
- Note: only square, diagonalisable matrices have eigen-decompositions
 - But *any* matrix has an SVD
 - The SVD is often more useful than an eigen-decomposition

Calculating the SVD

- We can calculate (most of) the SVD

Calculating the SVD

- We can calculate (most of) the SVD
- Form $B := A^T A$

Calculating the SVD

- We can calculate (most of) the SVD
- Form $B := A^T A$
- Calculate eigenvalues and eigenvectors of B

Calculating the SVD

- We can calculate (most of) the SVD
- Form $B := A^T A$
- Calculate eigenvalues and eigenvectors of B
- **Exercise:** Relate these to the singular values and singular vectors of A

Rank of a matrix and best approximation

- Number of non-zero singular values = column rank = **rank**

Rank of a matrix and best approximation

- Number of non-zero singular values = column rank = **rank**
- The SVD gives the **best approximation** to A by a matrix of rank k :

$$A_k := \sum_{i=1}^k \sigma_i \mathbf{u}_i \mathbf{v}_i^T$$

Rank of a matrix and best approximation

- Number of non-zero singular values = column rank = **rank**
- The SVD gives the **best approximation** to A by a matrix of rank k :

$$A_k := \sum_{i=1}^k \sigma_i \mathbf{u}_i \mathbf{v}_i^T$$

- For a data matrix the SVD gives the best approximation by a k -dimensional subspace

Rank of a matrix and best approximation

- Number of non-zero singular values = column rank = **rank**
- The SVD gives the **best approximation** to A by a matrix of rank k :

$$A_k := \sum_{i=1}^k \sigma_i \mathbf{u}_i \mathbf{v}_i^T$$

- For a data matrix the SVD gives the best approximation by a k -dimensional subspace
- Use for **low-rank approximation** of data
- PCA (Principal Component Analysis) in statistics

Rank of a matrix and best approximation

- Number of non-zero singular values = column rank = **rank**
- The SVD gives the **best approximation** to A by a matrix of rank k :

$$A_k := \sum_{i=1}^k \sigma_i \mathbf{u}_i \mathbf{v}_i^T$$

- For a data matrix the SVD gives the best approximation by a k -dimensional subspace
- Use for **low-rank approximation** of data
- PCA (Principal Component Analysis) in statistics

Summary

- Linear least squares for overdetermined systems
- Solvable using linear algebra
- Solution given by normal equations – linear system
- Solve using QR decomposition

- The SVD gives a matrix as rotation + stretch + rotation
- Closely related to eigendecomposition of $A^T A$

- Applications to data compression