

32. Spectral methods for boundary-value problems

Summary of the previous class

- Choosing the number of interpolation points:
 - Check that the Chebyshev coefficients have decayed
- Using Chebyshev technology:
 - Differentiation
 - Clenshaw–Curtis integration
 - Roots

Goals for today

- Spectral methods
- Boundary-value problems for ODEs
- Eigenvalue problems
- Time evolution

(Pseudo)-spectral methods for boundary-value problems

- We want to solve **boundary-value problems**
- E.g. The Poisson equation for the electric potential due to a distribution of electric charge

(Pseudo)-spectral methods for boundary-value problems

- We want to solve **boundary-value problems**
- E.g. The Poisson equation for the electric potential due to a distribution of electric charge
- Example:

$$u_{xx} = f(x) \quad \text{for } -1 < x < 1$$

with **boundary conditions** $u(-1) = u(+1) = 0$

(Pseudo)-spectral methods for boundary-value problems

- We want to solve **boundary-value problems**
- E.g. The Poisson equation for the electric potential due to a distribution of electric charge
- Example:

$$u_{xx} = f(x) \quad \text{for } -1 < x < 1$$

with **boundary conditions** $u(-1) = u(+1) = 0$

- Notation: derivative $u_x := u' := \frac{du}{dx}$
- $u_{xx} = 2\text{nd derivative } \frac{d^2u}{dx^2}$

Boundary-value problems

- There is no time in the problem, unlike for initial-value problems

Boundary-value problems

- There is no time in the problem, unlike for initial-value problems
- Everything must be satisfied *simultaneously*

Boundary-value problems

- There is no time in the problem, unlike for initial-value problems
- Everything must be satisfied *simultaneously*
- Information from *both* boundaries influences the value at all intermediate points

Boundary-value problems

- There is no time in the problem, unlike for initial-value problems
- Everything must be satisfied *simultaneously*
- Information from *both* boundaries influences the value at all intermediate points
- There are stepping methods (**shooting**) but they do not work well

Boundary-value problems

- There is no time in the problem, unlike for initial-value problems
- Everything must be satisfied *simultaneously*
- Information from *both* boundaries influences the value at all intermediate points
- There are stepping methods (**shooting**) but they do not work well
- Instead we try to reduce this to a linear algebra problem

Boundary-value problems

- There is no time in the problem, unlike for initial-value problems
- Everything must be satisfied *simultaneously*
- Information from *both* boundaries influences the value at all intermediate points
- There are stepping methods (**shooting**) but they do not work well
- Instead we try to reduce this to a linear algebra problem
- We will follow Trefethen, *Spectral Methods in MATLAB* (Chaps. 6 + 7)

Collocation methods

- We will use a **collocation method** to solve the problem

Collocation methods

- We will use a **collocation method** to solve the problem
- This is sometimes called a “pseudo-spectral” method

Collocation methods

- We will use a **collocation method** to solve the problem
- This is sometimes called a “pseudo-spectral” method
- We want to solve for an unknown function u that satisfies the ODE and the boundary conditions

Collocation methods

- We will use a **collocation method** to solve the problem
- This is sometimes called a “pseudo-spectral” method
- We want to solve for an unknown function u that satisfies the ODE and the boundary conditions
- Idea:
 - Choose $N + 1$ nodes x_j and solve for $N + 1$ *unknown* values $u_j := u(x_j)$

Collocation methods

- We will use a **collocation method** to solve the problem
- This is sometimes called a “pseudo-spectral” method
- We want to solve for an unknown function u that satisfies the ODE and the boundary conditions
- Idea:
 - Choose $N + 1$ nodes x_j and solve for $N + 1$ *unknown* values $u_j := u(x_j)$
- Impose that the ODE is satisfied at those nodes, i.e. that the residual is exactly 0 there

Collocation methods II

- Let $p(x)$ be the unique degree- N polynomial interpolating the u_j at x_j

Collocation methods II

- Let $p(x)$ be the unique degree- N polynomial interpolating the u_j at x_j
- We *impose* that the differential equation is satisfied at non-boundary nodes, so

$$p''(t_j) = u_j$$

Collocation methods II

- Let $p(x)$ be the unique degree- N polynomial interpolating the u_j at x_j
- We *impose* that the differential equation is satisfied at non-boundary nodes, so

$$p''(t_j) = u_j$$

- So

$$\mathbf{M}\mathbf{u} = \mathbf{f}$$

where $\mathbf{M} := \mathbf{D}_N^2$

Collocation methods II

- Let $p(x)$ be the unique degree- N polynomial interpolating the u_j at x_j
- We *impose* that the differential equation is satisfied at non-boundary nodes, so

$$p''(t_j) = u_j$$

- So

$$\mathbf{M}\mathbf{u} = \mathbf{f}$$

where $\mathbf{M} := \mathbf{D}_N^2$

- \mathbf{D}_N is differentiation matrix

Boundary conditions

- We also need to impose **boundary conditions**

Boundary conditions

- We also need to impose **boundary conditions**
- e.g. Dirichlet:

$$u(-1) = u(1) = 0$$

Boundary conditions

- We also need to impose **boundary conditions**
- e.g. Dirichlet:

$$u(-1) = u(1) = 0$$

- There are various ways to implement this

Boundary conditions

- We also need to impose **boundary conditions**
- e.g. Dirichlet:

$$u(-1) = u(1) = 0$$

- There are various ways to implement this
- E.g. **boundary bordering**:
 set rows 1 and $N + 1$ of M to rows of the identity matrix
 to impose $u_0 = u_N = 0$

Chebyshev spectral methods

- The formulation in terms of differentiation matrices can be used with finite differences . . .
- But using Chebyshev methods instead gives the usual advantage: **spectral convergence**

Summary

- Chebyshev spectral methods
- Collocation: Assume ODE is satisfied at interior points