

22. Chebyshev methods II

Summary of the previous class

- Convergence of the trapezoid method for periodic integrands
- Mapping Fourier to a non-periodic setting
- Chebyshev polynomials
- Chebyshev interpolation
- Discrete Cosine transform

Goals for today

- Choosing the number of interpolation points
- Operations in Chebyshev representation:
- Derivatives
- Integrals
- Roots

Chebyshev interpolation

- Recall: For a given $f : [-1, +1] \rightarrow \mathbb{R}$

Chebyshev interpolation

- Recall: For a given $f : [-1, +1] \rightarrow \mathbb{R}$
- We approximate $f \simeq f_n := \sum_{k=0}^n \alpha_k T_k$
- **Chebyshev polynomials** $T_k(x) := \cos(k \arccos(x))$
 - degree n polynomial

Chebyshev interpolation

- Recall: For a given $f : [-1, +1] \rightarrow \mathbb{R}$
- We approximate $f \simeq f_n := \sum_{k=0}^n \alpha_k T_k$
- **Chebyshev polynomials** $T_k(x) := \cos(k \arccos(x))$
– degree n polynomial
- Interpolate in **Chebyshev points** $x_j := \cos\left(\frac{\pi j}{N}\right)$
- $f_j := f(x_j)$ at $(n + 1)$ points x_j with $j = 0, \dots, n$

Chebyshev interpolation

- Recall: For a given $f : [-1, +1] \rightarrow \mathbb{R}$
- We approximate $f \simeq f_n := \sum_{k=0}^n \alpha_k T_k$
- **Chebyshev polynomials** $T_k(x) := \cos(k \arccos(x))$
– degree n polynomial
- Interpolate in **Chebyshev points** $x_j := \cos\left(\frac{\pi j}{N}\right)$
- $f_j := f(x_j)$ at $(n + 1)$ points x_j with $j = 0, \dots, n$
- Discrete Cosine Transformation (DCT):

$$\sum_k \alpha_k \cos\left(\frac{j k \pi}{n}\right) = f_j$$

Choosing the number of interpolation points

- Fundamental idea:

Represent / approximate function f by Chebyshev interpolant in Chebyshev points

Choosing the number of interpolation points

- Fundamental idea:

Represent / approximate function f by Chebyshev interpolant in Chebyshev points

- But *how many* points n should we choose?

Choosing the number of interpolation points

- Fundamental idea:

Represent / approximate function f by Chebyshev interpolant in Chebyshev points

- But *how many* points n should we choose?
- Enough that Chebyshev coefficients have decayed to ϵ_{mach}

Choosing number of interpolation points II

- Start with $n + 1$, e.g. $n = 8$

Choosing number of interpolation points II

- Start with $n + 1$, e.g. $n = 8$
- Interpolate f to get $f_j^{(n)}$

Choosing number of interpolation points II

- Start with $n + 1$, e.g. $n = 8$
- Interpolate f to get $f_j^{(n)}$
- Calculate α_k using the Fast Cosine Transform

Choosing number of interpolation points II

- Start with $n + 1$, e.g. $n = 8$
- Interpolate f to get $f_j^{(n)}$
- Calculate α_k using the Fast Cosine Transform
- Check if they are small, e.g. last two are $< 10^{-13}$

Choosing number of interpolation points II

- Start with $n + 1$, e.g. $n = 8$
- Interpolate f to get $f_j^{(n)}$
- Calculate α_k using the Fast Cosine Transform
- Check if they are small, e.g. last two are $< 10^{-13}$
- If not, **double** n and try again

Choosing number of interpolation points II

- Start with $n + 1$, e.g. $n = 8$
- Interpolate f to get $f_j^{(n)}$
- Calculate α_k using the Fast Cosine Transform
- Check if they are small, e.g. last two are $< 10^{-13}$

- If not, **double** n and try again
- Can reuse: $f_{2j}^{(2n)} = f_j^{(n)}$

Differentiation

- Suppose we have $f = \sum_{k=0}^n \alpha_k T_k$

Differentiation

- Suppose we have $f = \sum_{k=0}^n \alpha_k T_k$
- How can we calculate the derivative f' ?

Collaboration I

Differentiating a Chebyshev expansion

Suppose we have $f = \sum_{k=0}^n \alpha_k T_k$.

- 1 How could we calculate the derivative of the expansion directly?
- 2 What would we need to do so?
- 3 How could we calculate the derivative indirectly, without differentiating this expansion?

Differentiation II

- Suppose we have $f = \sum_{k=0}^n \alpha_k T_k$.

Differentiation II

- Suppose we have $f = \sum_{k=0}^n \alpha_k T_k$.
- The derivative will also be a sum of T_k :

$$f' = \sum_{k=1}^{n-1} \alpha_k T'_k$$

Differentiation II

- Suppose we have $f = \sum_{k=0}^n \alpha_k T_k$.
- The derivative will also be a sum of T_k :

$$f' = \sum_{k=1}^{n-1} \alpha_k T'_k$$

- This will be polynomial of degree $N - 1$

Differentiation III

- Let p be the unique polynomial that interpolates the f_j
- Set $w_j := p'(t_j)$

Differentiation III

- Let p be the unique polynomial that interpolates the f_j
- Set $w_j := p'(t_j)$
- Calculate using Lagrange interpolation since have explicit formulae

Differentiation III

- Let p be the unique polynomial that interpolates the f_j
- Set $w_j := p'(t_j)$
- Calculate using Lagrange interpolation since have explicit formulae
- Then

$$\mathbf{w} = \mathbf{D}_N \mathbf{f}$$

Differentiation III

- Let p be the unique polynomial that interpolates the f_j
- Set $w_j := p'(t_j)$
- Calculate using Lagrange interpolation since have explicit formulae
- Then

$$\mathbf{w} = D_N \mathbf{f}$$

- Where D_N is $(N + 1) \times (N + 1)$ **Chebyshev differentiation matrix**

Differentiation III

- Let p be the unique polynomial that interpolates the f_j
- Set $w_j := p'(t_j)$
- Calculate using Lagrange interpolation since have explicit formulae
- Then

$$\mathbf{w} = D_N \mathbf{f}$$

- Where D_N is $(N + 1) \times (N + 1)$ **Chebyshev differentiation matrix**
- Chapter 6 of Trefethen, *Spectral Methods in MATLAB* has explicit formulae for D_N

Differentiation IV

- The matrices D_N are *dense* matrices

Differentiation IV

- The matrices D_N are *dense* matrices
- Modern approach: Olver & Townsend 2014, “A fast and well-conditioned spectral method”

Differentiation IV

- The matrices D_N are *dense* matrices
- Modern approach: Olver & Townsend 2014, “A fast and well-conditioned spectral method”
- The derivative T'_k is an **ultraspherical polynomial**
- We get a **banded** matrix if we use the correct bases

Differentiation IV

- The matrices D_N are *dense* matrices
- Modern approach: Olver & Townsend 2014, “A fast and well-conditioned spectral method”
- The derivative T'_k is an **ultraspherical polynomial**
- We get a **banded** matrix if we use the correct bases
- “Differentiating scales the coefficients and changes the basis”

Numerical integration

- Given $f : [-1, +1] \rightarrow \mathbb{R}$
- How can we calculate $\int f = \int_{-1}^{+1} f(x) dx$?

Numerical integration

- Given $f : [-1, +1] \rightarrow \mathbb{R}$
- How can we calculate $\int f = \int_{-1}^{+1} f(x) dx$?
- Recall: trapezoid rule (equally-spaced points):
 $\mathcal{O}(n^{-2})$ convergence

Numerical integration

- Given $f : [-1, +1] \rightarrow \mathbb{R}$
- How can we calculate $\int f = \int_{-1}^{+1} f(x) dx$?
- Recall: trapezoid rule (equally-spaced points):
 $\mathcal{O}(n^{-2})$ convergence
- Can we do better?

Numerical integration

- Given $f : [-1, +1] \rightarrow \mathbb{R}$
- How can we calculate $\int f = \int_{-1}^{+1} f(x) dx$?
- Recall: trapezoid rule (equally-spaced points):
 $\mathcal{O}(n^{-2})$ convergence
- Can we do better?
- Clenshaw & Curtis (1960)

Collaboration II

Integration

Suppose we have $f = \sum_{k=0}^n \alpha_k T_k$.

- 1 How can we integrate f over $[-1, +1]$?
- 2 How accurate should the result be?

Clenshaw–Curtis integration

- We now have powerful **Chebyshev technology** at our disposal

Clenshaw–Curtis integration

- We now have powerful **Chebyshev technology** at our disposal
- Sample f at $n + 1$ Chebyshev points, giving f_j

Clenshaw–Curtis integration

- We now have powerful **Chebyshev technology** at our disposal
- Sample f at $n + 1$ Chebyshev points, giving f_j
- Find α_k in Chebyshev expansion

$$f \simeq f_N := \sum_{k=0}^n \alpha_k T_k$$
 via Fast Cosine Transform

Clenshaw–Curtis integration

- We now have powerful **Chebyshev technology** at our disposal
- Sample f at $n + 1$ Chebyshev points, giving f_j
- Find α_k in Chebyshev expansion

$$f \simeq f_N := \sum_{k=0}^n \alpha_k T_k$$
 via Fast Cosine Transform
- Double n until last few α_k are of order ϵ_{mach}

Clenshaw–Curtis integration II

- Now calculate $\int f$ as $\simeq \int f_n$:

$$\int f \simeq \sum_{k=0}^N \alpha_k \int T_k$$

Clenshaw–Curtis integration II

- Now calculate $\int f$ as $\simeq \int f_n$:

$$\int f \simeq \sum_{k=0}^N \alpha_k \int T_k$$

- Error $|\int f - \int f_N|$ decays as fast as $\|f - f_N\|$ does

Clenshaw–Curtis integration II

- Now calculate $\int f$ as $\simeq \int f_n$:

$$\int f \simeq \sum_{k=0}^N \alpha_k \int T_k$$

- Error $|\int f - \int f_N|$ decays as fast as $\|f - f_N\|$ does
- E.g. exponential (spectral) convergence if f is analytic

Clenshaw–Curtis integration III

- Need to calculate

$$\int f_N = \sum_{k=0}^N \alpha_k \int T_k$$

Clenshaw–Curtis integration III

- Need to calculate

$$\int f_N = \sum_{k=0}^N \alpha_k \int T_k$$

- Change variable in integral: $x = \cos(\theta)$

$$I_k = \int_{-1}^{+1} T_k(x) dx = \int_{-\pi}^{\pi} \cos(k\theta) \sin(\theta) d\theta$$

Clenshaw–Curtis integration III

- Need to calculate

$$\int f_N = \sum_{k=0}^N \alpha_k \int T_k$$

- Change variable in integral: $x = \cos(\theta)$

$$I_k = \int_{-1}^{+1} T_k(x) dx = \int_{-\pi}^{\pi} \cos(k\theta) \sin(\theta) d\theta$$

- Find $I_k = \frac{2}{1-k^2}$ if k is even, and 0 if k is odd

Clenshaw–Curtis integration IV

■ So we have

$$\int f = \sum_{k=0}^N \alpha_k I_k$$

Clenshaw–Curtis integration IV

- So we have

$$\int f = \sum_{k=0}^N \alpha_k I_k$$

- The integral becomes a dot product!

Clenshaw–Curtis integration IV

- So we have

$$\int f = \sum_{k=0}^N \alpha_k I_k$$

- The integral becomes a dot product!
- This is a version of the **Riesz representation theorem** for linear functional

Recurrence relation

- It turns out that there is a 3-term **recurrence relation** relating T_k to T_{k-1} and T_{k-2} :

$$T_k(x) = 2xT_{k-1}(x) - T_{k-2}(x)$$

Recurrence relation

- It turns out that there is a 3-term **recurrence relation** relating T_k to T_{k-1} and T_{k-2} :

$$T_k(x) = 2xT_{k-1}(x) - T_{k-2}(x)$$

- This allows us to calculate $T_k(x)$ explicitly, using $T_0 = 1$ and $T_1(x) = x$

Recurrence relation

- It turns out that there is a 3-term **recurrence relation** relating T_k to T_{k-1} and T_{k-2} :

$$T_k(x) = 2xT_{k-1}(x) - T_{k-2}(x)$$

- This allows us to calculate $T_k(x)$ explicitly, using $T_0 = 1$ and $T_1(x) = x$
- Where does this recurrence relation come from?

Recurrence relation II

- Consider $xT_k(x)$

Recurrence relation II

- Consider $xT_k(x)$
- T_k is a degree- k polynomial

Recurrence relation II

- Consider $xT_k(x)$
- T_k is a degree- k polynomial
- So $xT_k(x)$ is a degree- $(k + 1)$ polynomial

Recurrence relation II

- Consider $xT_k(x)$
- T_k is a degree- k polynomial
- So $xT_k(x)$ is a degree- $(k + 1)$ polynomial
- So we can write it as

$$xT_k(x) = \sum_{j=0}^{k+1} \alpha_{k,j} T_j(x)$$

Recurrence relation II

- Consider $xT_k(x)$
- T_k is a degree- k polynomial
- So $xT_k(x)$ is a degree- $(k + 1)$ polynomial
- So we can write it as

$$xT_k(x) = \sum_{j=0}^{k+1} \alpha_{k,j} T_j(x)$$

- Where $\alpha_{k,j} \propto (xT_k, T_j)$, i.e. equal up to normalization

Recurrence relation II

- We have

$$(xT_k, T_j) = \int_{-1}^1 xT_k(x)T_j(x)w(x)dx$$

- Change variables using $x = \cos(\theta)$:

$$(xT_k, T_j) = \int_0^{2\pi} c_1 c_k c_j d\theta$$

where $c_j(\theta) := \cos(j\theta)$

Recurrence relation II

- We have

$$(xT_k, T_j) = \int_{-1}^1 xT_k(x)T_j(x)w(x)dx$$

- Change variables using $x = \cos(\theta)$:

$$(xT_k, T_j) = \int_0^{2\pi} c_1 c_k c_j d\theta$$

where $c_j(\theta) := \cos(j\theta)$

- Use trigonometric relation

$$\cos(A)\cos(B) = \frac{1}{2}[\cos(A+B) + \cos(A-B)]$$

Recurrence relation III

■ We have

$$(xT_k, T_j) = \frac{1}{2} \int_0^{2\pi} c_1 [c_{k+j} + c_{k-j}]$$

Recurrence relation III

- We have

$$(xT_k, T_j) = \frac{1}{2} \int_0^{2\pi} c_1 [c_{k+j} + c_{k-j}]$$

- But $\int c_l c_m = 0$ if $l \neq m$

Recurrence relation III

- We have

$$(xT_k, T_j) = \frac{1}{2} \int_0^{2\pi} c_1 [c_{k+j} + c_{k-j}]$$

- But $\int c_l c_m = 0$ if $l \neq m$
- So (xT_k, T_j) is 0 unless $j = k + 1$ or $j = k - 1$

Recurrence relation III

- We have

$$(xT_k, T_j) = \frac{1}{2} \int_0^{2\pi} c_1 [c_{k+j} + c_{k-j}]$$

- But $\int c_l c_m = 0$ if $l \neq m$
- So (xT_k, T_j) is 0 unless $j = k + 1$ or $j = k - 1$
- Hence $xT_k = \alpha T_{k+1} + \beta T_{k-1}$
and we can calculate the constants α and β

Recurrence relation III

- We have

$$(xT_k, T_j) = \frac{1}{2} \int_0^{2\pi} c_1 [c_{k+j} + c_{k-j}]$$

- But $\int c_l c_m = 0$ if $l \neq m$
- So (xT_k, T_j) is 0 unless $j = k + 1$ or $j = k - 1$
- Hence $xT_k = \alpha T_{k+1} + \beta T_{k-1}$
and we can calculate the constants α and β

- In fact, *any* set of orthogonal polynomials have similar 3-term recurrence

Aliasing

- **Aliasing** is the following phenomenon

Aliasing

- **Aliasing** is the following phenomenon
- Consider sampling a Chebyshev polynomial T_k at Chebyshev points t_j
- This can give the *same function values* as sampling a *different* T_ℓ

Aliasing

- **Aliasing** is the following phenomenon
- Consider sampling a Chebyshev polynomial T_k at Chebyshev points t_j
- This can give the *same function values* as sampling a *different* T_ℓ
- On an $(n + 1)$ -point Chebyshev grid, the following have same values:
- $T_m, T_{2n-m}, T_{2n+m}, \dots$

Aliasing

- **Aliasing** is the following phenomenon
- Consider sampling a Chebyshev polynomial T_k at Chebyshev points t_j
- This can give the *same function values* as sampling a *different* T_ℓ
- On an $(n + 1)$ -point Chebyshev grid, the following have same values:
- $T_m, T_{2n-m}, T_{2n+m}, \dots$
- This gives a relationship between the coefficients of *truncation* of f and *interpolation* of f
- See Trefethen, *Approx. Theory*, Chap. 4

Root finding

- How can we find all **roots** of a complicated function $f : [-1, +1] \rightarrow \mathbb{R}$?

Root finding

- How can we find all **roots** of a complicated function $f : [-1, +1] \rightarrow \mathbb{R}$?
- Approximate f by a Chebyshev expansion p and find the roots of p

Root finding

- How can we find all **roots** of a complicated function $f : [-1, +1] \rightarrow \mathbb{R}$?
- Approximate f by a Chebyshev expansion p and find the roots of p
- Find a matrix C whose *eigenvalues* are the roots of p

Root finding

- How can we find all **roots** of a complicated function $f : [-1, +1] \rightarrow \mathbb{R}$?
- Approximate f by a Chebyshev expansion p and find the roots of p
- Find a matrix C whose *eigenvalues* are the roots of p
- In other words the characteristic polynomial of the matrix...

Root finding

- How can we find all **roots** of a complicated function $f : [-1, +1] \rightarrow \mathbb{R}$?
- Approximate f by a Chebyshev expansion p and find the roots of p
- Find a matrix C whose *eigenvalues* are the roots of p
- In other words the characteristic polynomial of the matrix...
- ... is p itself!

Root finding

- How can we find all **roots** of a complicated function $f : [-1, +1] \rightarrow \mathbb{R}$?
- Approximate f by a Chebyshev expansion p and find the roots of p
- Find a matrix C whose *eigenvalues* are the roots of p
- In other words the characteristic polynomial of the matrix...
- ... is p itself!
- For a polynomial written in the monomial basis we get the **companion** matrix

Root finding

- How can we find all **roots** of a complicated function $f : [-1, +1] \rightarrow \mathbb{R}$?
- Approximate f by a Chebyshev expansion p and find the roots of p
- Find a matrix C whose *eigenvalues* are the roots of p
- In other words the characteristic polynomial of the matrix...
- ... is p itself!
- For a polynomial written in the monomial basis we get the **companion** matrix
- In the Chebyshev basis it is called the **colleague matrix**

Colleague matrix

- The **colleague matrix** of the polynomial

$$p(x) = \sum_{k=0}^n a_k T_k(x), \quad a_n \neq 0$$

is

$$C := \begin{pmatrix} 0 & 1 & & & \\ \frac{1}{2} & 0 & & & \\ & \frac{1}{2} & 0 & & \\ & & \ddots & \ddots & \ddots \\ & & & \frac{1}{2} & 0 \end{pmatrix} - \frac{1}{2a_n} \begin{pmatrix} & & & & \\ & & & & \\ & & & & \\ & & & & \\ a_0 & a_1 & a_2 & \cdots & a_{n-1} \end{pmatrix}$$

Summary

- Spectral convergence gives excellent approximation of function
- Fundamental mathematical operations become “easy” once we have a spectral approximation
- This is (mostly) maintained by operations like differentiation, integration
- 3-term recurrence relation for Chebyshev polynomials
- Clenshaw–Curtis integration
- Spectrally accurate for analytic functions

Further reading

- Boyd, Finding the Zeros of a Univariate Equation: Proxy Rootfinders, Chebyshev Interpolation, and the Companion Matrix, SIAM Review **55**(2), 375–396.
<https://epubs.siam.org/doi/pdf/10.1137/110838297>
- Boyd, **Chebyshev and Fourier Spectral Methods**
- Trefethen, **Spectral Methods in Matlab**
- Trefethen, **Approximation Theory and Approximation Practice**