

13. The Singular-Value Decomposition (SVD) and applications

Last time

- Least-squares problems
- Minimise $\|A \mathbf{x} - \mathbf{b}\|^2$
- Optimization
- Solution via normal equations and QR
- Action of a matrix

Goals for today

- Action of a matrix A
- Eigen-factorisation (eigen-decomposition) for symmetric, square matrix

Goals for today

- Action of a matrix A
- Eigen-factorisation (eigen-decomposition) for symmetric, square matrix
- Singular-Value Decomposition (SVD)
 - Factorisation $A = U\Sigma V^T$ for any $(m \times n)$ matrix
- Applications to understanding and compressing data

The Singular-Value Decomposition (SVD)

How does a matrix act?

- Consider an arbitrary $(m \times n)$ matrix A

How does a matrix act?

- Consider an arbitrary $(m \times n)$ matrix A
- What is the *geometrical* action of the matrix?

How does a matrix act?

- Consider an arbitrary $(m \times n)$ matrix A
- What is the *geometrical* action of the matrix?
- For 2×2 matrices we know that they can:
 - do nothing (identity)
 - reflect
 - shear
 - rotate

How does a matrix act?

- Consider an arbitrary $(m \times n)$ matrix A
- What is the *geometrical* action of the matrix?
- For 2×2 matrices we know that they can:
 - do nothing (identity)
 - reflect
 - shear
 - rotate
- In general they send **parallelograms** to **parallelograms**

How does a matrix act?

- Consider an arbitrary $(m \times n)$ matrix A
- What is the *geometrical* action of the matrix?
- For 2×2 matrices we know that they can:
 - do nothing (identity)
 - reflect
 - shear
 - rotate
- In general they send **parallelograms** to **parallelograms**
- What about in higher dimensions?

Direction of maximal stretching

- One approach to this question is an eigen-factorisation
- We look for special directions \mathbf{v} such that $A\mathbf{v} = \lambda\mathbf{v}$

Direction of maximal stretching

- One approach to this question is an eigen-factorisation
- We look for special directions \mathbf{v} such that $A\mathbf{v} = \lambda\mathbf{v}$
- But that works *only* for (some) square matrices

Direction of maximal stretching

- One approach to this question is an eigen-factorisation
- We look for special directions \mathbf{v} such that $A\mathbf{v} = \lambda\mathbf{v}$
- But that works *only* for (some) square matrices
- We want a general approach that applies to *any* matrix

Direction of maximal stretching

- One approach to this question is an eigen-factorisation
 - We look for special directions \mathbf{v} such that $A\mathbf{v} = \lambda\mathbf{v}$
 - But that works *only* for (some) square matrices
 - We want a general approach that applies to *any* matrix
-
- Idea: Look for different special directions: Those associated with different amounts of **stretching**

Direction of maximal stretching

- One approach to this question is an eigen-factorisation
 - We look for special directions \mathbf{v} such that $A\mathbf{v} = \lambda\mathbf{v}$
 - But that works *only* for (some) square matrices
 - We want a general approach that applies to *any* matrix
-
- Idea: Look for different special directions: Those associated with different amounts of **stretching**
 - E.g. Which is the direction \mathbf{x} with *maximal* **stretching**?

Direction of maximal stretching

- We need to solve

$$\operatorname{argmax}_{\mathbf{x}} \frac{\|\mathbf{A}\mathbf{x}\|}{\|\mathbf{x}\|}$$

Direction of maximal stretching

- We need to solve

$$\operatorname{argmax}_{\mathbf{x}} \frac{\|\mathbf{A}\mathbf{x}\|}{\|\mathbf{x}\|}$$

- Equivalently: $\operatorname{argmax}_{\|x\|=1} \|\mathbf{A}\mathbf{x}\|$

Direction of maximal stretching

- We need to solve

$$\operatorname{argmax}_{\mathbf{x}} \frac{\|A\mathbf{x}\|}{\|\mathbf{x}\|}$$

- Equivalently: $\operatorname{argmax}_{\|x\|=1} \|A\mathbf{x}\|$
- Another optimisation problem
- Also solvable using linear algebra

Action of a matrix

- Let's study its action on the unit sphere \mathbb{S}^{n-1} in \mathbb{R}^n
- $\mathbb{S}^{n-1} := \{\mathbf{x} \in \mathbb{R}^n : \|\mathbf{x}\|^2 = 1\}$

Action of a matrix

- Let's study its action on the unit sphere \mathbb{S}^{n-1} in \mathbb{R}^n
- $\mathbb{S}^{n-1} := \{\mathbf{x} \in \mathbb{R}^n : \|\mathbf{x}\|^2 = 1\}$
- The image is a set in \mathbb{R}^m

Action of a matrix

- Let's study its action on the unit sphere \mathbb{S}^{n-1} in \mathbb{R}^n
- $\mathbb{S}^{n-1} := \{\mathbf{x} \in \mathbb{R}^n : \|\mathbf{x}\|^2 = 1\}$
- The image is a set in \mathbb{R}^m
- A visualisation suggests that $A(\mathbb{S}^{n-1})$ is an **ellipsoid** (hyper-ellipse) – a stretched and rotated sphere

Action of a matrix

- Let's study its action on the unit sphere \mathbb{S}^{n-1} in \mathbb{R}^n
- $\mathbb{S}^{n-1} := \{\mathbf{x} \in \mathbb{R}^n : \|\mathbf{x}\|^2 = 1\}$
- The image is a set in \mathbb{R}^m
- A visualisation suggests that $A(\mathbb{S}^{n-1})$ is an **ellipsoid** (hyper-ellipse) – a stretched and rotated sphere
- Key quantities:
 - Lengths σ_i of semi-axes – stretches
 - Directions \mathbf{u}_i of stretches

Collaboration I

Understanding the action of a matrix

Assume for now that A is square and invertible.

- 1 Suppose that \mathbf{y} lies in the *image* of the sphere under A . What can you say about \mathbf{y} ?
- 2 Can you expand this equation to get something more useful?

Action of a matrix II

- Let's think about the image of \mathbb{S}^{n-1} under A

Action of a matrix II

- Let's think about the image of \mathbb{S}^{n-1} under A
- Suppose $\mathbf{x} \in \mathbb{S}^{n-1}$
- Define $\mathbf{y} := A\mathbf{x}$ to be its image

Action of a matrix II

- Let's think about the image of \mathbb{S}^{n-1} under A
- Suppose $\mathbf{x} \in \mathbb{S}^{n-1}$
- Define $\mathbf{y} := A\mathbf{x}$ to be its image
- Suppose that A is square and invertible for now

Action of a matrix II

- Let's think about the image of \mathbb{S}^{n-1} under A
 - Suppose $\mathbf{x} \in \mathbb{S}^{n-1}$
 - Define $\mathbf{y} := A\mathbf{x}$ to be its image
 - Suppose that A is square and invertible for now
-
- Since $\|x\|^2 = 1$ we have $\|A^{-1}\mathbf{y}\|^2 = 1$
 - So $\mathbf{y}^T(A^{-1})^T A^{-1}\mathbf{y} = 1$

Action of a matrix II

- Let's think about the image of \mathbb{S}^{n-1} under A
- Suppose $\mathbf{x} \in \mathbb{S}^{n-1}$
- Define $\mathbf{y} := A\mathbf{x}$ to be its image
- Suppose that A is square and invertible for now
- Since $\|x\|^2 = 1$ we have $\|A^{-1}\mathbf{y}\|^2 = 1$
- So $\mathbf{y}^T(A^{-1})^T A^{-1}\mathbf{y} = 1$
- Hence $\mathbf{y}^T S \mathbf{y} = 1$
- Where $S := (A^{-1})^T(A^{-1})$ is symmetric

Spectral theorem for symmetric matrices

- Recall the **spectral theorem** for symmetric matrices

Spectral theorem for symmetric matrices

- Recall the **spectral theorem** for symmetric matrices
- It tells us that S has a *basis* of orthogonal eigenvectors, \mathbf{q}_i , for S :

$$S\mathbf{q}_i = \lambda_i\mathbf{q}_i$$

Collaboration II

Eigen-factorisation

- 1 Use the spectral theorem to find an equation involving S and a certain matrix that you can construct.
- 2 Hence write down a matrix factorisation of S , i.e. $S = \dots$
- 3 Can you use that to say something about the image vector \mathbf{y} ?

Eigen-factorisation

- Let's build an orthogonal matrix Q whose columns are the orthonormal basis vectors \mathbf{q}_i :

Eigen-factorisation

- Let's build an orthogonal matrix Q whose columns are the orthonormal basis vectors \mathbf{q}_i :

$$Q := [\mathbf{q}_1 | \mathbf{q}_2 | \cdots | \mathbf{q}_n]$$

Eigen-factorisation

- Let's build an orthogonal matrix Q whose columns are the orthonormal basis vectors \mathbf{q}_i :

$$Q := [\mathbf{q}_1 | \mathbf{q}_2 | \cdots | \mathbf{q}_n]$$

- Since $S\mathbf{q}_i = \lambda_i\mathbf{q}_i$ we get $SQ = Q\Lambda$
- With

$$\Lambda := \begin{bmatrix} \lambda_1 & & & \\ & \lambda_2 & & \\ & & \ddots & \\ & & & \lambda_m \end{bmatrix}$$

Eigen-factorisation

- Let's build an orthogonal matrix Q whose columns are the orthonormal basis vectors \mathbf{q}_i :

$$Q := [\mathbf{q}_1 | \mathbf{q}_2 | \cdots | \mathbf{q}_n]$$

- Since $S\mathbf{q}_i = \lambda_i\mathbf{q}_i$ we get $SQ = Q\Lambda$
- With

$$\Lambda := \begin{bmatrix} \lambda_1 & & & \\ & \lambda_2 & & \\ & & \ddots & \\ & & & \lambda_m \end{bmatrix}$$

- Hence $S = Q\Lambda Q^T$ – the **eigen-factorisation**

Action of a matrix III

■ So $\mathbf{y}^T \mathbf{Q} \mathbf{\Lambda} \mathbf{Q}^T \mathbf{y} = 1$

Action of a matrix III

- So $\mathbf{y}^T \mathbf{Q} \mathbf{\Lambda} \mathbf{Q}^T \mathbf{y} = 1$
- Hence $\mathbf{z}^T \mathbf{\Lambda} \mathbf{z} = 1$
where $\mathbf{z} := \mathbf{Q}^T \mathbf{y}$

Action of a matrix III

- So $\mathbf{y}^T \mathbf{Q} \mathbf{\Lambda} \mathbf{Q}^T \mathbf{y} = 1$

- Hence $\mathbf{z}^T \mathbf{\Lambda} \mathbf{z} = 1$

where $\mathbf{z} := \mathbf{Q}^T \mathbf{y}$

- So $\lambda_1 z_1^2 + \cdots + \lambda_n z_n^2 = 1$

Action of a matrix III

- So $\mathbf{y}^T \mathbf{Q} \mathbf{\Lambda} \mathbf{Q}^T \mathbf{y} = 1$
- Hence $\mathbf{z}^T \mathbf{\Lambda} \mathbf{z} = 1$
where $\mathbf{z} := \mathbf{Q}^T \mathbf{y}$
- So $\lambda_1 z_1^2 + \cdots + \lambda_n z_n^2 = 1$
- Can show $(\mathbf{A}^T \mathbf{A})^{-1}$ has eigenvalues $\lambda_i > 0$ (exercise)

Action of a matrix III

- So $\mathbf{y}^T \mathbf{Q} \mathbf{\Lambda} \mathbf{Q}^T \mathbf{y} = 1$
- Hence $\mathbf{z}^T \mathbf{\Lambda} \mathbf{z} = 1$
 where $\mathbf{z} := \mathbf{Q}^T \mathbf{y}$
- So $\lambda_1 z_1^2 + \dots + \lambda_n z_n^2 = 1$
- Can show $(\mathbf{A}^T \mathbf{A})^{-1}$ has eigenvalues $\lambda_i > 0$ (exercise)
- Hence \mathbf{z} lies on an **ellipse**, with semi-axis lengths
 $\sigma_i := \sqrt{\lambda_i}$

Action of a matrix III

- So $\mathbf{y}^T \mathbf{Q} \mathbf{\Lambda} \mathbf{Q}^T \mathbf{y} = 1$
- Hence $\mathbf{z}^T \mathbf{\Lambda} \mathbf{z} = 1$
 where $\mathbf{z} := \mathbf{Q}^T \mathbf{y}$
- So $\lambda_1 z_1^2 + \dots + \lambda_n z_n^2 = 1$
- Can show $(\mathbf{A}^T \mathbf{A})^{-1}$ has eigenvalues $\lambda_i > 0$ (exercise)
- Hence \mathbf{z} lies on an **ellipse**, with semi-axis lengths
 $\sigma_i := \sqrt{\lambda_i}$
- Thus \mathbf{y} lies on a rotated ellipse !

Singular values and singular vectors

- σ_i are **singular values** of A – this is a terrible name!

Singular values and singular vectors

- σ_i are **singular values** of A – this is a terrible name!
- The \mathbf{u}_i , the columns of Q , are the **left singular vectors**
- I.e. the directions in which the ellipse is stretched

Singular values and singular vectors

- σ_i are **singular values** of A – this is a terrible name!
- The \mathbf{u}_i , the columns of Q , are the **left singular vectors**
- I.e. the directions in which the ellipse is stretched
- \mathbf{v}_i such that $A\mathbf{v}_i = \sigma_i\mathbf{u}_i$ are the **right singular vectors**

Singular values and singular vectors

- σ_i are **singular values** of A – this is a terrible name!
- The \mathbf{u}_i , the columns of Q , are the **left singular vectors**
- I.e. the directions in which the ellipse is stretched
- \mathbf{v}_i such that $A\mathbf{v}_i = \sigma_i\mathbf{u}_i$ are the **right singular vectors**
- We arrange the singular values in decreasing order:
$$\sigma_1 \geq \sigma_2 \geq \dots$$

Singular-value decomposition (SVD)

- Any $(m \times n)$ matrix A has an SVD:

$$A = U\Sigma V^T = \sum_i \sigma_i \mathbf{u}_i \mathbf{v}_i^T$$

- the columns of U are \mathbf{u}_i
- the columns of V are \mathbf{v}_i
- Σ is diagonal matrix with σ_i on diagonal

Singular-value decomposition (SVD)

- Any $(m \times n)$ matrix A has an SVD:

$$A = U\Sigma V^T = \sum_i \sigma_i \mathbf{u}_i \mathbf{v}_i^T$$

- the columns of U are \mathbf{u}_i
 - the columns of V are \mathbf{v}_i
 - Σ is diagonal matrix with σ_i on diagonal
-
- Note: only square, diagonalisable matrices have eigen-factorisations
 - But *any* matrix has an SVD

Singular-value decomposition (SVD)

- Any $(m \times n)$ matrix A has an SVD:

$$A = U\Sigma V^T = \sum_i \sigma_i \mathbf{u}_i \mathbf{v}_i^T$$

- the columns of U are \mathbf{u}_i
 - the columns of V are \mathbf{v}_i
 - Σ is diagonal matrix with σ_i on diagonal
- Note: only square, diagonalisable matrices have eigen-factorisations
- But *any* matrix has an SVD
- The SVD is often more useful than an eigen factorisation

Collaboration III

Calculating the SVD

- 1 How could you *calculate* the SVD, using what we have seen so far in the course?

Calculating the SVD

- We can *calculate* the SVD
- There are more efficient algorithms that are beyond the scope of the course

Calculating the SVD

- We can *calculate* the SVD
- There are more efficient algorithms that are beyond the scope of the course
- Form $B := A^T A$

Calculating the SVD

- We can *calculate* the SVD
- There are more efficient algorithms that are beyond the scope of the course
- Form $B := A^T A$
- Calculate the eigenvalues and eigenvectors of the symmetric matrix B

Calculating the SVD

- We can *calculate* the SVD
- There are more efficient algorithms that are beyond the scope of the course
- Form $B := A^T A$
- Calculate the eigenvalues and eigenvectors of the symmetric matrix B
- Do the same for $C := A A^T$

Calculating the SVD

- We can *calculate* the SVD
- There are more efficient algorithms that are beyond the scope of the course
- Form $B := A^T A$
- Calculate the eigenvalues and eigenvectors of the symmetric matrix B
- Do the same for $C := A A^T$
- **Exercise:** Relate these to the singular values and singular vectors of A

Rank of a matrix

- The SVD gives as information about the action of a matrix

Rank of a matrix

- The SVD gives as information about the action of a matrix
- E.g. the **rank** of a matrix is the dimension of the *image* of \mathbb{R}^n (i.e. the column rank)

Rank of a matrix

- The SVD gives as information about the action of a matrix
- E.g. the **rank** of a matrix is the dimension of the *image* of \mathbb{R}^n (i.e. the column rank)
- This is given by the *number of non-zero singular values* !

Rank of a matrix

- The SVD gives as information about the action of a matrix
- E.g. the **rank** of a matrix is the dimension of the *image* of \mathbb{R}^n (i.e. the column rank)
- This is given by the *number of non-zero singular values* !
- Usually floating-point round-off error will make the singular values non-zero
- So we need to look at the **numerical rank**
 - the number of “non-zero” singular values above a threshold

Best-fitting subspace

- Suppose that we want to find the best-fitting plane, or dimension- k hyperplane, to a cloud of data

Best-fitting subspace

- Suppose that we want to find the best-fitting plane, or dimension- k hyperplane, to a cloud of data
- It turns out that the SVD solves this problem *immediately*!

Best-fitting subspace

- Suppose that we want to find the best-fitting plane, or dimension- k hyperplane, to a cloud of data
- It turns out that the SVD solves this problem *immediately*!
- We first **de-mean** the data (subtract the mean from each component)
- Then the SVD gives the **best approximation** to a matrix A by a matrix of rank k as

$$A_k := \sum_{i=1}^k \sigma_i \mathbf{u}_i \mathbf{v}_i^T$$

Best-fitting subspace

- Suppose that we want to find the best-fitting plane, or dimension- k hyperplane, to a cloud of data
- It turns out that the SVD solves this problem *immediately*!
- We first **de-mean** the data (subtract the mean from each component)
- Then the SVD gives the **best approximation** to a matrix A by a matrix of rank k as

$$A_k := \sum_{i=1}^k \sigma_i \mathbf{u}_i \mathbf{v}_i^T$$

- For a data matrix the SVD gives the best approximation by a k -dimensional subspace

Principal Component Analysis (PCA)

- Given an $(m \times n)$ **data matrix**

Principal Component Analysis (PCA)

- Given an $(m \times n)$ **data matrix**
- How many “important directions” are there in the data?
- i.e. what is the “dimension” of the data

Principal Component Analysis (PCA)

- Given an $(m \times n)$ **data matrix**
- How many “important directions” are there in the data?
- i.e. what is the “dimension” of the data

- Idea: We want to ignore “noise”

Principal Component Analysis (PCA)

- Given an $(m \times n)$ **data matrix**
- How many “important directions” are there in the data?
- i.e. what is the “dimension” of the data

- Idea: We want to ignore “noise”
- This is basically asking for the numerical rank!

PCA II

- Calculate the SVD

PCA II

- Calculate the SVD
- Plot the “importance” of each direction

$$\sigma_i \text{ or } \sigma_i^2 \text{ or } \frac{\sigma_i^2}{\sum_j \sigma_j^2}$$

as a function of i

PCA II

- Calculate the SVD
- Plot the “importance” of each direction

$$\sigma_i \text{ or } \sigma_i^2 \text{ or } \frac{\sigma_i^2}{\sum_j \sigma_j^2}$$

as a function of i

- Called a “**scree plot**”
- It often has an “elbow” after which the σ_i are small
- This helps to choose the k for the low-rank approximation

$$A_k := \sum_{i=1}^k \sigma_i \mathbf{u}_i \mathbf{v}_i^T$$

Summary

- The SVD decomposes the action of a matrix
 - rotation + stretch + rotation
- *Any* $(m \times n)$ matrix A has an SVD

$$A = U\Sigma V^T = \sum_i \sigma_i \mathbf{u}_i \mathbf{v}_i^T$$

- The SVD is closely related to the eigen-factorisation of the symmetric matrices $A^T A$ and AA^T
- Applications to analysing and compressing data