Nutritional Counseling - 7th Edition

Lesson 3 Test Questions

1. Approximately what proportion of infants who die within the first year of life die within the first month after birth?
a. 1/3
b. 2/3
c. 1/2
d. 3/4
e. 5/8

2. How can the change in lipid blood levels that occurs during pregnancy best be described?
a. Cholesterol and triglyceride levels decrease due to the increased water volume in the blood
b. Cholesterol and triglyceride levels remain the same as pre-pregnancy levels
c. Cholesterol and triglyceride levels increase dramatically from pre-pregnancy levels
d. Cholesterol and triglyceride levels fluctuate daily depending on when the fetus is building nerve cells

3. All of the following substances are transported through the placenta easily, with the exception of _____, which is not transferred at all.
a. water
b. cholesterol
c. oxygen
d. ketones
e. insulin

4. A critical period of spinal cord development following conception is:
A. 1-2 weeks after conception.
B. 3-4 weeks after conception.
C. 5-6 weeks after conception.
D. 7-8 weeks after conception.
E. 9-10 weeks after conception.

5. The recommended daily protein (g) intake for pregnant women is _____.
a. 30 g
b. 71 g
c. 90 g
d. 120 g

6. Research is emerging that suggests pregnant women should increase their food sources of docosahexaenoic acid (DHA). Which of the following foods would provide the most DHA to pregnant women?
a. 1/2 tuna salad sandwich
b. 1/4 cup granola containing 1 tsp sunflower seed
c. 8 oz glass whole milk
d. 1 cup broccoli

7. Pregnant women of _____ ethnicity are more likely than those of other ethnicities to have an eating disorder known as pica.
a. Chinese
b. African American
c. White Caucasian
d. Hispanic
e. Hmong

8. _____ do(es) NOT pose any foodborne bacterial risks in pregnant women.
a. Brie cheese
b. Ready-to-eat deli meats
c. Raw oysters
d. Unpasteurized milk
e. Organic bananas

9. Why must volume expansion occur BEFORE maternal nutrient stores accumulate?
a. In order to support large gains in fetal weight
b. In order to provide the fetus with sufficient energy, nutrients, and oxygen
c. In order to provide the mother with plenty of fluids
d. In order to dilute the high concentration of nutrients in pregnant women
e. In order to make room for growing organs

10. Which of the following is a major function of the placenta?
a. Hormone and enzyme production
b. Nutrient and gas exchange between mother and fetus
c. Removal of waste products from the fetus
d. Barrier to drugs and alcohol
e. a, b, and c
f. all of the above

Use the following information to answer questions 11-13:

Jane is 5'7" (1.70 m) and weighs 160 pounds (72.7 kg).

11. What is Jane's body mass index?
a. 19.7
b. 25.1
c. 31.4
d. 43

12. Based on her BMI, Jane would be considered:
a. underweight.
b. normal weight.
c. overweight.
d. obese.
e. morbidly obese.

13. Jane becomes pregnant. How many pounds should Jane gain during her pregnancy, based on her BMI?
a. 11 to 20
b. 15 to 25
c. 25 to 35
d. 28 to 40
e. 35 to 45

14. The recommended daily dietary intake of folate during pregnancy is:
a. 200 mcg from fortified food or supplements.
b. 400 mcg from fortified food or supplements.
c. 600 mcg total with 400 mcg from fortified food or supplements.
d. 800 mcg total with 600 mcg from fortified food or supplements.
e. 4000 mcg total in any form.

15. Hypertension is defined as having a blood pressure of _____.
a. ≥120 mm Hg systolic or ≥70 mm Hg diastolic
b. ≥140 mm Hg systolic or ≥90 mm Hg diastolic
c. ≥120 mm Hg systolic or ≥90 mm Hg diastolic
d. ≥140 mm Hg systolic or ≥70 mm Hg diastolic
e. ≥160 mm Hg systolic or ≥110 mm Hg diastolic

16. A worldwide leading cause of MATERNAL mortality is:
a. gestational diabetes.
b. hypertensive disorders of pregnancy.
c. AIDS/HIV.
d. multifetal pregnancies.
e. adolescent pregnancy complications.

17. Maternal organs most affected by small blot clots and reduced blood flow in preeclampsia include all of the following EXCEPT:
a. the placenta
b. the kidneys.
c. the brain.
d. the appendix.
e. the liver.

18. Nutritional and physical activity recommendations that may benefit women at risk for preeclampsia include all of the following EXCEPT:
a. high-dose iron supplements.
b. 5 or more servings of fruits and vegetables daily.
c. moderate exercise.
d. adequate fiber intake
e. adequate vitamin D.

19. High maternal blood glucose levels cause the fetus to _____.
a. increase insulin output
b. convert glucose into triglycerides and store it as fat
c. potentially develop type 2 diabetes later in life
d. All of the above

20. Which type of diabetes is potentially the most hazardous to the mother and fetus?
a. Type 1 diabetes
b. Type 2 diabetes
c. Gestational diabetes
d. They are all equally hazardous.

21. The vanishing twin phenomenon refers to:
a. the fact that more triplets and quadruplets are being born because of fertility drugs.
b. the disappearance of embryos within 13 weeks of conception.
c. the decreasing percentage of dizygotic twins in favor of monozygotes.
d. None of the above

22. The estimated minimum Calorie level for an overweight woman with gestational diabetes weighing 200 lb (91 kg) is _____.
a. 1200 Calories
b. 1820 Calories
c. 2000 Calories
d. 2275 Calories
e. 2500 Calories

23. A multifetal pregnancy increases the mother's risk for:
a. development of preeclampsia
b. development of iron-deficiency anemia
c. preterm delivery
d. cesarean delivery
e. All of the above

24. To achieve a 40-pound weight gain, women with twins need to consume approximately _____ more calories during pregnancy than do women with singleton pregnancies.
a. 15,000
b. 25,000
c. 35,000
d. 45,000
e. 50,000

25. "Best practice" vitamin and mineral supplement recommendations for women during multifetal pregnancies include which of the following statements?
a. There is no need to take a vitamin and mineral supplement if you are eating enough food.
b. Women with twins need to take more iron.
c. Women should use a prenatal vitamin and mineral supplement.
d. Women with twins need to take twice the recommended amount of a supplement.