NC Lesson Test 7

1. Studies show that children of parents who impose authoritarian methods of control on their eating are:
a. likely to have healthy eating habits
b. less likely to regulate their intake
c. more likely to become overweight
d. prone to developing fewer eating-related issues
e. Both b and c

2. What is the recommended adequate intake of total fiber for children between the ages of 4 and 8?
a. 25 g/day
b. 29 g/day
c. 31 g/day
d. 35 g/day

3. Which of the following foods would be MOST likely to promote dental caries?
a. Raisins
b. Apple slices
c. Cheddar cheese
d. Peanut butter
e. Plain yogurt

4. “BMI rebound” can be defined as:
a. the period in which your weight “rebounds” and settles in at your normal weight.
b. the normal increase in percent body fat which occurs after percent body fat declines and reaches its lowest point.
c. the point at which you have the greatest amount of fat you will ever have, typically between the ages of 8 and 10.
d. the weight you attain by the age of 13 after you have had a growth spurt.
e. the loss of all a child’s baby fat, which occurs at about 6 years of age.

5. The most significant predictor of childhood obesity has been found to be:
a. low family income.
b. food insecurity.
c. Maternal obesity.
d. lower cognitive stimulation.
e. use of formula instead of breastfeeding.

6. Which of the following foods would be the best choice for a parent trying to increase fiber in her/his child’s diet?
a. Protein or energy bar
b. Fresh fruits, vegetables and whole-grain breads

c. Fresh juice and enriched bread
d. Canned fruits
e. All of the above would be great sources of fiber.

7. The American Academy of Pediatrics (AAP) advocates for the elimination of sweetened beverages in school and suggests replacing them with all of the following EXCEPT:
a. Low-fat chocolate milk
b. Low-fat milk
c. Vegetable juice
d. 100% fruit juice
e. Sports beverages

8. **Using the following information, answer questions 8-11.**

Jane is an 8-year-old girl that weighs 56.5 pounds (25.6 kg), is 51 inches tall, and enjoys lots of outdoor activities. Her total caloric intake is 1800 Calories per day consisting of 80 grams of fat, 225 grams of carbohydrate, and 45 grams of protein. She likes vegetables and eats lots of fruit; the only food she truly dislikes is milk.

8. Jane gets _____% of her energy from fat.
Select one:
a. 20
b. 25
c. 30
d. 35
e. 40

9. Is this within the Acceptable Macronutrient Distribution Range (AMDR) for fat for a child her age?
a. Yes
b. No
c. Cannot be determined from the information provided

10. How many grams of protein per day are recommended for Jane?
a. 24 g
b. 15 g
c. 30 g
d. 20 g
e. 50 g

11. Is she eating more, less, or the recommended amount of protein compared to the DRI for a girl her age?
a. More
b. Less
c. Exactly as much as the DRI recommends

12. Examples of conditions that INCREASE protein needs include:
a. wound healing.
b. cystic fibrosis.
c. PKU.
d. All of the above
e. a and b only

13. Fat-soluble vitamin supplements are recommended for children with _____ due to poor intestinal absorption of these nutrients.
a. diabetes
b. seizures
c. PKU
d. cystic fibrosis
e. all of the above

14. Energy and protein needs for children with cystic fibrosis are _____ those of healthy children.
a. less than
b. the same as
c. two to four times greater than
d. greater than four times

15. Type 1 diabetes is related to immune function and results in virtually _____.
a. no immune system regulation
b. no colds for the first few years of the disease
c. no ability to process all macronutrients
d. no regulation of appetite
e. no insulin production

16. Nutritional consequences of spastic quadriplegia include:
a. rapid weight gain.
b. predictable 2-fold increases in energy needs.
c. changes in body composition.
d. extreme vitamin and mineral deficiencies.

17. A psychostimulant is a medication given to children with _____.
a. seizures
b. attention deficit hyperactivity disorder
c. phenylketonuria
d. type 1 diabetes

18. What is the importance of taking pancreatic enzymes when a child has cystic fibrosis?
a. These enzymes help digest foods and when they are missing, malabsorption of various nutrients occurs
b. Pancreatic enzymes also supply the fat-soluble vitamins, which are necessary for normal growth.
c. It is not necessary to take pancreatic enzymes with cystic fibrosis; you need to take nutritional
supplements, however.
d. Without these enzymes, inflammation occurs.
e. These enzymes help with calcium absorption.

19. Which of the following foods would be the best breakfast choice for a child on a ketogenic diet?
a. A bowl of frosted Mini-Wheat, milk, and orange juice
b. Homemade oatmeal with milk, raisins, and brown sugar
c. Scrambled eggs with bacon and cheese
d. Toast and jelly with a glass of grape juice
e. A doughnut with a glass of chocolate milk

20. Which of the following is NOT a secondary effect of cerebral palsy?
a. Dehydration
b. Constipation
c. Contractures
d. Gastroesophageal reflux