Lesson Test 8

1. In order to improve the eating habits of youth, nutrition interventions should be aimed at three levels of influence. Which of the following would NOT be one of these levels?
a. Environmental factors
b. Macrosystem factors
c. Transportation factors
d. Personal factors

2. Snacks account for up to _____% of daily food intake among adolescents.
a. 15-20
b. 23-27
c. 27-35
d. 41-44
e. 51-55

3. The best reason for advising young adolescents to consume foods that provide adequate calcium is _____.
a. young adolescents expend more energy from increased physical activity and need more calcium
b. young adolescents do not have much money and dairy foods are the least expensive to buy
c. young adolescents retain four times as much calcium than adults and build bone mass
d. young adolescents fight with their parents a lot and this would eliminate one food battle

4. All of the following are factors that increase a teen's need for iron EXCEPT:
a. A rapid rate of linear growth.
b. The onset of menarche for females.
c. An increase in blood volume.
d. Greater intake of heme than nonheme iron sources.
e. All of the above factors increase a teen's need for iron.

5.Susan is a 15-year-old who likes fruits, some vegetables, and breads (whole-grain and enriched) but refuses to consume milk, seafood, and eggs. A recent doctor visit revealed normal hemoglobin and plasma ferritin levels. What would you advise?
a. A folate supplement (400 micrograms/day)
b. A vitamin D supplement (400 IU/day)
c. An iron supplement (10 mg/day)
d. No changes in nutrient intake

6. An underweight 17-year-old male was 15 minutes late to his nutrition counseling appointment and he has to leave for a school-related practice in 20 minutes. What dietary assessment method would be the most appropriate for the dietitian to begin assessing his food intake pattern as well as a method that she could reliably complete on the telephone at a later time?
a. 24-hour recall

b. Food frequency
c. Food record
d. Diet history

7. Which of these messages is best for motivating adolescents to eat a healthy diet?
a. “Eating well will give you energy for sports or social activities.”
b. ”Eat what you don’t like because it’s good for you.”
c. ”Eating well may prevent heart disease in the long term.”
d. ”Eat well now so you won’t be frail or disabled when you are old.”

8. For which 16-year-old teen would the dietitian have the greatest concern about folate status?
a. Male whose estimated intake is 450 micrograms per day
b. Female who consumes orange juice and cereal for breakfast
c. Male with serum folate levels just below the normal range
d. Female who is sexually active and skips breakfast

9. Youth with body mass index (BMI) values > 50th percentile, but < 75th percentile are considered:
a. normal weight.
b. at risk for overweight.
c. overweight.
d. obese.
e. morbidly obese.

10. As many as _____ of overweight adolescents can be expected to remain overweight into adulthood.
a. 40%
b. 50%
c. 60%
d. 70%
e. 90%

11. Each pound of body weight lost during an activity requires ingestion of _____ ounces of fluid
following the activity to maintain proper hydration.
a. 8
b. 16
c. 20
d. 32
e. 48

Use the Tapango's menu provided below to answer questions 12-14.

Breakfast Burrito
nutritional information: 412 Calories, 17g fat, 6.6g saturated fat, 44.9g carbs per 200g burrito
ingredients: flour tortilla, scrambled eggs, salsa, and cheddar cheese
Breakfast Burrito with Bacon
nutritional information: 657 Calories, 35.9g fat, 12.9g saturated fat, 45.5g carbs per 245g burrito
ingredients: flour tortilla, scrambled eggs, bacon, salsa, and cheddar cheese
Primo Breakfast Burrito
nutritional information: 761 Calories, 43.6g fat, 16.8g saturated fat, 54.2g carbs per 402g burrito
ingredients: flour tortilla, scrambled eggs, bacon, cheddar cheese, salsa, lettuce, chili sauce, sour cream, tomatoes, olives, onions, jalapeño peppers
Breakfast Quesadilla
nutritional information: 478 Calories, 23.9g fat, 11.8g saturated fat, 41.7g carbs per 171g quesadilla
ingredients: flour tortilla, scrambled eggs, and cheddar cheese
Breakfast Quesadilla with Bacon
nutritional information: 726 Calories, 43g fat, 18.1g saturated fat, 42.5g carbs per 217g quesadilla
ingredients: flour tortilla, scrambled eggs, bacon, and cheddar cheese
Primo Nachos
nutritional information: 964 Calories, 49.3g fat, 14.3g saturated fat, 102.7g carbs per 445g nachos
ingredients: tortilla chips, nacho cheese, refried beans, chicken (chicken breast, green chili peppers, olive oil, onions, garlic), salsa, lettuce, chili sauce, sour cream, tomatoes, olives, onions, and jalapeño peppers
Primo Taco Salad
nutritional information: 502 Calories, 20.9g fat, 5.5g saturated fat, 62.6g carbs per 448g taco salad
ingredients: lettuce, chicken (chicken breast, green chili peppers, olive oil, onions, garlic, and spices), tortilla chips, refried beans, Spanish rice (rice, olive oil, tomato sauce, onions, garlic, and
spices), salsa, lettuce, chili sauce, sour cream, tomatoes, olives, onions, and jalapeño peppers
Beef Taco
nutritional information: 214 Calories, 11.7g fat, 5.1g saturated fat, 14.5 carbs per 128g taco
ingredients: corn tortilla, beef (ground beef, tomato sauce, onions, and spices), chili sauce, lettuce, tomatoes, and cheddar cheese
Chicken Taco
nutritional information: 154 Calories, 5.3g fat, 2.4g saturated fat, 14.1g carbs per 128g taco
ingredients: corn tortilla, chicken (chicken breast, green chili peppers, olive oil, onions, garlic, and spices), chili sauce, lettuce, tomatoes, and cheddar cheese

12. An adolescent is considering Tapango's menu and would like to choose the food with the lowest percentage of fat calories. This would be the:
a. breakfast burrito w/ bacon.
b. breakfast quesadilla w/ bacon.
c. primo taco salad.
d. beef taco.
e. primo nachos.

13. Would this food fit within the current recommendations for fat consumption?
a. No, it is high at 37% fat
b. No, it is high at 47% fat
c. No, it is low at 17% fat
d. Yes, it is only 30% fat
e. Need more information

14. What beverage would be a good accompaniment to this meal and provide a nutrient that is usually scarce in adolescent diets?
a. Non-fat milk
b. Diet soda
c. Water
d. Regular soda
e. Iced tea with sweetener

15. A juice or sports drink that contains no more than _____ carbohydrate may allow for better hydration and physical performance than those that are more concentrated.
a. 2-3%
b. 4-5%
c. 5-6%
d. 6-8%
e. 9-10%

16. A teenage girl ate Christmas dinner at her step-mother's house and following a creamy dessert, announced, "I am fatter than all of my classmates; I am going to stop eating for 5 days!" Important behaviors the step-mother should model include all of the following EXCEPT:
a. eating well-balanced meals as part of her normal routine.
b. showing the step-daughter love without using food.
c. identifying strengths of the step-daughter that are not eating related.
d. beginning a subscription to Weight Watcher's magazine for the stepdaughter.

17. Binge-eating disorder differs from bulimia nervosa in that binge-eating disorder:
a. does not include binges on "fast" or "junk" foods.
b. does not involve eating large amounts of food in a short period of ,me.
c. does include large amounts of fruits and vegetables during a binge episode.
d. does not involve compensatory behaviors.

e. involves stool softeners, not laxatives.

18. Key goals of nutritional care for someone in treatment for an eating disorder would include:
a. establishing a regular pattern of nutritionally balanced meals.
b. gradually introducing forbidden foods into the diet.
c. avoiding dieting behaviors and excessive exercise.
d. thoroughly assessing dietary intake and adequacy.
e. All of the above are key goals

19. An estimated _____ percent of children and adolescents with a chronic condition or disability also have nutrition risk factors that warrant a referral to a dietitian.
a. 20
b. 30
c. 40
d. 50

Use the following nutrition label to answer questions 20-22.

Nutrition Facts Serving Size 1 Drink Box (6.75 oz) Servings Per Container 1	
Amount Per Serving Calories 100	Calories from Fat 0
	% Daily Value*
Total Fat 0g	0%
Saturated Fat 0g	0%
Trans Fat 0g	
Cholesterol 0mg	0%
Sodium 25mg	1%
Total Carbohydrate 27g	10%
Dietary Fiber 0g	
Sugars 24g	
Protein 1g	2%
Vitamin A 0% •	Vitamin C 2%
Calcium 10% •	Iron 0%

20. A coach bought fruit juice boxes (see food label above) for the wrestling team to provide energy to wrestlers after they weighed in. What is the best recommendation for teens when they consume this 30 minutes before the meet?
a. Dilute half of this juice with 6.75 oz cold water
b. Drink three juice boxes to get adequate fluid volume before the match.
c. Only drink 20 oz of water before the match.
d. Find a vending machine and purchase a sports drink containing caffeine.

21. Based on the calories in this drink, calculate the percentage of total calories from carbohydrates it provides.
a. 14
b. 40
c. 50
d. 100

22. Would this beverage be a good choice to drink during a strenuous activity?
a. Yes: it is 100% fruit juice, so it is a good choice
b. No: it is too high in carbohydrate and may cause cramping
c. Not enough information is provided on the label

23. What iron-rich food source would be the best choice for an adolescent who is a lactovegetarian?
a. Pinto beans and diced tomatoes
b. Scrambled eggs with cheese
c. Tuna sandwich
d. Flour tortilla with cheese
e. Fish sandwich with lettuce

24. What dietary components are MOST effective at raising muscle glycogen concentrations?
a. Fats
b. Proteins
c. Carbohydrates
d. Chromium & iron-containing supplements

25. The energy needs of adolescents are influenced by basal metabolic rate, activity level, and increased requirements to support pubertal growth and development.
True
False