

1) Consider the unconstrained optimization problem

$$\begin{cases} \min & 2x_1^2 + x_2^2 - x_1x_2 + e^{x_1+2x_2} \\ & x \in \mathbb{R}^2 \end{cases}$$

(a) Prove that the problem admits a global minimum;

(b) Apply the gradient method with an inexact line search, setting  $\bar{t} = 1, \alpha = 0.1, \gamma = 0.8$ , with starting point  $x^0 = (-10, 8)$  and using  $\|\nabla f(x)\| < 10^{-6}$  as stopping criterion. How many iterations are needed by the algorithm? Write explicitly the vectors found at the last three iterations.

(c) Is the obtained solution a global minimum of the given problem? Justify the answer.

### SOLUTION

(a) The objective function  $f(x) = f_1(x) + f_2(x)$ , where  $f_1 = 2x_1^2 + x_2^2 - x_1x_2$  is strongly convex and  $f_2(x) = e^{x_1+2x_2}$  is convex, being  $f_2 = \psi \circ g$ , with  $\psi(y) = e^y$  convex and  $g(x) = x_1 + 2x_2$  linear. Therefore  $f_1 + f_2$  is strongly convex being the sum of a strongly convex plus a convex function. Consequently,  $f$  admits a unique global minimum point.

(b) We notice that

$$\nabla f(x_1, x_2) = \begin{pmatrix} 4x_1 - x_2 + e^{x_1+2x_2} \\ -x_1 + 2x_2 + 2e^{x_1+2x_2} \end{pmatrix}$$

### Matlab solution

```
%% Data
```

```
alpha = 0.1;
gamma = 0.8;
tbar = 1;
x0 = [-10;8];
tolerance = 10^(-6) ;
```

```
X=[ ];
```

```
ITER = 0 ;
x = x0;
```

```
while true
    [v, g] = f(x);
```

```
    X=[X;ITER,x(1),x(2),v,norm(g)];
```

```
    % stopping criterion
    if norm(g) < tolerance
        break
    end
```

```
    d = -g; % search direction
```

```
    t = tbar ; % Armijo inexact line search
    while f(x+t*d) > v + alpha*g'*d*t
        t = gamma*t ;
    end
```

```
    x = x + t*d ; % new point
    ITER = ITER + 1 ;
end
```

```
disp('optimal solution')
x
v
norm(g)
ITER
```

```
function [v, g] = f(x)

v = 2*x(1)^2 + x(2)^2 - x(1)*x(2) + exp(x(1)+2*x(2)) ;

g = [4*x(1)-x(2)+exp(x(1)+2*x(2));
     -x(1)+2*x(2)+2*exp(x(1)+2*x(2))];

end
```

We obtain the following solution:

```
x =

    -0.1952
    -0.4393

v =

    0.5251

ans =

    6.6986e-07
```

```
ITER =

    24
```

In particular, the gradient norm evaluated at the final point is:

```
ans =

    6.6986e-07
```

The iterations of the algorithm are 24.

The vectors found at the last three iterations are:

```
-0.1952    -0.4393
-0.1953    -0.4393
-0.1952    -0.4393
```

(b) The found point  $x = (-0.1952, -0.4393)$  is a global minimum since the objective function is strongly convex as shown in point (a).

2) Consider a regression problem with the following data set where the points  $(x_i, y_i)$ ,  $i = 1, 31$ , are given by the row vectors of the matrices:

$$\begin{pmatrix} -3.0000 & 4.58 \\ -2.8000 & 7.19 \\ -2.6000 & 8.22 \\ -2.4000 & 16.06 \\ -2.2000 & 16.42 \\ -2.0000 & 17.53 \\ -1.8000 & 11.48 \\ -1.6000 & 14.10 \\ -1.4000 & 16.82 \\ -1.2000 & 16.15 \\ -1.0000 & 11.68 \\ -0.8000 & 6.00 \\ -0.6000 & 7.82 \\ -0.4000 & 2.82 \\ -0.2000 & 2.71 \\ 0 & 1.16 \end{pmatrix} \quad \begin{pmatrix} 0.2000 & -1.42 \\ 0.4000 & -3.84 \\ 0.6000 & -4.71 \\ 0.8000 & -8.15 \\ 1.0000 & -7.33 \\ 1.2000 & -13.64 \\ 1.4000 & -15.26 \\ 1.6000 & -14.87 \\ 1.8000 & -9.92 \\ 2.0000 & -10.50 \\ 2.2000 & -7.72 \\ 2.4000 & -11.78 \\ 2.6000 & -10.26 \\ 2.8000 & -7.13 \\ 3.0000 & -2.11 \end{pmatrix}$$

- Write the dual formulation of a nonlinear  $\varepsilon$ -SV regression model with  $C = 5$ ,  $\varepsilon = 3$  and a polynomial kernel  $k(x, y) := (x^T y + 1)^4$ ;
- Solve the problem in (a) and find the regression function;
- Find the support vectors;
- Find the points of the data set that are outside the  $\varepsilon$ -tube, by making use of the dual solution.

### SOLUTION

(a) Let  $\ell = 31$ ,  $(x_i, y_i)$ ,  $i = 1, \dots, \ell$  be the  $i$ -th element of the data set,  $C = 5$ ,  $\varepsilon = 3$ ,  $k(x, y) := (x^T y + 1)^4$ . The dual formulation of a nonlinear  $\varepsilon$ -SV regression model is

$$\begin{cases} \max_{(\lambda^+, \lambda^-)} & -\frac{1}{2} \sum_{i=1}^{\ell} \sum_{j=1}^{\ell} (\lambda_i^+ - \lambda_i^-)(\lambda_j^+ - \lambda_j^-) k(x_i, x_j) \\ & -\varepsilon \sum_{i=1}^{\ell} (\lambda_i^+ + \lambda_i^-) + \sum_{i=1}^{\ell} y_i (\lambda_i^+ - \lambda_i^-) \\ \sum_{i=1}^{\ell} (\lambda_i^+ - \lambda_i^-) & = 0 \\ \lambda_i^+, \lambda_i^- & \in [0, C], i = 1, \dots, \ell \end{cases} \quad (1)$$

### (b) Matlab solution

```
data = [
-3.0000    4.58
-2.8000    7.19
-2.6000    8.22
-2.4000   16.06
-2.2000   16.42
-2.0000   17.53
-1.8000   11.48
-1.6000   14.10
-1.4000   16.82
-1.2000   16.15
-1.0000   11.68
-0.8000    6.00
-0.6000    7.82
-0.4000    2.82
-0.2000    2.71
    0     1.16
 0.2000   -1.42
 0.4000   -3.84
 0.6000   -4.71
 0.8000   -8.15
 1.0000   -7.33
 1.2000  -13.64
 1.4000  -15.26
 1.6000  -14.87
```

```

1.8000    -9.92
2.0000   -10.50
2.2000    -7.72
2.4000   -11.78
2.6000   -10.26
2.8000    -7.13
3.0000    -2.11
    ];

x = data(:,1) ;
y = data(:,2) ;
l = length(x) ;

epsilon = 3 ;
C = 5;

X = zeros(l,l);
for i = 1 : l
    for j = 1 : l
        X(i,j) = kernel(x(i),x(j)) ;
    end
end
Q = [ X -X ; -X X ];
c = epsilon*ones(2*l,1) + [-y;y];

sol = quadprog(Q,c,[],[],[ones(1,l) -ones(1,l)],0,zeros(2*l,1),C*ones(2*l,1));
lap = sol(1:l);
lam = sol(l+1:2*l);

% compute b
ind = find(lap > 1e-3 & lap < C-1e-3);
if isempty(ind)==0
    i = ind(1);
    b = y(i) - epsilon;
    for j = 1 : l
        b = b - (lap(j)-lam(j))*kernel(x(i),x(j));
    end
else
    ind = find(lam > 1e-3 & lam < C-1e-3);
    i = ind(1);
    b = y(i) + epsilon ;
    for j = 1 : l
        b = b - (lap(j)-lam(j))*kernel(x(i),x(j));
    end
end

z = zeros(l,1);
for i = 1 : l
    z(i) = b ;
    for j = 1 : l
        z(i) = z(i) + (lap(j)-lam(j))*kernel(x(i),x(j));
    end
end
zp = z + epsilon ;
zm = z - epsilon ;

sv = [find(lap > 1e-3);find(lam > 1e-3)];
sv = sort(sv);

plot(x,y,'b.',x(sv),y(sv),'ro',x,z,'k-',x,zp,'r-',x,zm,'r-');
disp('Support vectors')

```

% find regression and epsilon-tube

% find support vectors

% plot the solution

```
[sv,x(sv),y(sv),lam(sv),lap(sv)]    % Indexes of support vectors, support vectors, lambda_-,lambda_+
```

```
function v = kernel(x,y)
p = 4 ;
v = (x'*y + 1)^p;
end
```

Let  $\lambda_-$  and  $\lambda_+$  be the vectors given by the Matlab solutions lam, lap. In particular we find,  $b = -0.008$ .

The regression function is:

$$f(x) = \sum_{i=1}^{\ell} (\lambda_i^+ - \lambda_i^-) k(x_i, x) + b = \sum_{i=1}^{31} (\lambda_i^+ - \lambda_i^-) (x_i x + 1)^4 - 0.008$$

(c) We obtain the support vectors (columns 2-3) and corresponding  $\lambda_-$  and  $\lambda_+$  (columns 4-5) :

ans =

4.0000	-2.4000	16.0600	0.0000	1.2569
7.0000	-1.8000	11.4800	5.0000	0.0000
9.0000	-1.4000	16.8200	0.0000	1.3176
10.0000	-1.2000	16.1500	0.0000	5.0000
12.0000	-0.8000	6.0000	0.5720	0.0000
23.0000	1.4000	-15.2600	4.7410	0.0000
27.0000	2.2000	-7.7200	0.0000	5.0000
29.0000	2.6000	-10.2600	2.2616	0.0000

(d) Consider the feasibility condition of the primal formulation of the regression problem:

$$y_i - f(x_i) - \varepsilon - \xi_i^+ \leq 0, \quad y_i - f(x_i) + \varepsilon + \xi_i^- \geq 0, \quad i = 1, \dots, \ell$$

If a point  $(x_i, y_i)$  is outside the  $\varepsilon$ -tube then  $\xi_i^+ > 0$  or  $\xi_i^- > 0$ .

Given the dual optimal solution  $(\lambda_+, \lambda_-)$  of (1), we can find the errors  $\xi_i^+$  and  $\xi_i^-$  associated with the point  $(x_i, y_i)$  by the complementarity conditions:

$$\begin{cases} \lambda_i^+ [y_i - f(x_i) - \varepsilon - \xi_i^+] = 0, & i = 1, \dots, \ell \\ \lambda_i^- [y_i - f(x_i) + \varepsilon + \xi_i^-] = 0, & i = 1, \dots, \ell \\ \xi_i^+ (C - \lambda_i^+) = 0, & i = 1, \dots, \ell \\ \xi_i^- (C - \lambda_i^-) = 0, & i = 1, \dots, \ell \end{cases} \quad (2)$$

it follows that a necessary condition for a point  $(x_i, y_i)$  to be outside the  $\varepsilon$ -tube is that  $\lambda_i^+ = C = 5$  or  $\lambda_i^- = C = 5$ . We find that  $\lambda_i^- = 5$ , for  $i = 7$ ,  $\lambda_i^+ = 5$ , for  $i = 10, 27$  which correspond to the points

$$(x_7, y_7) = (-1.8, 11.48), \quad (x_{10}, y_{10}) = (-1.2, 16.15), \quad (x_{27}, y_{27}) = (2.2, -7.72)$$

3) Consider the following unconstrained multiobjective optimization problem:

$$\begin{cases} \min f(x_1, x_2) = (x_1 x_2, x_1^2 + 2x_2^2 - 2x_2) \\ x \in \mathbb{R}^2 \end{cases}$$

- (a) Prove that the problem admits a Pareto minimum point.  
 (b) Find a suitable subset of Pareto minima, by means of the scalarization method.

### SOLUTION

- (a) We observe that the function  $f_2$  is strongly convex, so that its minimum point is a Pareto minimum of the given problem.  
 (b) Consider the scalarized problem  $(P_{\alpha_1})$ , where  $0 \leq \alpha_1 \leq 1$ , i.e.

$$\begin{cases} \min \alpha_1 x_1 x_2 + (1 - \alpha_1)(x_1^2 + 2x_2^2 - 2x_2) =: \psi_{\alpha_1}(x) \\ x \in \mathbb{R}^2 \end{cases}$$

We note that the objective function  $\psi_{\alpha_1}(x)$  is quadratic with Hessian given by

$$Q = \begin{pmatrix} 2(1 - \alpha_1) & \alpha_1 \\ \alpha_1 & 4(1 - \alpha_1) \end{pmatrix}$$

We observe that  $Q$  is positive definite for  $0 \leq \alpha_1 < \frac{8-2\sqrt{2}}{7} \approx 0.7388$  so that for such values  $P_{\alpha_1}$  is strongly convex and admits a unique global solution which is a Pareto minimum.

$P_{\alpha_1}$  can be solved by Matlab for  $0 \leq \alpha_1 \leq 1$ :

```
Q1 = [0 1; 1 0];
Q2 = [2 0; 0 4];
c1=[0 0]';
c2=[0 -2]';
```

```
MINIMA=[]; SOL=[];
```

```
for alfa1 = 0 : 0.001 : 1
    eigQalfa1= eig(alfa1*Q1+(1-alfa1)*Q2);
    eigQ=[eigQ;alfa1,eigQalfa1']; % Eigenvalues of the hessian of \psi_alfa1(x)
```

```
if (eigQ > 0.001)
    [x,fval,exitflag,exitflag] = fminunc(@(x) 0.5*x'*(alfa1*Q1+(1-alfa1)*Q2)*x + (alfa1*c1+(1-alfa1)*c2)'*x, [0,0]);
    MINIMA=[MINIMA; alfa1 x'];
```

```
else
    [x,fval,exitflag,exitflag] = fminunc(@(x) 0.5*x'*(alfa1*Q1+(1-alfa1)*Q2)*x + (alfa1*c1+(1-alfa1)*c2)'*x, [0,0]);
    SOL=[SOL; alfa1 x'];
end
end
```

```
plot(MINIMA(:,2),MINIMA(:,3), 'r*')
```

For every  $0 \leq \alpha_1 \leq 0.738$ , the problem admits a unique solution which is a Pareto minimum (MINIMA(:,2:3)). In particular,

- For  $\alpha_1 = 0$  we obtain the point  $(0, 0.5)$
- For  $\alpha_1 = 0.736$  we obtain the point  $(-24.4839, 17.5645)$ ;
- For  $\alpha_1 = 0.737$  we obtain  $(-38.0695, 27.1704)$
- For  $\alpha_1 = 0.738$  we obtain  $(-85.7834, 60.9085)$
- For  $\alpha_1 \geq 0.739$  the problem  $P_{\alpha_1}$  is unbounded.

The previous values lead us to conjecture that the sequence of minima diverges as  $\alpha_1 \rightarrow \frac{8-2\sqrt{2}}{7} \approx 0.7388$

We note that the previous solutions can also be obtained by the stationarity conditions for  $(P_{\alpha_1})$  which are necessary for a weak minimum point. Then, the following system provides a necessary condition for an optimal solution of  $(P_{\alpha_1})$  and such condition is also sufficient for  $0 \leq \alpha_1 < \frac{8-2\sqrt{2}}{7}$ :

$$\begin{cases} \alpha_1 x_2 + 2(1 - \alpha_1)x_1 = 0 \\ \alpha_1 x_1 + (1 - \alpha_1)(4x_2 - 2) = 0 \\ 0 \leq \alpha_1 \leq 1, \end{cases}$$

Sistema KKT

We obtain:

$$\begin{cases} x_1 = \frac{2\alpha_1(1 - \alpha_1)}{-7\alpha_1^2 + 16\alpha_1 - 8}, \\ x_2 = \frac{4(1 - \alpha_1)^2}{7\alpha_1^2 - 16\alpha_1 + 8} \\ 0 \leq \alpha_1 \leq 1, \\ \alpha_1 \neq \frac{8-2\sqrt{2}}{7} \end{cases} \quad (3)$$

In particular, for  $\alpha_1 = \frac{8-2\sqrt{2}}{7}$  the previous system is impossible.

We obtain the set of points

$$\begin{cases} x_1 = \frac{2\alpha_1(1 - \alpha_1)}{-7\alpha_1^2 + 16\alpha_1 - 8}, \\ x_2 = \frac{4(1 - \alpha_1)^2}{7\alpha_1^2 - 16\alpha_1 + 8} \\ 0 \leq \alpha_1 < \frac{8-2\sqrt{2}}{7} \end{cases}$$

which are Pareto minima.

For  $1 \geq \alpha_1 > \frac{8-2\sqrt{2}}{7}$  is not a minimum of  $(P_{\alpha_1})$  being the Hessian matrix  $Q$  indefinite as can also be checked by the previous Matlab program.

4) Consider the following matrix game:

$$C = \begin{pmatrix} 1 & 4 & -1 & 5 & 2 \\ 2 & 1 & 3 & 3 & 5 \\ 2 & 3 & -2 & 3 & 1 \\ 1 & 1 & 5 & 2 & 3 \end{pmatrix}$$

- (a) Find the dominated strategies and reduce the cost matrix accordingly;
- (b) Find the set of pure strategies Nash equilibria, if any. Alternatively, show that no pure strategies Nash equilibrium exists.
- (c) Find a mixed strategies Nash equilibrium.

### SOLUTION

(a) Strategy 1 of Player 2 is dominated by Strategy 4, so that column 1 can be deleted. The reduced game is given by the matrix

$$C_{R1} = \begin{pmatrix} 4 & -1 & 5 & 2 \\ 1 & 3 & 3 & 5 \\ 3 & -2 & 3 & 1 \\ 1 & 5 & 2 & 3 \end{pmatrix}$$

Now, Strategy 1 of Player 1 is dominated by Strategy 3, so that row 1 can be deleted and the reduced matrix becomes:

$$C_{R2} = \begin{pmatrix} 1 & 3 & 3 & 5 \\ 3 & -2 & 3 & 1 \\ 1 & 5 & 2 & 3 \end{pmatrix}$$

(b) We observe that no minimum component on the columns of the reduced matrix is a maximum on the respective row, so that no pure strategies Nash equilibrium exists.

(c) The optimization problem associated with Player 1 is

$$\begin{cases} \min v \\ v \geq x_1 + 2x_2 + 2x_3 + x_4 \\ v \geq 4x_1 + x_2 + x_3 + x_4 \\ v \geq -x_1 + 3x_2 - 2x_3 + 5x_4 \\ v \geq 5x_1 + 3x_2 + 3x_3 + 2x_4 \\ v \geq 2x_1 + 5x_2 + x_3 + 3x_4 \\ x_1 + x_2 + x_3 + x_4 = 1 \\ x \geq 0 \end{cases} \quad (4)$$

The previous problem can be solved by Matlab.

#### Matlab solution

```
C=[1,4,-1,5,2; 2 1 3 3 5; 2 3 -2 3 1;1 1 5 2 3]
```

```
m = size(C,1);
n = size(C,2);
c=[zeros(m,1);1];
A= [C', -ones(n,1)]; b=zeros(n,1); Aeq=[ones(1,m),0]; beq=1;
lb= [zeros(m,1);-inf]; ub=[ ];
[sol,Val,exitflag,output,lambda] = linprog(c, A,b, Aeq, beq, lb, ub);
x = sol(1:m)
y = lambda.ineqlin
```

We obtain the optimal solution  $(x, v) = (0, 0, \frac{3}{8}, \frac{5}{8}, 2.375)$ . The optimal solution of the dual of (4) is given by  $(y, w) = (0, 0, \frac{1}{8}, \frac{7}{8}, 0, 2.375)$ .  $y$  can be found in the vector *lambda.ineqlin* given by the Matlab function *linprog*.

Therefore,

$$(x_1, x_2, x_3, x_4) = (0, 0, \frac{3}{8}, \frac{5}{8}), \quad (y_1, y_2, y_3, y_4, y_5) = (0, 0, \frac{1}{8}, \frac{7}{8}, 0),$$

is a mixed strategies Nash equilibrium.



