



Special article

Nutritional support of the elderly cancer patient: Long-term nutritional support



Stéphane M. Schneider M.D., Ph.D., F.E.B.G.H. *, Xavier Hébuterne M.D., Ph.D.

Faculté de Médecine, Gastroentérologie et Nutrition Clinique, Université de Nice Sophia-Antipolis, Pôle Digestif-Gynécologie-Obstétrique, Centre Hospitalier Universitaire de Nice, Nice, France

ARTICLE INFO

Article history:

Received 17 November 2014

Accepted 24 November 2014

Keywords:

Elderly

Cancer

Home artificial nutrition

Enteral nutrition

Parenteral nutrition

ABSTRACT

Elderly cancer patients account for a growing part of home artificial nutrition patients. Long-term enteral or parenteral nutrition in the older patient with cancer is prescribed for sequels after treatment (dysphagia, intestinal failure) or for bowel obstruction. Home artificial nutrition should benefit from a specialized follow-up. For patients out of therapy, the goal of nutritional care is to optimize quality of life and comfort.

© 2015 Elsevier Inc. All rights reserved.

Remarkably few studies have addressed the subject of long-term (>3 mo) nutritional support in older patients, let alone those with cancer. Therefore, recommendations tend to be made based on those available for adult patients.

Indications for long-term artificial nutrition

Indications for prolonged enteral nutrition (EN) in patients with usually include dysphagia and mild intestinal failure (gastrectomy and Whipple's procedure), both induced by cancer treatment. Indications for prolonged parenteral nutrition (PN) in these patients include severe intestinal failure (surgical resection, radiation enteritis) induced by cancer treatment. They also include chronic obstruction that may be due to peritoneal carcinomatosis, nowadays compatible with prolonged survival (especially when the primary tumor is ovarian) [1]. These indications are similar to those in younger patients, both in the hospital and at home. Guidelines from the European Society for Clinical Nutrition and Metabolism (ESPEN) state that, although indications of PN should be limited to situations when EN is contraindicated or poorly tolerated, age per se is not a reason for excluding patients from PN [2].

In patients with advanced cancer and inoperable bowel obstruction, multicenter studies on home PN (HPN) have demonstrated a positive effect on nutritional status and quality

of life in those who survived >3 mo and had a Karnofsky Performance Status score of >50 at the onset of HPN [3,4]. In patients where death is imminent (i.e., within the next 4 wk), PN or hydration should only be used after careful and interdisciplinary reflection, especially because the ability to estimate the length of survival is quite limited.

Patient comfort is the highest priority, and nutritional support should be given in conjunction with other palliative treatments [2].

In Europe in 1997, individuals age >61 represented 28% of all HPN patients [5], a percentage comparable to that observed in the United States (22%) [6]. Between 2001 and 2005, 81% of home EN (HEN) patients in the Veneto region of Italy were age >65, and the majority of head and neck and abdominal cancers were in this age group [7]. In the U.S. registry analysis of geriatric patients, cancer was the main reason for HEN and HPN (43% for each) [6].

ESPEN guidelines on EN in the geriatric population state that for long-term nutritional support, gastrostomy tubes may be preferred to nasogastric tubes because they are associated with fewer treatment failures, better nutritional status, and are more convenient for the patient [6]. Similarly, HPN should be performed via a specialized central venous line (tunneled catheters or implanted ports) [8].

Long-term artificial nutrition needs to be performed at home or in an institution, not in a hospital. Patients who receive geriatric care at home, including nutritional support, require considerable assistance from family members. Appropriateness

* Corresponding author. Tel.: +33 49 203 6018; fax: +33 49 203 6575.

E-mail address: stephane.schneider@unice.fr (S. M. Schneider).

of artificial nutrition should be considered with caution, taking into account the patient's particular circumstances, such as probable survival, rehabilitation potential, and complication risks [2]. When EN is to be delivered for long periods of time (i.e., prolonged dysphagia) the patient and his or her family members must be trained for home management. Furthermore, HEN and HPN patients and their family members should receive written instructions for dealing with common problems, as well as for routine procedures [9]. This is one of the tasks that can be handled at specialized artificial nutrition centers, where patients receiving long-term nutritional support may ideally be referred.

Outcome of artificial nutrition

The effectiveness of artificial nutrition on nutritional status may be compromised by poor compliance with the tubes (for EN) and by side effects. Many patients are bedridden and consequent immobility further enhances muscle wasting and prevents lean body mass gain. Weighing the patient also may prove to be problematic.

Survival, rehabilitation, and ability to resume oral intake are lower among older patients on home artificial nutrition compared with middle-aged and young patients [6,7,10]. A United Kingdom survey of 188 patients (963 patient-years of HPN) over a 25-y period found a lower survival probability among older patients compared with a younger population. However, HPN dependence among survivors was not affected by age [11]. In contrast, probably due to age-induced anorexia, patients will probably be more difficult to wean from EN. In a retrospective study of 102 patients post-chemoradiation for head and neck cancer and placement of a percutaneous endoscopic gastrostomy, EN was significantly ($P = 0.001$) prolonged as the age of patients increased [12].

Complications of HEN and HPN are similar to those observed in other age groups. They include tube complications (obstruction, removal) and diarrhea/constipation for enterally fed patients and catheter-related complications (infections, thrombosis, obstruction) for parenterally fed patients. Specifically, advanced age has been associated with a higher risk for

central catheter vascular erosion [13], but not for bloodstream infections [14].

References

- [1] Bozzetti F, Cozzaglio L, Biganzoli E, Chiavenna G, De Cicco M, Donati D, et al. Quality of life and length of survival in advanced cancer patients on home parenteral nutrition. *Clin Nutr* 2002;21:281–8.
- [2] Sobotka L, Schneider SM, Berner YN, Cederholm T, Krznaric Z, Shenkin A, et al. ESPEN Guidelines on Parenteral Nutrition: Geriatrics. *Clin Nutr* 2009;28:461–6.
- [3] Cozzaglio L, Balzola F, Cosentino F, DeCicco M, Fellagara P, Gaggiotti G, et al. Outcome of cancer patients receiving home parenteral nutrition. Italian Society of Parenteral and Enteral Nutrition (S.I.N.P.E.). *JPEN J Parenter Enteral Nutr* 1997;21:339–42.
- [4] Pasanisi F, Orban A, Scalfi L, Alfonsi L, Santarpia L, Zurlo E, et al. Predictors of survival in terminal-cancer patients with irreversible bowel obstruction receiving home parenteral nutrition. *Nutrition* 2001;17:581–4.
- [5] Bakker H, Bozzetti F, Staun M, Leon-Sanz M, Hebutterne X, Pertkiewicz M, et al. Home parenteral nutrition in adults: a European multicentre survey in 1997. ESPEN-Home Artificial Nutrition Working Group. *Clin Nutr* 1999;18:135–40.
- [6] Howard L, Malone M. Clinical outcome of geriatric patients in the United States receiving home parenteral and enteral nutrition. *Am J Clin Nutr* 1997;66:1364–70.
- [7] Paccagnella A, Baruffi C, Pizzolato D, Favaro V, Marcon ML, Morello M, et al. Home enteral nutrition in adults: a five-year (2001–2005) epidemiologic analysis. *Clin Nutr* 2008;27:378–85.
- [8] Staun M, Pironi L, Bozzetti F, Baxter J, Forbes A, Joly F, et al. ESPEN Guidelines on Parenteral Nutrition: home parenteral nutrition (HPN) in adult patients. *Clin Nutr* 2009;28:467–79.
- [9] De Cicco M, Bortolussi R, Fantin D, Matovic M, Fracasso A, Fabiani F, et al. Supportive therapy of elderly cancer patients. *Crit Rev Oncol Hematol* 2002;42:189–211.
- [10] Schneider SM, Raina C, Pugliese P, Pouget I, Rampal P, Hebutterne X. Outcome of patients treated with home enteral nutrition. *JPEN J Parenter Enteral Nutr* 2001;25:203–9.
- [11] Lloyd DA, Vega R, Bassett P, Forbes A, Gabe SM. Survival and dependence on home parenteral nutrition: experience over a 25-year period in a UK referral centre. *Aliment Pharmacol Ther* 2006;24:1231–40.
- [12] Lawson JD, Gaultney J, Saba N, Grist W, Davis L, Johnstone PA. Percutaneous feeding tubes in patients with head and neck cancer: rethinking prophylactic placement for patients undergoing chemoradiation. *Am J Otol* 2009;30:244–9.
- [13] Walshe C, Phelan D, Bourke J, Buggy D. Vascular erosion by central venous catheters used for total parenteral nutrition. *Intensive Care Med* 2007;33:534–7.
- [14] Bozzetti F, Mariani L, Bertinet DB, Chiavenna G, Crose N, De Cicco M, et al. Central venous catheter complications in 447 patients on home parenteral nutrition: an analysis of over 100,000 catheter days. *Clin Nutr* 2002;21:475–85.