

# Postpartum Haemorrhage

Tutorial

Week 3 · Medical disorders &amp; PPH

Obstetrics

Dr Kavuma

Primary PPH is blood loss of 500 mL or more within 24 hours of birth (or 300 mL or more with abnormal vital signs, WHO 2025); it is the leading killer of Ugandan mothers, and survival depends on measuring blood loss objectively and starting the E-MOTIVE bundle within minutes, not on estimating by eye.

## Must know before the ward round

1. Primary PPH:  $\geq 500$  mL within 24 h of birth; WHO/FIGO/ICM 2025 also say act at  $\geq 300$  mL plus any abnormal vital sign. Severe/major PPH:  $\geq 1000$  mL. Secondary PPH: 24 h to 12 weeks.
2. Causes = the 4 Ts: Tone (about 70%), Trauma (about 20%), Tissue (about 10%), Thrombin (under 1%). Always check all four.
3. Prevention (AMTSL): oxytocin 10 IU IM/IV within 1 minute of birth, delayed cord clamping, controlled cord traction, then check uterine tone every 15 minutes for 2 hours.
4. E-MOTIVE bundle, started together as soon as PPH is diagnosed: uterine Massage, Oxytocics, Tranexamic acid, IV fluids, Vaginal/genital-tract Examination, then Escalate if bleeding continues.
5. Treatment oxytocin: 10 IU IV slowly (or IM), then 20 IU in 1 L NS/RL at 60 drops/min (maintenance 40 drops/min); no more than 3 L of oxytocin-containing fluid.
6. Tranexamic acid 1 g IV over 10 minutes within 3 hours of birth; a second 1 g if bleeding continues after 30 minutes or restarts within 24 hours (WOMAN trial). No benefit after 3 hours.
7. Second-line uterotonics: misoprostol 800  $\mu$ g sublingual; ergometrine 0.2 mg IM (max 1 mg/24 h; not in hypertension or heart disease); carboprost 0.25 mg deep IM (max 2 mg; not in asthma).
8. Calibrated drape: E-MOTIVE (NEJM 2023) cut severe PPH, laparotomy or death from bleeding by 60% (1.6% vs 4.3%).
9. Visual estimation underestimates blood loss by 30–50%; weigh pads (1 g  $\approx$  1 mL). Shock index (HR  $\div$  SBP)  $\geq 0.9$  is abnormal and  $\geq 1.4$  needs urgent action; hypotension is a late sign.
10. Firm uterus but still bleeding = trauma (inspect cervix and vagina in good light) or thrombin (bedside clotting test: no clot by 7 min), not a reason for another uterotonic.
11. Retained placenta: not out 30 minutes after birth  $\rightarrow$  empty the bladder, extra oxytocin 10 IU, repeat CCT, then manual removal with a single-dose antibiotic. Never ergometrine.
12. Shock out of proportion to visible loss + fundus not palpable = uterine inversion: stop oxytocin, replace immediately, then give oxytocin.

- |                                 |                             |
|---------------------------------|-----------------------------|
| 1 In a nutshell                 | 8 Investigations            |
| 2 Definitions & classification  | 9 Management                |
| 3 Epidemiology & Uganda context | 10 Complications            |
| 4 Physiology & pathophysiology  | 11 Ward-round & exam pearls |
| 5 Clinical acumen               | 12 Mnemonics & memory aids  |
| 6 Clinical workup               | 13 High-yield summary       |
| 7 Differential diagnosis        |                             |

# In a nutshell

**Postpartum haemorrhage (PPH)** is excessive bleeding from the genital tract after birth. It is the **number-one direct cause of maternal death in Uganda** and worldwide. Most of these deaths happen **within 24 hours of birth, often within the first 2 hours, and most are preventable**.

The modern approach, in the **WHO 2023 recommendations** and the **WHO/FIGO/ICM 2025 consolidated PPH guideline**, rests on two ideas:

1. **Detect early:** *measure* blood loss (calibrated drape) instead of guessing, and act at **300 mL if vital signs are abnormal**.
2. **Treat as a bundle:** give the **E-MOTIVE** first-response bundle all at once and quickly, not one drug at a time waiting to see what works.

## CLINICAL PEARL

The one thing never to forget: **a woman who bleeds after delivery can die within 2 hours**. Call for help early, measure, massage, give oxytocin **and** tranexamic acid, run fluids, look for the cause (4 Ts), and escalate. Don't wait for her blood pressure to fall.

## Definitions & classification

| Term                       | Definition                                                                                                                                                                                                                             |
|----------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Primary PPH</b>         | Blood loss <b>≥500 mL</b> from the genital tract within <b>24 hours</b> of birth (any route). The WHO/FIGO/ICM <b>2025</b> guideline adds: diagnose PPH at <b>≥300 mL plus any abnormal vital sign</b> (e.g. tachycardia, hypotension) |
| <b>Severe / major PPH</b>  | <b>≥1000 mL</b> (RCOG: moderate 1000–2000 mL; severe >2000 mL). Minor = 500–1000 mL                                                                                                                                                    |
| <b>PPH after caesarean</b> | Classically <b>≥1000 mL</b> ; but treat <b>any bleeding that threatens haemodynamic stability</b> as PPH                                                                                                                               |
| <b>Clinical definition</b> | Any blood loss enough to cause haemodynamic instability. A small, anaemic woman (Hb 8 g/dL) may collapse after "only" 400 mL                                                                                                           |
| <b>Secondary PPH</b>       | Abnormal or excessive bleeding from <b>24 hours to 12 weeks</b> after birth                                                                                                                                                            |
| <b>Refractory PPH</b>      | Bleeding that continues despite first-line uterotonics and the first-response bundle                                                                                                                                                   |
| <b>Retained placenta</b>   | Placenta not delivered within <b>30 minutes</b> of birth with active management (WHO; UK uses 60 minutes)                                                                                                                              |

## The 4 Ts (cause-based classification)

| T                   | Approx. share | Examples                                                                                                                                                                                                                   |
|---------------------|---------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Tone</b> (atony) | ~70%          | Overdistension (twins, polyhydramnios, macrosomia), prolonged or precipitate labour, prolonged oxytocin augmentation, chorioamnionitis, grand multiparity, fibroids, MgSO <sub>4</sub> , general anaesthesia, full bladder |
| <b>Trauma</b>       | ~20%          | Perineal, vaginal and cervical tears, episiotomy extension, uterine rupture, uterine inversion, broad-ligament haematoma, CS extensions                                                                                    |
| <b>Tissue</b>       | ~10%          | Retained placenta, retained cotyledon or membranes, placenta accreta spectrum (PAS), clots in the cavity                                                                                                                   |

| T        | Approx. share | Examples                                                                                                                                                                                 |
|----------|---------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Thrombin | <1%           | DIC (abruption, amniotic fluid embolism, sepsis, IUFD, severe pre-eclampsia/HELLP), anticoagulants, inherited bleeding disorders, <b>dilutional coagulopathy after large transfusion</b> |

## Epidemiology & Uganda context

- Globally PPH affects millions of women every year. It is the single largest cause of maternal death, most of it in sub-Saharan Africa and South Asia.
- Uganda:** the MMR fell from **336 per 100,000 live births (UDHS 2016) to 189 (UDHS 2022)**. That is still far above the SDG target of <70. **Haemorrhage remains the leading direct cause** of maternal death in MPDSR reports, together with hypertensive disorders, sepsis and abortion complications.
- Ugandan studies report PPH ( $\geq 500$  mL) in about **9% of births**, with severe PPH ( $\geq 1000$  mL) in about 1%. That is probably an underestimate, because blood loss is estimated by eye.

### UGANDA CONTEXT

The realities on a Ugandan labour ward matter. **Blood** is often unavailable at HC IVs and even at regional referral hospitals (RRHs), where it comes from the Uganda Blood Transfusion Service. **Oxytocin quality** suffers when the cold chain fails (hence interest in **heat-stable carbetocin**). Many women are **anaemic** before they bleed. Night-time deliveries are covered by one midwife. Referral from HC III to HC IV to RRH can take hours. Your management must work with what is on the shelf: oxytocin, misoprostol, ergometrine, TXA, IV fluids, a Foley catheter and a condom, and your hands.

**Risk factors** (but **two-thirds of PPH happens in women with no risk factor**, so prepare for every birth):

| Antenatal                                                                                                                                                                                                                        | Intrapartum                                                                                                                                                                                                           |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Previous PPH (strongest), grand multiparity ( $\geq 5$ ), multiple pregnancy, polyhydramnios, macrosomia, anaemia (Hb <9 g/dL), placenta praevia/PAS, abruption, pre-eclampsia, fibroids, obesity, maternal age >35, previous CS | Prolonged labour (1st or 2nd stage), prolonged oxytocin use, induction, chorioamnionitis/fever, instrumental birth, CS (especially emergency), episiotomy, retained placenta, general anaesthesia, precipitate labour |

## Physiology & pathophysiology

### Why the placental bed bleeds so much

- At term, **uterine blood flow is 500–800 mL/min** (about 10–15% of cardiac output). The spiral arteries have been converted by trophoblast into **wide, low-resistance, non-muscular channels**. They **cannot constrict themselves**.
- After the placenta separates, bleeding is stopped mainly by **mechanical compression**. The criss-cross (interlacing) fibres of the myometrium contract and kink the vessels running between them. These are the **"living ligatures"**, and they explain why **uterine tone is everything**. Clotting and local thrombosis are secondary.
- So:** a soft, "boggy" uterus = open vessels = torrential bleeding (tone). A well-contracted uterus that still bleeds must be bleeding from **somewhere else** (trauma or thrombin), or something is preventing contraction (**tissue:** a retained cotyledon holds the uterus open).

### How the body compensates, and why that fools us

- Pregnancy increases blood volume by about **40–50%** (roughly 1.5 L extra). A healthy woman at term can lose **1000 mL with only mild tachycardia**. Young women hold their blood pressure through vasoconstriction

until very late.

- **Hypotension is a late sign** (often after 30–40% of blood volume is lost). **Tachycardia, pallor, cool peripheries, restlessness, reduced urine output and a rising shock index** come first.
- **Shock index (SI) = heart rate ÷ systolic BP**. Normal postpartum SI is 0.7–0.9. **SI ≥ 0.9** should prompt concern and **SI ≥ 1.4** means urgent action and transfer.

## The lethal triad

Continued bleeding leads to **hypothermia** (cold fluids, exposure), **acidosis** (hypoperfusion) and **coagulopathy** (consumption plus dilution with crystalloid). Each worsens the others. Fibrinogen falls first in obstetric haemorrhage, and **fibrinogen < 2 g/L predicts progression to severe PPH**. This is why we warm fluids, limit crystalloid, give **tranexamic acid early** (to block fibrinolysis before clots dissolve), and transfuse blood and clotting factors rather than litres of saline.

## Why tranexamic acid must be early

TXA blocks plasminogen activation (antifibrinolytic). Fibrinolysis rises sharply within minutes to hours of birth and bleeding. In the **WOMAN trial** (20,060 women), TXA **reduced death from bleeding by about 30% when given within 3 hours**. Given later there was no benefit. Every 15 minutes of delay reduces its benefit.

# Clinical acumen

## Focused history

In the acute setting, history is taken **while** resuscitation happens, often from the notes and the midwife.

| Question                                                                  | Why it matters                                                                                                           |
|---------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------|
| When and how did she deliver? (SVD, vacuum, CS; time of birth)            | Instrumental birth → trauma; CS → intra-abdominal bleeding; time since birth tells you whether TXA can still help (<3 h) |
| Was AMTSL done? Which uterotonic, when? Is the placenta out and complete? | No oxytocin → atony likely; incomplete placenta → tissue                                                                 |
| Length of labour, oxytocin augmentation, fever in labour                  | Prolonged/augmented labour and chorioamnionitis → atony                                                                  |
| Baby's weight, twins, polyhydramnios                                      | Overdistension → atony                                                                                                   |
| Parity; previous PPH, previous CS or uterine surgery                      | Grand multipara → atony and rupture; previous CS + praevia → accreta                                                     |
| Antenatal Hb, HIV status, blood group                                     | Anaemic women decompensate early; plan transfusion                                                                       |
| Pre-eclampsia? BP in labour?                                              | <b>Contraindicates ergometrine</b> ; HELLP → thrombin                                                                    |
| Asthma? Heart disease?                                                    | <b>Carboprost contraindicated in asthma</b> ; ergometrine avoided in cardiac disease                                     |
| Bleeding disorders, anticoagulants, abruption, IUFD                       | Thrombin cause                                                                                                           |
| Symptoms: dizziness, thirst, breathlessness, confusion                    | Hypovolaemia; confusion = late, severe shock                                                                             |

## Examination

1. **General**: conscious level, pallor, sweating, cold clammy peripheries, capillary refill >2 s, restlessness.
2. **Vital signs**: pulse (rate and volume), BP, respiratory rate, SpO<sub>2</sub>, temperature, **shock index**, **urine output** (catheter; target >30 mL/h).

3. **Abdomen: feel the fundus.** Is it **soft and above the umbilicus** (atony) or firm and central (look elsewhere)?  
A deviated fundus suggests a full bladder. Tenderness and peritonism after CS or obstructed labour suggest rupture or intra-abdominal bleeding. **Absent fundus on palpation plus shock out of proportion to bleeding = uterine inversion.**
4. **Blood loss:** measure it. Read the **calibrated drape**, weigh pads and linen (1 g  $\approx$  1 mL), count soaked swabs. Also look under the woman and on the floor.
5. **Genital tract:** in good light with a speculum and an assistant. Look at perineum, vagina, cervix (all quadrants), and for a vulval or vaginal haematoma.
6. **Placenta and membranes:** complete? Missing cotyledon? Vessels running off the edge (succenturiate lobe)?
7. **Clotting:** does the blood clot? **Bedside clotting test** (WHO): 2 mL in a small dry glass tube held warm in your fist; tilt at 4 min, then every minute; **no clot by 7 minutes**, or a soft clot that breaks up, = coagulopathy. (Some units use the 20-minute whole-blood clotting test instead.) Oozing from drip sites also suggests coagulopathy.

## Red flags

### RED FLAG

- Pulse >100–110, SBP <90–100 mmHg, **shock index  $\geq 0.9$**  ( $\geq 1.4$  = critical)
- Restlessness, confusion, air hunger, oliguria (<30 mL/h)
- Bleeding continuing after the first uterotonic, or a uterus that keeps relaxing
- **Shock out of proportion to visible bleeding:** think concealed bleeding (uterine rupture, broad-ligament or vulval haematoma, intra-abdominal bleeding after CS) or **uterine inversion**
- Blood that doesn't clot, oozing from cannula sites: DIC
- Severe perineal or rectal pain after delivery: expanding haematoma

## Thinking like a clinician

- **"Is the uterus hard?"** is the first branching question. Soft  $\rightarrow$  Tone (massage, oxytocics, empty bladder, check for tissue). Hard but still bleeding  $\rightarrow$  Trauma (inspect genital tract) or Thrombin (clotting test).
- **Bright red trickle with a well-contracted uterus** = a tear until proven otherwise, usually cervical or high vaginal. Clamp and suture; a second uterotonic won't fix it.
- **Uterus contracts then relaxes again and again**  $\rightarrow$  think **retained tissue** (placenta incomplete) or overdistension/infection. Explore the uterus.
- **The fundus can't be felt, the woman is in profound shock (partly neurogenic, with bradycardia) and there is a mass in the vagina**  $\rightarrow$  **uterine inversion**. Replace it immediately, before the cervix clamps down. **Stop the oxytocin until it is replaced.**
- **Previous CS + prolonged labour + sudden collapse**  $\rightarrow$  uterine rupture: laparotomy, not another uterotonic.
- **After CS:** rising pulse, abdominal distension, falling Hb with little vaginal loss  $\rightarrow$  intra-abdominal bleeding; return to theatre.
- **Anaemic woman:** her "500 mL" is someone else's 1500 mL. Treat the woman, not the number.
- **Common student mistakes:** underestimating loss by eye; waiting for hypotension; giving drugs one at a time; forgetting to catheterise; forgetting TXA; giving ergometrine to a hypertensive woman; forgetting to examine the placenta and the cervix; not documenting times.

### EXAM FAVOURITE

Examiners love this: "**The uterus is well contracted but she is still bleeding. What will you do?**" Answer: inspect the genital tract under good light for trauma (cervical and vaginal tears, haematoma), check the placenta for completeness, do a bedside clotting test, and continue resuscitation.

## Clinical workup

In PPH the workup and the treatment happen **at the same time**. You never finish the history before you touch the uterus. Think of the steps below as what the team leader is checking off while the bundle runs.

### Step 0: First 5 minutes

1. **Recognise and name it.** Say out loud: "This is PPH. Time is 10:05." Triggers: calibrated drape  $\geq 500$  mL, or  $\geq 300$  mL plus any abnormal vital sign (WHO/FIGO/ICM 2025), or any bleeding with a woman who looks unwell.
2. **Call for help:** second midwife, the most senior clinician, anaesthetic officer; alert theatre and the blood bank. Assign roles (uterus, IV and drugs, observations, scribe/timekeeper).
3. **Hand on the fundus at once.** Soft and high  $\rightarrow$  start **massage** now. This is both the first examination and the first treatment.
4. **ABC and vitals:** airway and conscious level, RR, SpO<sub>2</sub>, pulse, BP, temperature. Calculate the **shock index** (HR  $\div$  SBP):  $\geq 0.9$  abnormal,  $\geq 1.4$  urgent,  $\geq 1.7$  critical.
5. **Position and oxygen:** lie flat (legs raised if shocked), oxygen 6–8 L/min by mask, keep her warm.
6. **Two large-bore cannulae (14–16G).** From the first cannula draw blood for **Hb, group and cross-match**, and fill a **dry glass tube for the bedside clotting test**.
7. **Start the bundle together:** oxytocin 10 IU IV slowly then infusion, **TXA 1 g IV over 10 min** (if within 3 h of birth), warmed crystalloid, catheterise the bladder.

### RED FLAG

Call the most senior person available **immediately** if: shock index  $\geq 1.4$ , SBP  $< 90$  mmHg, confusion or drowsiness, loss  $\geq 1000$  mL, blood that does not clot, or shock out of proportion to what you can see (think rupture, inversion, concealed haematoma, intra-abdominal bleeding after CS).

### Step 1: Focused history

Take it from the midwife, the partograph and the file while others treat. Each item changes the plan:

- ☐ **Time of birth**  $\rightarrow$  still within 3 h? (TXA works only then; also tells you whether this is primary or secondary PPH.)
- ☐ **Mode of birth:** SVD, vacuum, CS, fundal pressure  $\rightarrow$  instrumental = tear; CS = intra-abdominal bleeding; fundal pressure/cord traction on an unseparated placenta = inversion.
- ☐ **Was AMTSL given? Which drug, when?**  $\rightarrow$  no prophylactic oxytocin makes atony more likely; you also need the total uterotonic dose so far.
- ☐ **Placenta out? Complete?**  $\rightarrow$  retained or incomplete placenta = tissue; plan MRP.
- ☐ **Labour:** length, oxytocin augmentation, fever or chorioamnionitis, precipitate birth  $\rightarrow$  atony; fever  $\rightarrow$  sepsis-related atony and DIC.
- ☐ **Baby:** weight, twins, polyhydramnios, stillbirth  $\rightarrow$  overdistension; IUFD or abruption  $\rightarrow$  DIC.
- ☐ **Parity and uterine history:** grand multipara, previous PPH, previous CS/myomectomy  $\rightarrow$  atony, rupture, accreta.
- ☐ **Blood pressure / pre-eclampsia, heart disease**  $\rightarrow$  no ergometrine; HELLP  $\rightarrow$  low platelets.

- ☐ **Asthma → no carboprost.**
- ☐ **Antenatal Hb, blood group, HIV status, known bleeding disorder or anticoagulant** → how little reserve she has; Rh status for anti-D; thrombin cause.
- ☐ **Symptoms:** dizziness, thirst, breathlessness, severe perineal/rectal pain (haematoma), abdominal pain (rupture), confusion (late shock).

## Step 2: Focused examination

**General:** conscious level (AVPU/GCS), pallor, sweating, cold peripheries, capillary refill (>2 s abnormal), air hunger.

**Vital signs** every 5–15 min: pulse (rate, volume), BP, RR, SpO<sub>2</sub>, temperature, shock index, **urine output** via catheter (target  $\geq 30$  mL/h).

### Abdomen:

- **Fundus:** height, consistency (soft/boggy vs firm), central or deviated (full bladder), does it relax again after massage (tissue)?
- Tenderness, guarding, distension, free fluid, easily felt fetal parts or a scar (rupture; post-CS bleeding).
- **Fundus not palpable** → uterine inversion.

### Genital tract (in good light, with an assistant and a speculum or retractors):

- Perineum and episiotomy; vaginal walls and fornices; **all four quadrants of the cervix** (walk sponge forceps around it).
- Vulval or paravaginal swelling, severe pain, bluish tense mass = haematoma; rectal exam if a haematoma is suspected.
- A mass in the vagina or at the introitus = inversion (or a fibroid polyp, much rarer).

**Placenta and membranes:** lay the placenta out; missing cotyledons, ragged membranes, vessels running to a torn edge (succenturiate lobe).

**Signs of coagulopathy:** oozing from cannula sites, the episiotomy or gums, petechiae, blood in the drape that does not clot.

| Document these positives                          | And these negatives                   |
|---------------------------------------------------|---------------------------------------|
| Measured loss (drape + weighed pads) and the time | "Placenta and membranes complete"     |
| Fundal height and tone after massage              | "Cervix inspected all round, no tear" |
| Shock index and urine output                      | "No haematoma, no inversion"          |
| Site and degree of any tear                       | "Blood clots in the 20-min test"      |

## Step 3: Bedside tests

| Test                               | How                                                                                 | What it tells you                                                                     |
|------------------------------------|-------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------|
| <b>Calibrated drape reading</b>    | Read the level every 5–15 min; note the time                                        | Objective loss; the trigger for the bundle                                            |
| <b>Weigh pads, linen and swabs</b> | Weigh soaked items and subtract the weight of the same items dry; <b>1 g ≈ 1 mL</b> | Adds the loss the drape missed (after the drape is removed, on the floor, in the bed) |
| <b>Shock index</b>                 | HR ÷ SBP, repeated with each set of vitals                                          | Rising SI = ongoing bleeding even if BP is "normal"                                   |

| Test                                                   | How                                                                                                                                              | What it tells you                                                             |
|--------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------|
| <b>HemoCue / point-of-care Hb</b>                      | Finger-prick or venous sample                                                                                                                    | Baseline only; the <b>first Hb underestimates the loss</b> (see Step 5)       |
| <b>Bedside clotting test</b>                           | 2 mL venous blood in a clean, dry <b>glass</b> tube; hold it warm, tilt at 4 min then every minute (WHO MCPC); some units read at 20 min instead | No clot by 7 min, or a clot that breaks up = coagulopathy (DIC)               |
| <b>Hourly urine output</b>                             | Foley catheter to a urometer or measured bag                                                                                                     | <30 mL/h = under-resuscitated or AKI                                          |
| <b>Point-of-care ultrasound</b> (if skilled)           | Uterine cavity, abdomen                                                                                                                          | Retained products/clots in the cavity; free fluid (rupture, post-CS bleeding) |
| <b>Urine dipstick for protein, BP</b>                  |                                                                                                                                                  | Undiagnosed pre-eclampsia → avoid ergometrine; think HELLP                    |
| <b>Malaria RDT, temperature</b> (secondary PPH, fever) |                                                                                                                                                  | Co-existing malaria, sepsis                                                   |

## Step 4: Laboratory & imaging

| Test                                  | When                                                                                | Expected / abnormal finding                                                                                     | How it changes management                                                                                                                                                                               |
|---------------------------------------|-------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Group and cross-match</b>          | Every PPH, with the first cannula                                                   | Blood group; 2–4 compatible units                                                                               | Transfuse early in major PPH. <b>HC IV:</b> only where there is a blood fridge and a supply from the regional blood bank (UBTS); otherwise refer. <b>RRH/NRH:</b> usually available, but stocks run out |
| <b>Full blood count</b>               | Baseline, then after resuscitation and at 24 h                                      | Hb (baseline often low from antenatal anaemia); platelets $<75 \times 10^9/L$ abnormal                          | Transfusion and platelet decisions; Hb at 24 h guides iron vs transfusion. Available at most HC IVs and hospitals                                                                                       |
| <b>PT/INR, APTT</b>                   | Major PPH ( $\geq 1000$ mL), abnormal clotting test, abruption, IUFD, sepsis, HELLP | Prolonged ( $>1.5 \times$ normal)                                                                               | FFP 15 mL/kg. <b>Mainly RRH/NRH/private hospitals;</b> often not at night                                                                                                                               |
| <b>Fibrinogen</b>                     | As above                                                                            | <b><math>&lt;2</math> g/L = severe coagulopathy</b> and predicts progression (normal in late pregnancy 4–6 g/L) | Cryoprecipitate (or fresh whole blood if no components). <b>NRH and some private hospitals only</b>                                                                                                     |
| <b>Urea, creatinine, electrolytes</b> | Major PPH, oliguria                                                                 | Creatinine rising above $\sim 80\text{--}90 \mu\text{mol/L}$ ; high $K^+$                                       | AKI: fluid balance, avoid nephrotoxins (gentamicin dose), renal review. RRH/NRH                                                                                                                         |
| <b>LFTs</b>                           | Hypertensive woman, low platelets                                                   | Raised ALT/AST (with low platelets, haemolysis)                                                                 | HELLP → coagulopathy, $\text{MgSO}_4$ , no ergometrine                                                                                                                                                  |
| <b>Lactate / blood gas</b>            | Severe shock, HDU/ICU                                                               | Lactate $>4$ mmol/L, base deficit                                                                               | Severity of shock; response to resuscitation. NRH/ICU only                                                                                                                                              |

| Test                       | When                                              | Expected / abnormal finding                         | How it changes management                                                                         |
|----------------------------|---------------------------------------------------|-----------------------------------------------------|---------------------------------------------------------------------------------------------------|
| <b>Ultrasound (formal)</b> | Retained tissue suspected; secondary PPH; post-CS | Echogenic mass in the cavity; free fluid; haematoma | Evacuation vs laparotomy vs expectant. Most HC IVs and all hospitals, if a sonographer is on site |
| <b>Blood cultures, HVS</b> | Secondary PPH or fever                            | Organisms and sensitivities                         | Tailor antibiotics. RRH/NRH (culture capacity limited)                                            |
| <b>β-hCG</b>               | Secondary PPH that persists after evacuation      | Raised or plateauing                                | Suspect gestational trophoblastic neoplasia; send tissue for histology                            |

## Step 5: Putting it together

### Model one-line summary

"A 32-year-old para 5, 40 minutes after SVD of a 4.1 kg baby following an augmented labour, with primary PPH of 900 mL (drape plus pads), shock index 1.2, a soft uterus that firms with massage, a complete placenta and a second-degree tear: atonic PPH with a minor trauma component, haemodynamically compensated."

### Problem list

1. Primary PPH (900 mL and still being measured), cause mainly **Tone**, plus **Trauma** (tear).
2. Hypovolaemia (SI 1.2), on the bundle.
3. Pre-existing anaemia (antenatal Hb 9.5 g/dL) → low reserve.
4. Risk of coagulopathy (continue clotting tests if loss exceeds 1000 mL).

**Working diagnosis:** *primary atonic PPH with a second-degree perineal tear, compensated hypovolaemic shock.*

#### EXAM FAVOURITE

**Worked example 1: shock index and blood-loss estimation.** The drape reads **650 mL** when it is removed. After that, four soaked pads and a draw sheet weigh **620 g**; the same items dry weigh **120 g**. Extra loss =  $620 - 120 = 500 \text{ g} \approx 500 \text{ mL}$ . **Total  $\approx 1150 \text{ mL}$  = severe PPH.** Pulse 126, BP 90/60 → **SI =  $126 \div 90 = 1.4$**  → urgent: second cannula, blood now, senior at the bedside, prepare theatre. The eye estimate in the notes said "about 500 mL": a classic underestimate. Her admission BP was 120/80 with pulse 86: SBP has fallen by 30 mmHg and pulse has risen by 40. By the **Rule of 30** (quick reference) she has lost roughly **30% of her blood volume** (moderate shock), which fits the weighed total.

#### EXAM FAVOURITE

**Worked example 2: the first Hb and the clotting test.** Thirty minutes after a 1500 mL PPH following abruption with IUFD, her Hb is **9.8 g/dL** (antenatal 11.2). This does **not** mean the loss was small: she has lost whole blood and there has been no time for haemodilution, so the Hb falls further over hours (and with crystalloid). Treat the **clinical picture and the measured loss**, not the first Hb. The bedside clotting test shows **no clot at 7 minutes**. Later labs: platelets  **$62 \times 10^9/\text{L}$** , PT prolonged, fibrinogen **1.3 g/L**. Interpretation: **DIC** from abruption plus consumption. Action: TXA (if within 3 h), red cells plus **FFP 15 mL/kg, cryoprecipitate** (fibrinogen  $<2 \text{ g/L}$ ), **platelets** ( $<75 \times 10^9/\text{L}$ ); fresh whole blood if components are not available; keep her warm; ensure the uterus is empty and contracted.

### UGANDA CONTEXT

At most HC IVs you will have the **drape, a scale, a HemoCue, a glass tube and your hands**, and often no coagulation lab. That is enough to diagnose PPH, grade shock, find the cause (4 Ts) and detect coagulopathy. What you often **cannot** do is transfuse components or operate at night, so the workup must end in a clear decision: **manage here or refer now** (with NASG, tamponade, oxytocin running and an escort).

## Differential diagnosis

These are differentials for the **cause** of bleeding (and for shock after delivery).

| Condition                               | Distinguishing features                                                                                  | Key investigation / action                                    |
|-----------------------------------------|----------------------------------------------------------------------------------------------------------|---------------------------------------------------------------|
| Uterine atony                           | Soft, boggy, large uterus above umbilicus; bleeding with clots; improves with massage                    | Palpation; response to massage/oxytocin                       |
| Genital-tract trauma                    | Uterus firm; continuous bright-red bleeding; instrumental or precipitate birth                           | Speculum exam in good light (±EUA)                            |
| Retained placenta / tissue              | Placenta not delivered or incomplete; uterus relaxes repeatedly                                          | Inspect placenta; manual exploration; ultrasound if available |
| Uterine rupture                         | Previous CS/obstructed labour; abdominal pain, shock > visible loss, fetal parts easily felt, haematuria | Clinical; laparotomy                                          |
| Uterine inversion                       | Fundus not palpable abdominally; mass at introitus/in vagina; severe pain; shock with bradycardia        | Clinical; immediate replacement                               |
| Coagulopathy / DIC                      | Non-clotting blood, oozing from sites; abruption, IUFD, AFE, sepsis, HELLP                               | 20-min WBCT, platelets, PT/APTT, fibrinogen                   |
| Vulval/vaginal/broad-ligament haematoma | Severe pain, rectal pressure, unexplained tachycardia, swelling; little external blood                   | Examination; ultrasound; drainage                             |
| Amniotic fluid embolism                 | Sudden collapse, hypoxia, hypotension, seizure, then DIC                                                 | Clinical diagnosis; ICU                                       |
| Other shock (sepsis, PE, cardiac)       | Fever, chest pain, dyspnoea; uterus fine and little bleeding                                             | Clinical picture; ECG, etc.                                   |

## Investigations

The full test-by-test table (with Ugandan availability) is in **Clinical workup, Step 4** above. Three principles to remember:

1. **Send bloods while you resuscitate**; never delay treatment for a result. The first cannula gives you Hb, group and cross-match and the clotting-test tube.
2. **Bedside tests beat the lab at night**: the calibrated drape, weighed pads, shock index, HemoCue, the bedside clotting test and hourly urine output can all be done at a HC IV.
3. **Repeat, don't rely on one value**: the first Hb underestimates loss, and a normal first clotting test can become abnormal after a further litre of bleeding. Repeat Hb after resuscitation and at 24 h, and repeat clotting tests if loss exceeds about 1000 mL or bleeding continues.

# Management

## A. Prevention: at every birth

**Antenatal:** treat anaemia (iron + folic acid; Hb target  $\geq 11$  g/dL), identify high-risk women (previous PPH, praevia, grand multipara, twins), **birth preparedness** (deliver in a CEmONC facility, identify blood donors, transport money).

**Active management of the third stage of labour (AMTSL)**, per WHO:

1. **Exclude a second baby** (palpate the abdomen).
2. **Oxytocin 10 IU IM** (or IV slowly) **within 1 minute of birth**. This is the **most important component**.
3. **Delayed cord clamping** (1–3 minutes) unless the baby needs resuscitation.
4. **Controlled cord traction (CCT)** with counter-traction above the pubis, only during a contraction, by a skilled attendant. It is optional where attendants are not skilled.
5. **Assess uterine tone** abdominally **every 15 minutes for 2 hours** (sustained continuous massage is no longer recommended if oxytocin was given, but tone checks are).
6. **Examine the placenta and membranes** for completeness, and the genital tract for tears.
7. **Measure blood loss objectively** (calibrated drape placed under the buttocks after birth, WHO 2023).

| Prophylactic uterotonic            | Dose & route                       | Notes                                                                                                                                                    |
|------------------------------------|------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Oxytocin</b> (first choice)     | <b>10 IU IM or IV</b> (slow IV)    | Store 2–8 °C; loses potency in heat                                                                                                                      |
| <b>Heat-stable carbetocin</b>      | <b>100 µg IM or IV</b> single dose | Non-inferior to oxytocin (CHAMPION trial, which included Uganda). Use where oxytocin quality can't be assured. <b>For prevention only, not treatment</b> |
| <b>Misoprostol</b>                 | <b>400–600 µg orally</b>           | Where injectables are not available (home births, community distribution)                                                                                |
| Ergometrine / oxytocin-ergometrine | 0.2 mg IM / 5 IU + 0.5 mg IM       | Only if no hypertension; more side effects                                                                                                               |
| Tranexamic acid (at CS)            | —                                  | Not routine for all births. Some units give 1 g IV at CS for high-risk women. Follow local protocol                                                      |

### UGANDA CONTEXT

In Uganda, **oxytocin 10 IU IM** is standard AMTSL at every facility. **Misoprostol 600 µg orally** (3 × 200 µg tablets) has been distributed for women who may deliver at home, to be taken immediately after the baby is born, and never before. Heat-stable carbetocin has been introduced in some humanitarian settings. At caesarean, oxytocin is given IV **slowly** (a rapid bolus causes hypotension).

## B. Treatment: first response (the E-MOTIVE bundle)

**Diagnose early.** Calibrated drape reading  $\geq 500$  mL, or  $\geq 300$  mL **plus abnormal vital signs or clinical concern**, triggers the bundle. Start all components **together, within 15 minutes**.

**Emergency algorithm (vaginal birth, any facility):**

1. **Call for help.** Shout for the midwife and the most senior clinician; assign roles (one on airway/obs, one on IV and drugs, one on the uterus, one scribe/timekeeper). Note the time.
2. **ABC:** lie flat, raise legs; **oxygen** 6–8 L/min by mask; keep her warm.
3. **Two large-bore IV cannulae** (14–16G). Draw blood for Hb, group and cross-match, and a clotting test.

4. **M: uterine Massage.** Rub the fundus firmly to stimulate a contraction and expel clots. Continue until firm.
5. **O: Oxytocin** 10 IU IV slowly (or IM), then an infusion (see drug table).
6. **T: Tranexamic acid 1 g IV over 10 minutes** (within 3 hours of birth). Repeat 1 g after 30 minutes if bleeding continues.
7. **I: IV fluids:** warmed crystalloid (Ringer's lactate or normal saline). Give a 500 mL–1 L bolus quickly, then reassess. Avoid more than about 2 L of crystalloid before blood; start **blood** as soon as available.
8. **Empty the bladder** (Foley catheter, left in to measure urine output hourly).
9. **V: Vaginal/genital-tract Examination.** Is the placenta out and complete? Look for tears (repair them), haematoma or inversion.
10. **E: Escalate** if bleeding continues despite the bundle. Move to second-line drugs, bimanual compression, tamponade, theatre, and **refer early** from a HC III/IV that cannot operate or transfuse.
11. **Monitor** pulse, BP, RR, SpO<sub>2</sub> and blood loss every 15 minutes; urine hourly; **document** drug times and volumes.

#### NOTE

Why a bundle? In the **E-MOTIVE trial** (210,132 vaginal births, 78 hospitals in Kenya, Nigeria, South Africa and Tanzania), the calibrated drape plus the bundle **reduced the composite of severe PPH ( $\geq 1000$  mL), laparotomy for bleeding or maternal death from bleeding from 4.3% to 1.6% (RR 0.40).** Treatment was started early and together instead of sequentially after each "wait and see". WHO adopted it in 2023.

## C. Uterotonic and haemostatic drugs for treatment

#### DRUG DOSING

Confirm the dose on your unit protocol. These are the WHO / Uganda regimens.

| Drug                                             | Dose                                                                                                                     | Route                                             | Frequency / max                                                                                                             | Contraindications, side effects & notes                                                                                                                           |
|--------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Oxytocin</b> (first line)                     | <b>10 IU</b> , then infusion <b>20 IU in 1 L NS/RL at 60 drops/min</b> ; maintenance 20 IU in 1 L at <b>40 drops/min</b> | IV slowly or IM; infusion IV                      | Continue for at least 2–6 h after bleeding stops. <b>No more than 3 L of oxytocin-containing fluid</b> (water intoxication) | Rapid IV bolus → hypotension. Very safe otherwise. Some Ugandan units use 20–40 IU in 1 L; follow your unit                                                       |
| <b>Misoprostol</b>                               | <b>800 µg</b> (4 × 200 µg)                                                                                               | <b>Sublingual</b>                                 | Single dose                                                                                                                 | Use if oxytocin unavailable or as an add-on. Shivering and fever (up to 40 °C) are common and harmless; don't mistake them for sepsis. Slower onset than oxytocin |
| <b>Ergometrine</b> (ergonovine)                  | <b>0.2 mg</b>                                                                                                            | IM, or IV <b>slowly</b>                           | Repeat 0.2 mg IM after 15 min, then 0.2 mg every 4 h; <b>max 5 doses (1 mg) in 24 h</b>                                     | <b>Contraindicated in hypertension, pre-eclampsia, heart disease, retained placenta.</b> Causes vomiting and severe hypertension; can cause stroke or eclampsia   |
| <b>Carboprost</b> (15-methyl PGF <sub>2α</sub> ) | <b>0.25 mg</b>                                                                                                           | <b>Deep IM</b> (or intramyometrial by specialist) | Every 15 min; <b>max 8 doses (2 mg)</b>                                                                                     | <b>Contraindicated in asthma</b> (bronchospasm). Diarrhoea, vomiting, fever. Expensive and rarely stocked in Uganda                                               |
| <b>Tranexamic acid</b>                           | <b>1 g</b> (10 mL of 100 mg/mL)                                                                                          | IV over <b>10 min</b> (1 mL/min)                  | Second 1 g if bleeding continues after 30 min or restarts within 24 h                                                       | Give <b>within 3 h of birth</b> for <b>all causes</b> of PPH (tone, trauma, tissue). Too-fast injection causes hypotension                                        |

#### CLINICAL PEARL

**Carbetocin is for prevention only.** It is not a treatment drug for established PPH. And there is no point giving IV ergometrine to a woman whose BP is 160/110. If you remember nothing else about contraindications: "**Ergometrine: not for the hypertensive heart; carboprost: not for the wheezy chest.**"

### D. Escalation: bleeding continues after the bundle

Work through these **in parallel**, not one by one, while someone prepares theatre and blood.

#### 1. Bimanual uterine compression (buys time):

- One gloved fist in the anterior vaginal fornix pushes against the anterior uterine wall. The other hand, on the abdomen, presses the fundus down and forward onto the fist. The uterus is compressed between the two hands.
- Maintain firm pressure for at least 5–10 minutes (up to 20–30), until bleeding stops or help arrives.
- **External aortic compression** (fist pressed just above and left of the umbilicus onto the aorta until the femoral pulse disappears) is a temporising alternative, especially during transfer.

#### 2. Uterine balloon tamponade (UBT): for atonic PPH not responding to uterotonics.

- The WHO (2021) recommends UBT **where standard PPH care, including surgery and blood, is accessible.** In practice in Uganda the **condom catheter** is widely used as a bridge to theatre or during referral.
- **Condom catheter:** tie a condom over a Foley catheter (or use a purpose-made balloon). Insert it into the uterine cavity past the cervix. Fill with **300–500 mL** warm saline through a giving set until bleeding stops, and pack the vagina with gauze to hold it in. Continue the **oxytocin infusion**, give **prophylactic antibiotics** (e.g. ampicillin 2 g IV single dose, or a cephalosporin), catheterise the bladder, monitor closely. Deflate gradually and remove after **6–24 hours** (usually 12–24 h), with theatre and blood ready.
- **Tamponade test:** if bleeding stops, the tamponade is working. If bleeding continues through or around the balloon, go straight to laparotomy.

**3. Non-pneumatic anti-shock garment (NASG):** neoprene segments wrapped from the ankles upwards to the abdomen. They compress the lower body and pelvis, squeezing blood back to the heart, brain and lungs and reducing pelvic bleeding. It is a **first-aid device to buy time**, particularly for **transfer from HC III/IV to hospital**. Apply ankle segments first; remove only when stable (SBP  $\geq 90$  and pulse  $< 100$ , blood running), **from the ankles upwards**, one segment at a time, rechecking BP after each (a drop of 20 mmHg or a pulse rise of 20 means reapply).

**4. Transfuse:** whole blood or packed red cells as soon as possible. In major haemorrhage give **FFP** (15 mL/kg) when more than about 4 units of red cells have been given or clotting is abnormal, **cryoprecipitate** if fibrinogen is  $< 2$  g/L, and **platelets** if  $< 75 \times 10^9$ /L (availability at RRH/NRH). Aim for Hb  $> 8$  g/dL once bleeding is controlled.

#### 5. Surgery (laparotomy, at CEmONC level), from the most conservative step upward:

| Step                                               | What / how                                                                                                                                                                                          | Why                                                                        |
|----------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------|
| Exploration                                        | Evacuate clots, check for rupture, repair lacerations                                                                                                                                               | Finds the trauma you missed                                                |
| B-Lynch compression suture                         | Absorbable suture (e.g. No. 1 or 2 chromic/polyglactin on a large needle) that "braces" the uterus over the fundus; test first with bimanual compression (if that stops bleeding, B-Lynch will too) | Mechanically compresses an atonic uterus while preserving fertility        |
| Bilateral uterine artery ligation (O'Leary stitch) | Ligate the ascending branches at the level of the lower segment, including some myometrium                                                                                                          | The uterine arteries carry about 90% of uterine blood; cuts pulse pressure |

| Step                                           | What / how                                           | Why                                                                                                                                                 |
|------------------------------------------------|------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Utero-ovarian / internal iliac ligation</b> | Specialist procedures                                | Further devascularisation                                                                                                                           |
| <b>Subtotal (or total) hysterectomy</b>        | <b>Decide early</b> ; don't wait until she is in DIC | Life-saving. Subtotal is faster and safer in an unstable woman; total is needed if the lower segment or cervix bleeds (praevia/PAS, cervical tears) |

#### EXAM FAVOURITE

"When would you do a hysterectomy?" Answer: when **conservative measures have failed** and the mother's life is at risk, **before** she becomes moribund. A living mother without a uterus is better than a dead mother with one. Get consent early (from the woman or her next of kin) and involve a senior.

## E. Cause-specific management

### Trauma

- **Perineal/vaginal tears:** repair in good light with analgesia (lidocaine 1%). Secure the **apex first** (start the stitch 1 cm above it), because retracted vessels keep bleeding.
- **Cervical tears:** sponge forceps on either side, suture from the apex. Refer if you cannot see the apex.
- **Vulval/vaginal haematoma:** small (<5 cm) and not expanding → ice, analgesia, observe. Large or expanding → incise, evacuate, ligate bleeders, pack; catheterise.
- **Uterine rupture:** resuscitate, then **laparotomy**: repair (if clean tear and she wants more children) or hysterectomy; inspect the bladder.
- **Uterine inversion:** call for help, **stop oxytocin**, treat shock, and **replace immediately**. **Johnson manoeuvre:** push the fundus back through the cervix with the palm, the part that came out last going in first; don't remove the placenta first if still attached. If that fails, use **O'Sullivan hydrostatic** replacement (warm saline into the vagina with the introitus sealed by hand or a ventouse cup), or surgical correction (Huntington, Haultain). **After replacement, give oxytocin** and hold the uterus in place until it contracts. Give antibiotics.

### Tissue: retained placenta

1. If the placenta has not delivered **30 minutes** after birth: empty the bladder, **oxytocin 10 IU IV/IM** (if not already given as part of AMTSL, or a repeat), and repeat **CCT**. **Don't give ergometrine** (tetanic contraction can trap the placenta).
2. If still retained, or if bleeding: **manual removal of placenta (MRP)** under analgesia or anaesthesia (ketamine, spinal or pethidine + diazepam). Give **single-dose prophylactic antibiotics** (e.g. ampicillin 2 g IV, or cefazolin 2 g IV). The hand follows the cord into the uterus and finds the cleavage plane with the ulnar edge of the fingers while the other hand steadies the fundus. Remove it whole, check it, re-explore, and follow with an oxytocin infusion.
3. If there is **no cleavage plane**, suspect **placenta accreta spectrum**. Stop, don't pull pieces out, and call a senior. Laparotomy or hysterectomy may be needed.
4. **Retained fragments:** manual exploration or **MVA / blunt curettage** with a large curette (the soft postpartum uterus perforates easily).

### Thrombin (coagulopathy)

- Treat the cause (deliver an abruption, treat sepsis), give **TXA**, fresh whole blood (contains clotting factors and platelets, useful where components are unavailable), FFP, cryoprecipitate and platelets. Keep her **warm**; correct calcium and acidosis.

## F. Setting-specific plan

| Level                                  | What to do                                                                                                                                                                                                                                                                                               |
|----------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| HC II / community                      | Misoprostol 600 µg orally for prevention; uterine massage; refer urgently                                                                                                                                                                                                                                |
| HC III (BEmONC)                        | AMTSL; full E-MOTIVE bundle (massage, oxytocin, TXA, fluids, genital exam, repair tears); MRP; MVA for fragments; bimanual compression; condom tamponade and NASG; <b>early referral</b> to HC IV/hospital with IV running, oxytocin infusion, NASG on, health worker escort, referral note, phone ahead |
| HC IV / general hospital (CEmONC)      | All of the above <b>plus</b> blood transfusion, theatre: laparotomy, B-Lynch, uterine artery ligation, hysterectomy                                                                                                                                                                                      |
| RRH / NRH (e.g. Mulago-Kawempe, Mengo) | Blood components (FFP, cryo, platelets), HDU/ICU, specialist surgery for PAS, interventional options where available                                                                                                                                                                                     |

## G. Secondary PPH (24 h to 12 weeks)

Causes: **endometritis** (most common), **retained products**, subinvolution of the placental site, rarely trophoblastic disease (GTN) or arteriovenous malformation.

1. Assess and resuscitate as for primary PPH. Most secondary PPH is modest, but it can be massive.
2. **Investigations:** FBC, blood culture and high vaginal swab, pelvic ultrasound (retained products), **β-hCG if bleeding persists** (to exclude GTN), malaria test if febrile.
3. **Antibiotics** if infection is suspected: **ampicillin 2 g IV 6-hourly + gentamicin 5 mg/kg IV once daily + metronidazole 500 mg IV 8-hourly** until afebrile for 48 hours, then oral amoxicillin plus metronidazole to complete 7 days (or local regimen, e.g. ceftriaxone + metronidazole).
4. **Uterotonics:** oxytocin infusion, misoprostol 800 µg SL, or ergometrine if no hypertension.
5. **Evacuation of retained products** by an experienced person (MVA or gentle curettage), **after antibiotics have started** (the infected uterus perforates easily).
6. Persistent bleeding: surgical options as above.

## H. After the bleeding stops

- Continue the oxytocin infusion for several hours. Monitor vital signs, fundus and loss every 15 min for 2 hours, then half-hourly and hourly; fluid balance chart; urine output.
- Hb check at 24 h; **treat anaemia** (ferrous sulphate 200 mg + folic acid; IV iron at referral level; transfusion if symptomatic or Hb <7 g/dL).
- Thromboprophylaxis once bleeding is controlled (massive PPH is a VTE risk).
- **Debrief** the woman and family; explain the risk next pregnancy (recurrence about 15%); document well and plan her next delivery at a CEmONC facility.
- Watch for **Sheehan's syndrome** (failure of lactation, amenorrhoea, fatigue after major PPH).
- **Near-miss audit or MPDSR** review if there was a death.

## Complications

| Maternal     |                                                                                                                                                                                  |
|--------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Immediate    | Hypovolaemic shock, death, DIC, transfusion reactions, cardiac arrest                                                                                                            |
| Organ damage | <b>Acute kidney injury</b> (ATN, cortical necrosis), ARDS, <b>Sheehan's syndrome</b> (pituitary necrosis → failed lactation, amenorrhoea, hypothyroidism, adrenal insufficiency) |

| Maternal       |                                                                                                                                                                                      |
|----------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| From treatment | Hysterectomy (loss of fertility), bladder/ureteric injury, infection after MRP or tamponade, dilutional coagulopathy, fluid overload/pulmonary oedema, water intoxication (oxytocin) |
| Later          | Severe anaemia, fatigue, failure of breastfeeding, <b>VTE</b> , post-traumatic stress and depression, Asherman's syndrome (after curettage)                                          |

## Ward-round & exam pearls

### Model summary line:

"Mrs N, a 32-year-old para 5 at 39 weeks, delivered a 4.1 kg baby by SVD 40 minutes ago after a 16-hour oxytocin-augmented labour. She developed primary PPH from uterine atony, with measured loss of 900 mL on the calibrated drape. Her pulse was 118, BP 96/60, shock index 1.2. The E-MOTIVE bundle was started at 10:05: massage, oxytocin 10 IU IV plus infusion, TXA 1 g IV, 1 L Ringer's, catheterised, and genital tract inspected (second-degree tear repaired, placenta complete). The uterus is now firm, pulse 100, BP 110/70, urine output 40 mL/h. Hb 8.1 g/dL; 2 units cross-matched. Plan: continue the oxytocin infusion, 15-minute observations, transfuse 1 unit, repeat Hb at 24 h, oral iron, and document as severe PPH for review."

### Likely viva questions

1. *Define PPH.*  $\geq 500$  mL within 24 h of birth; severe  $\geq 1000$  mL. WHO 2025: act at  $\geq 300$  mL with abnormal vital signs. Secondary: 24 h–12 weeks.
2. *What are the components of AMTSL?* Uterotonic within 1 min (oxytocin 10 IU IM), CCT, and uterine tone assessment (plus delayed cord clamping and examining the placenta).
3. *Name the components of the E-MOTIVE bundle.* Massage, Oxytocics, TXA, IV fluids, Examination, Escalation, triggered by objective measurement with a calibrated drape.
4. *Why is TXA time-critical?* Fibrinolysis peaks early; benefit is seen only within 3 h and falls by about 10% for every 15 minutes' delay.
5. *Contraindications to ergometrine?* Hypertension/pre-eclampsia, cardiac disease, retained placenta (and a second twin undelivered).
6. *What is a Couvelaire uterus, and is it an indication for hysterectomy?* Blood extravasated into the myometrium from abruption. **No**; the uterus usually contracts.
7. *How do you manage uterine inversion?* Call help, stop uterotonics, resuscitate, immediate manual replacement (Johnson), then oxytocin.

**OSCE tips:** on the PPH drill station, **say "I will call for help" first**, state the time, assign roles, verbalise ABC, and use the E-MOTIVE sequence out loud. Demonstrate bimanual compression correctly (fist in the anterior fornix). Always **check the placenta**. Mention **documentation** and **debrief**.

## Mnemonics & memory aids

- **4 Ts:** Tone, Trauma, Tissue, Thrombin (70/20/10/1).
- **E-MOTIVE** (E is for early detection): **M**assage, **O**xytocics, **T**ranexamic acid, **I**V fluids, **V**aginal (genital-tract) examination, **E**scalation.
- **HAEMOSTASIS** (the classic RCOG escalation ladder): **H**elp and hands on the uterus, **A**ssess and resuscitate, **E**stablish aetiology / ecobolics, **M**assage, **O**xytocin infusion / prostaglandins, **S**hift to theatre (EUA, bimanual compression), **T**amponade (balloon), **A**pply compression sutures (B-Lynch), **S**ystematic pelvic devascularisation, **I**nterventional radiology, **S**ubtotal or total hysterectomy.
- **"Ergo: no BP, no heart; carbo: no wheeze."**

- **TXA "1-10-3-30"**: 1 g, over **10** minutes, within **3** hours, repeat after **30** minutes if still bleeding.

## High-yield summary

### Numbers & doses to know

| Item                      | Value                                                                                                                                             |
|---------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------|
| Primary PPH               | <b>≥500 mL within 24 h</b> of birth; or <b>≥300 mL + any abnormal vital sign</b> (WHO/FIGO/ICM 2025)                                              |
| Severe / major PPH        | <b>≥1000 mL</b> (RCOG: 1000–2000 moderate, >2000 severe)                                                                                          |
| Secondary PPH             | <b>24 h to 12 weeks</b> after birth                                                                                                               |
| Retained placenta         | Not delivered <b>30 min</b> after birth (WHO; UK 60 min)                                                                                          |
| AMTSL uterotonic          | <b>Oxytocin 10 IU IM</b> within <b>1 min</b> of birth; tone checks <b>every 15 min for 2 h</b>                                                    |
| Treatment oxytocin        | <b>10 IU IV slowly/IM</b> , then <b>20 IU in 1 L at 60 drops/min</b> , maintenance <b>40 drops/min</b> ; max <b>3 L</b> oxytocin-containing fluid |
| Tranexamic acid           | <b>1 g IV over 10 min</b> , within <b>3 h</b> ; repeat 1 g after <b>30 min</b> if still bleeding ("1-10-3-30")                                    |
| Misoprostol               | Prevention <b>600 µg orally</b> (400–600 µg); treatment <b>800 µg sublingual</b>                                                                  |
| Ergometrine               | <b>0.2 mg IM/slow IV</b> , repeat after 15 min then 4-hourly; <b>max 1 mg (5 doses)/24 h</b>                                                      |
| Carboprost                | <b>0.25 mg deep IM</b> every 15 min; <b>max 2 mg (8 doses)</b>                                                                                    |
| Carbetocin (heat-stable)  | <b>100 µg IM/IV</b> , prevention only                                                                                                             |
| Shock index (HR ÷ SBP)    | Normal 0.7–0.9; <b>≥0.9 abnormal; ≥1.4 urgent; ≥1.7 critical</b>                                                                                  |
| Pad weighing              | <b>1 g ≈ 1 mL</b> (subtract dry weight)                                                                                                           |
| Clotting test             | 2 mL in a dry glass tube; <b>no clot by 7 min</b> = coagulopathy (WHO)                                                                            |
| Fibrinogen                | Normal at term 4–6 g/L; <b>&lt;2 g/L</b> = severe coagulopathy → cryoprecipitate                                                                  |
| Platelets                 | Transfuse if <b>&lt;75 × 10<sup>9</sup>/L</b> in active bleeding                                                                                  |
| FFP                       | <b>15 mL/kg</b> after about 4 units of red cells or with abnormal clotting                                                                        |
| Urine output target       | <b>≥30 mL/h</b>                                                                                                                                   |
| Balloon tamponade         | <b>300–500 mL</b> warm saline; remove after <b>6–24 h</b>                                                                                         |
| NASG removal              | When <b>SBP ≥90 and pulse &lt;100</b> , ankles first; reapply if SBP falls 20 or pulse rises 20                                                   |
| Recurrence next pregnancy | About <b>15%</b>                                                                                                                                  |

### Classic exam traps

- **"Wait until the drape reaches 500 mL."** Wrong if vital signs are abnormal: at **≥300 mL plus an abnormal vital sign**, it is already PPH.
- **"BP is normal, so the loss is small."** Hypotension is late; look at the **pulse and shock index**.
- **"Give a second uterotonic"** for a **firm** uterus that keeps bleeding. Wrong: find the **tear** or check **clotting**.
- **"Give drugs one at a time and wait to see."** Wrong: the E-MOTIVE bundle is given **together**.
- **"TXA helps whenever it is given."** Wrong: only within **3 h** of birth.
- **Ergometrine for retained placenta** (traps it) or for a hypertensive woman (stroke, eclampsia). Give oxytocin.
- **"Hb 10 at 30 minutes, so no transfusion needed."** The first Hb underestimates acute loss.

- **Oxytocin before replacing an inverted uterus.** Replace first, then give oxytocin.

## Questions seniors ask

**What is your first action when the midwife calls "PPH"?** Call for help and note the time, put a hand on the fundus and massage, and start the bundle together while assessing ABC and the shock index.

**The uterus is firm but she is still bleeding. What now?** Inspect the genital tract in good light (cervix all round, vagina, perineum), check the placenta is complete, and do a bedside clotting test; continue resuscitation and give TXA if within 3 h.

**How do you measure blood loss here?** Calibrated drape under the buttocks from birth, then weigh pads and linen (1 g  $\approx$  1 mL, minus dry weight); visual estimates are 30–50% too low.

**Why give TXA even for atony?** Fibrinolysis rises quickly after birth for every cause of PPH; WOMAN showed about a 30% reduction in death from bleeding when given within 3 h.

**Which uterotonic would you avoid in this woman with pre-eclampsia, and what would you use?** Avoid ergometrine (and Syntometrine); use oxytocin, then misoprostol 800  $\mu$ g SL (carboprost if available and she is not asthmatic).

**Her Hb is 9.8 g/dL after a 1.5 L bleed. Is she fine?** No. The first Hb has not equilibrated; repeat it after resuscitation and at 24 h, and transfuse on clinical grounds and measured loss.

**When would you refer from a HC III, and how?** As soon as bleeding is not controlled by the bundle, or she needs blood or theatre: with IV oxytocin running, NASG and/or tamponade, a health worker who can do bimanual compression, a referral note, and a phone call ahead.

**When would you do a hysterectomy?** When conservative measures fail and her life is at risk, decided early by a senior, before she is moribund in DIC.

# Flashcards

26 cards. Cover the right-hand column and test yourself.

| #  | Question                                                    | Answer                                                                                                                                                                                                                                                                    |
|----|-------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 1  | Define primary PPH (classic and WHO/FIGO/ICM 2025).         | Classic: blood loss <b>≥500 mL</b> from the genital tract within <b>24 h</b> of birth. WHO/FIGO/ICM 2025 adds: diagnose and act at <b>≥300 mL plus any abnormal vital sign</b> . Severe PPH <b>≥1000 mL</b> .                                                             |
| 2  | Define secondary PPH and name its commonest cause.          | Abnormal bleeding from <b>24 h to 12 weeks</b> after birth. Commonest cause: <b>endometritis</b> , often with <b>retained products</b> .                                                                                                                                  |
| 3  | The 4 Ts of PPH with approximate proportions.               | <b>Tone</b> ~70%, <b>Trauma</b> ~20%, <b>Tissue</b> ~10%, <b>Thrombin</b> <1%.                                                                                                                                                                                            |
| 4  | Components of active management of the third stage (AMTSL). | <ol style="list-style-type: none"> <li><b>Oxytocin 10 IU IM</b> within 1 min of birth (most important)</li> <li>Controlled cord traction</li> <li>Uterine tone assessment every 15 min for 2 h</li> </ol> Plus delayed cord clamping (1–3 min) and checking the placenta. |
| 5  | What does E-MOTIVE stand for?                               | Early detection with a calibrated drape, then the bundle: <b>Massage</b> , <b>Oxytocic</b> drugs, <b>Tranexamic acid</b> , <b>IV fluids</b> , <b>Vaginal/genital-tract examination</b> , <b>Escalation</b> .                                                              |
| 6  | Main result of the E-MOTIVE trial (NEJM 2023).              | Severe PPH, laparotomy for bleeding or death from bleeding fell from <b>4.3% to 1.6%</b> (RR 0.40, i.e. a <b>60% reduction</b> ) in 210,132 vaginal births in Kenya, Nigeria, South Africa and Tanzania.                                                                  |
| 7  | Tranexamic acid for PPH: dose, route, timing, repeat.       | <b>1 g IV over 10 min, within 3 h of birth</b> ; repeat 1 g if bleeding continues after <b>30 min</b> or restarts within 24 h (WOMAN trial). Memory: <b>1-10-3-30</b> .                                                                                                   |
| 8  | Oxytocin regimen for treating PPH (WHO).                    | <b>10 IU IV slowly/IM</b> , then <b>20 IU in 1 L NS/RL at 60 drops/min</b> , then <b>20 IU in 1 L at 40 drops/min</b> . Max <b>3 L</b> of oxytocin-containing fluid (water intoxication).                                                                                 |
| 9  | Ergometrine: dose, maximum, contraindications.              | <b>0.2 mg IM</b> (or slow IV); repeat after 15 min, then 4-hourly; <b>max 5 doses (1 mg)</b> . Contraindicated in <b>hypertension/pre-eclampsia</b> , <b>heart disease</b> , retained placenta.                                                                           |
| 10 | Carboprost: dose, maximum, key contraindication.            | <b>0.25 mg deep IM</b> every 15 min, <b>max 8 doses (2 mg)</b> . Contraindicated in <b>asthma</b> (bronchospasm).                                                                                                                                                         |
| 11 | Misoprostol doses: prevention vs treatment of PPH.          | Prevention: <b>400–600 µg orally</b> (Uganda community distribution: 600 µg). Treatment: <b>800 µg sublingual</b> . Side effects: shivering, fever.                                                                                                                       |
| 12 | Heat-stable carbetocin: dose and role.                      | <b>100 µg IM/IV</b> single dose, for <b>prevention only</b> where oxytocin quality or cold chain can't be assured. Non-inferior to oxytocin (CHAMPION trial). <b>Not</b> for treatment.                                                                                   |
| 13 | Shock index: formula and thresholds in obstetrics.          | <b>HR ÷ SBP</b> . Normal postpartum 0.7–0.9. <b>≥0.9</b> = concern; <b>≥1.4</b> = critical, needs urgent intervention or transfer. It rises before BP falls.                                                                                                              |
| 14 | How do you perform bimanual uterine compression?            | Fist in the <b>anterior vaginal fornix</b> pressing on the anterior wall; the other hand on the abdomen pushes the fundus down and forward onto the fist. Hold firm pressure until bleeding stops (at least 5–10 min).                                                    |

| #  | Question                                                  | Answer                                                                                                                                                                                                                                                                                                                                                                  |
|----|-----------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 15 | Condom catheter tamponade: key steps.                     | Condom tied over a Foley catheter, inserted into the uterine cavity, filled with <b>300–500 mL</b> warm saline until bleeding stops; vaginal pack; continue <b>oxytocin infusion</b> ; <b>antibiotic prophylaxis</b> ; catheterise; remove after <b>6–24 h</b> . If bleeding continues, go to laparotomy.                                                               |
| 16 | When is a placenta 'retained' and what is the first step? | Not delivered <b>30 min</b> after birth (WHO). Empty the bladder, give <b>oxytocin 10 IU IM/IV</b> , repeat CCT. If unsuccessful or bleeding → <b>manual removal</b> under anaesthesia with <b>single-dose prophylactic antibiotic</b> . Don't give ergometrine.                                                                                                        |
| 17 | Uterine inversion: signs and immediate management.        | Fundus not palpable abdominally, mass in the vagina, <b>shock out of proportion to blood loss</b> (often bradycardia). Call help, <b>stop oxytocin</b> , resuscitate, <b>immediate manual replacement (Johnson)</b> , then give oxytocin. Hydrostatic (O'Sullivan) if that fails.                                                                                       |
| 18 | Surgical options for refractory atonic PPH, in order.     | Exploration → <b>B-Lynch</b> compression suture → <b>bilateral uterine artery ligation</b> → internal iliac ligation (specialist) → <b>subtotal/total hysterectomy</b> . Decide on hysterectomy early.                                                                                                                                                                  |
| 19 | Bedside test for coagulopathy at a HC IV.                 | <b>20-minute whole blood clotting test</b> : 2 mL of venous blood in a clean dry glass tube; if no clot at 20 minutes, suspect coagulopathy (DIC).                                                                                                                                                                                                                      |
| 20 | Sheehan's syndrome: what, and how does it present?        | Ischaemic necrosis of the enlarged pituitary after massive PPH. Presents with <b>failure of lactation</b> , amenorrhoea, fatigue, hypothyroidism, adrenal insufficiency (hyponatraemia, hypoglycaemia).                                                                                                                                                                 |
| 21 | NASG: what it does and how to remove it.                  | Non-pneumatic anti-shock garment: compresses the legs and pelvis, raising central blood volume and reducing pelvic bleeding. A <b>first-aid bridge for transfer</b> . Remove only when stable (SBP $\geq 90$ , pulse $< 100$ ), <b>starting at the ankles</b> , one segment at a time; reapply if BP falls by 20 mmHg or pulse rises by 20.                             |
| 22 | PPH: what must be done in the first 5 minutes?            | Say 'PPH' and note the time; <b>call for help</b> and assign roles; <b>hand on the fundus and massage</b> ; ABC, vitals and <b>shock index</b> ; lie flat, oxygen, keep warm; <b>two 14–16G cannulae</b> with bloods (Hb, group and cross-match, <b>20-min clotting tube</b> ); start the <b>bundle together</b> (oxytocin, TXA, fluids, catheter, genital-tract exam). |
| 23 | How do you add weighed pads to the drape reading?         | Weigh the soaked pads/linen, <b>subtract the weight of the same items dry</b> , and count <b>1 g <math>\approx</math> 1 mL</b> . Example: 620 g wet – 120 g dry = <b>500 mL</b> extra. Visual estimates underestimate loss by 30–50%.                                                                                                                                   |
| 24 | Why is the first Hb after a PPH misleading?               | She has lost <b>whole blood</b> and there has not been time for <b>haemodilution</b> , so the first Hb <b>underestimates</b> the loss. Treat the measured loss and the clinical picture; repeat Hb after resuscitation and at <b>24 h</b> .                                                                                                                             |
| 25 | Blood-product triggers in major PPH.                      | Red cells early; <b>FFP 15 mL/kg</b> after about 4 units of red cells or with abnormal clotting; <b>cryoprecipitate</b> if fibrinogen <b><math>&lt; 2</math> g/L</b> ; <b>platelets</b> if <b><math>&lt; 75 \times 10^9/L</math></b> . Fresh whole blood if components are unavailable. Keep her warm.                                                                  |
| 26 | Pulse 126, BP 90/60 after birth. Shock index and meaning? | <b>SI = <math>126 \div 90 = 1.4</math></b> → needs <b>urgent intervention</b> : senior at the bedside, blood now, prepare theatre or refer. ( $\geq 0.9$ abnormal; $\geq 1.7$ critical.)                                                                                                                                                                                |

## SELF-TEST

# Single best answer questions

15 questions. The answers and explanations follow at the end.

**1.** A 29-year-old para 4 delivers a 4.2 kg baby at a HC IV. Ten minutes after the placenta delivers complete, the calibrated drape reads 350 mL. Her pulse is 112/min and BP 98/60 mmHg (shock index 1.14). The uterus is soft. Which is the MOST appropriate action?

- A Observe for 15 minutes and act if the drape reaches 500 mL
- B Give ergometrine 0.5 mg IV and wait to see the response
- C Diagnose PPH and immediately start the full E-MOTIVE bundle
- D Refer to the regional referral hospital without treatment
- E Give misoprostol 800 µg sublingually alone and recheck in 30 minutes

**2.** A 24-year-old primigravida delivers by vacuum extraction. The placenta is complete and the uterus is firm and well contracted at the umbilicus, but a steady stream of bright red blood continues. Pulse 104/min. What is the MOST likely cause of her bleeding?

- A Genital tract trauma
- B Uterine atony
- C Retained placental tissue
- D Disseminated intravascular coagulation
- E Uterine inversion

**3.** A 34-year-old para 3 with severe pre-eclampsia (BP 162/108 mmHg on MgSO<sub>4</sub>) develops atonic PPH after vaginal birth. She has received oxytocin 10 IU IV, an oxytocin infusion, TXA 1 g and uterine massage, but the uterus remains soft. Which additional uterotonic is MOST appropriate?

- A Ergometrine 0.2 mg IV
- B Oxytocin-ergometrine (Syntometrine) 1 ampoule IM
- C Carboprost 0.25 mg IV bolus
- D Misoprostol 800 µg sublingual
- E Methylergometrine 0.2 mg IM

**4.** A woman delivered 4 hours ago at home and arrives at a HC III with ongoing PPH. She is pale, pulse 124/min. A student suggests tranexamic acid. Which statement is MOST accurate? Advanced

- A TXA should be given because it helps whenever it is given
- B TXA is not expected to help because it is more than 3 hours since birth; focus on uterotonics, fluids, the cause and referral
- C TXA should be given as 1 g IM because IV access is difficult
- D TXA is contraindicated in all postpartum women because of thrombosis
- E TXA is only for trauma causes of PPH, not atony

**5.** Thirty minutes after a normal delivery at a HC III, the placenta of a para 2 has not delivered despite oxytocin 10 IU IM at birth and one attempt at controlled cord traction. There is minimal bleeding and she is stable. What is the BEST next step?

- A Give ergometrine 0.5 mg IV to expel the placenta
- B Inject oxytocin into the umbilical vein
- C Pull harder on the cord with fundal pressure
- D Wait another 2 hours before any action
- E Empty the bladder, give a further 10 IU oxytocin and repeat CCT; if unsuccessful, manual removal with antibiotic cover

**6.** Immediately after a fundal-pressure delivery and vigorous cord traction, a para 1 collapses: pulse 52/min, BP 70/40 mmHg, with only about 400 mL of visible blood loss. The fundus cannot be palpated abdominally and a mass is felt in the vagina. What is the MOST important immediate action? Advanced

- A Start an oxytocin infusion at 60 drops/min
- B Attempt manual removal of the placenta
- C Call for help, stop uterotonics, and immediately replace the uterus manually
- D Give atropine and wait for the heart rate to recover
- E Arrange a pelvic ultrasound scan

**7.** Which intervention is the MOST effective single measure to prevent PPH at a vaginal birth?

- A Routine episiotomy
- B Sustained uterine massage for 30 minutes in every woman
- C Early cord clamping within 30 seconds
- D Prophylactic uterotonic (oxytocin 10 IU IM/IV) within 1 minute of birth
- E Routine tranexamic acid for all vaginal births

**8.** A 30-year-old para 6 at a regional referral hospital has atonic PPH continuing despite the full E-MOTIVE bundle, misoprostol 800 µg SL, and 20 minutes of bimanual compression. Measured loss is 1800 mL; she has received 2 units of blood; theatre is being prepared. What is the MOST appropriate temporising measure while theatre is prepared? Advanced

- A Uterine balloon (condom catheter) tamponade
- B Third dose of TXA
- C Carboprost 0.25 mg IV
- D Vaginal pack of gauze only
- E Stop fluids to allow permissive hypotension for 2 hours

**9.** A 27-year-old has had a PPH of 1500 mL after delivery of a stillborn baby with a large retroplacental clot. Blood is oozing from her cannula sites and the vaginal blood does not clot. The HC IV has no laboratory at night. What is the BEST bedside test to confirm your suspicion?

- A Kleihauer–Betke test
- B Urine dipstick for protein
- C Random blood sugar
- D 20-minute whole blood clotting test
- E Peripheral smear for malaria parasites

- 10.** At a HC III 3 hours' drive from the nearest hospital, a woman has PPH controlled only while bimanual compression is maintained. The bundle has been given. What is the BEST plan for transfer?
- A Send her by boda-boda with her relatives and a referral note
  - B Wait until the bleeding stops completely before referral
  - C Apply NASG and/or balloon tamponade, keep IV oxytocin running, and transfer by ambulance with a health worker after calling ahead
  - D Give ergometrine 0.2 mg IV every 15 minutes during transport regardless of BP
  - E Remove the IV line so it does not come out during transport
- 11.** A 25-year-old known asthmatic has atonic PPH not responding to oxytocin and misoprostol. Which drug should be AVOIDED?
- A Ergometrine
  - B Tranexamic acid
  - C Oxytocin infusion
  - D Carboprost (15-methyl PGF<sub>2</sub>α)
  - E Ringer's lactate
- 12.** Six hours after an emergency caesarean section for obstructed labour, a 31-year-old has pulse 128/min, BP 88/56 mmHg and a distended, tender abdomen. Vaginal loss is minimal and the uterus feels firm. Hb has fallen from 10.5 to 6.8 g/dL. What is the MOST appropriate management? Advanced
- A Increase the oxytocin infusion and observe
  - B Give misoprostol 800 µg rectally
  - C Insert a balloon tamponade
  - D Transfuse 2 units and discharge to the ward
  - E Resuscitate with blood and return to theatre for re-laparotomy
- 13.** What is the main mechanism by which bleeding from the placental bed is controlled after delivery?
- A Contraction of the interlacing myometrial fibres compressing the spiral arteries
  - B Vasoconstriction of the spiral arteries by their smooth muscle
  - C Platelet plug formation alone
  - D Reduction in cardiac output after delivery
  - E Increase in plasma fibrinolytic activity
- 14.** A para 3 is readmitted on day 10 after SVD with heavy fresh vaginal bleeding, temperature 38.6 °C, foul-smelling lochia and a tender, bulky uterus. Pulse 108/min, BP 110/70 mmHg. What is the BEST initial management?
- A Immediate curettage before antibiotics
  - B IV ampicillin, gentamicin and metronidazole plus uterotonics, ultrasound, and evacuation of any retained products once antibiotics have started
  - C Oral amoxicillin and discharge home
  - D Emergency hysterectomy
  - E Tranexamic acid alone

**15.** During laparotomy for atonic PPH in a 22-year-old para 1 who strongly wishes more children, bimanual compression of the uterus on the operating table stops the bleeding. What is the MOST appropriate next surgical step? Advanced

- A Total abdominal hysterectomy
- B Internal iliac artery ligation by the general surgeon
- C Close the abdomen and give more oxytocin
- D Subtotal hysterectomy
- E B-Lynch uterine compression suture

## Answers & explanations

**1. C** · Diagnose PPH and immediately start the full E-MOTIVE bundle

C is right: WHO/FIGO/ICM 2025 advise diagnosing PPH at **≥300 mL with abnormal vital signs**. Tachycardia and a shock index >0.9 qualify, so the whole bundle (massage, oxytocin, TXA, IV fluids, examination, escalation) should start now. A is wrong: waiting for 500 mL is exactly the delay the new threshold aims to prevent. B is wrong: ergometrine is second-line and 0.5 mg is the wrong dose (0.2 mg), and giving single drugs sequentially is less effective than the bundle. D is wrong: never refer without first-response treatment; the HC IV can give all of the bundle. E is wrong: misoprostol alone, with a 'wait and see' interval, is not the bundle; oxytocin is first line.

**2. A** · Genital tract trauma

A is right: a **firm uterus with continuing bright-red bleeding after an instrumental birth** = trauma (cervical or vaginal laceration) until proven otherwise; inspect under good light and repair. B is wrong: atony gives a soft, boggy uterus. C is wrong: the placenta is complete, and retained tissue usually causes a uterus that relaxes repeatedly. D is wrong: DIC is rare without a trigger (abruption, AFE, sepsis) and presents with non-clotting blood and oozing. E is wrong: in inversion the fundus is not palpable at the umbilicus.

**3. D** · Misoprostol 800 µg sublingual

D is right: **misoprostol 800 µg SL** is a safe add-on uterotonic in hypertension. A, B and E are wrong: ergometrine and its combinations and derivatives cause vasoconstriction and severe hypertension; they are **contraindicated in pre-eclampsia** (risk of stroke or eclampsia). C is wrong: carboprost is safe in hypertension but is given **deep IM (0.25 mg)**, never as an IV bolus, and it is contraindicated in asthma.

**4. B** · TXA is not expected to help because it is more than 3 hours since birth; focus on uterotonics, fluids, the cause and referral

B is right: in the WOMAN trial, TXA reduced death from bleeding only when given **within 3 hours**, and there was no benefit (and possible harm) later. Management should focus on massage, uterotonics, fluids, finding the cause (4 Ts), NASG and urgent referral. A is wrong: benefit is time-dependent and falls about 10% per 15 min of delay. C is wrong: the recommended route is **IV over 10 minutes**; get access (IO or cut-down if needed). D is wrong: WOMAN showed no increase in thromboembolic events. E is wrong: TXA is recommended for PPH of **any cause** when given within 3 h.

**5. E** · Empty the bladder, give a further 10 IU oxytocin and repeat CCT; if unsuccessful, manual removal with antibiotic cover

E is right: for a retained placenta at 30 minutes, WHO recommends **additional oxytocin 10 IU** with CCT; a **full bladder** can prevent separation. If this fails, proceed to **manual removal** under anaesthesia with **single-dose prophylactic antibiotics**. A is wrong: ergometrine causes a tetanic contraction that can trap the placenta, and 0.5 mg is the wrong dose. B is wrong: WHO does not recommend umbilical vein oxytocin. C is wrong: excessive traction with fundal pressure risks cord avulsion and **uterine inversion**. D is wrong: prolonged waiting increases PPH and infection risk.

**6. C** · Call for help, stop uterotonics, and immediately replace the uterus manually

C is right: this is **acute uterine inversion**. Shock out of proportion to blood loss, with **vagal bradycardia**, and an absent fundus. Immediate manual replacement (Johnson manoeuvre) while the cervix is still lax is the priority, along with resuscitation; give oxytocin only **after** replacement. A is wrong: oxytocin before replacement contracts the cervical ring and traps the inverted fundus. B is wrong: removing the placenta before replacement increases bleeding. D is wrong: atropine doesn't correct the cause; delay allows the cervix to clamp down. E is wrong: it is a clinical diagnosis and imaging only delays treatment.

**7. D** · Prophylactic uterotonic (oxytocin 10 IU IM/IV) within 1 minute of birth

D is right: the **prophylactic uterotonic** is the most important component of AMTSL and reduces PPH by about 50–60%. A is wrong: routine episiotomy increases blood loss and trauma. B is wrong: sustained massage is not recommended when oxytocin is given; only tone checks every 15 min. C is wrong: **delayed** cord clamping (1–3 min) is recommended; early clamping doesn't prevent PPH and harms the baby's iron stores. E is wrong: TXA is not recommended routinely for prevention in all vaginal births.

**8. A** · Uterine balloon (condom catheter) tamponade

A is right: **uterine balloon tamponade** is recommended for refractory atonic PPH where surgery and blood are available, and it often stops bleeding or buys time; if bleeding continues through it, proceed straight to laparotomy. B is wrong: TXA is limited to a second dose, and further doses aren't recommended. C is wrong: carboprost is given IM or intramyometrially, not IV. D is wrong: a vaginal pack alone conceals ongoing intrauterine bleeding. E is wrong: prolonged permissive hypotension isn't appropriate; resuscitate with blood.

**9. D** · 20-minute whole blood clotting test

D is right: abruption with fetal death can cause **DIC**. The **20-minute whole blood clotting test** (no clot in a dry glass tube at 20 min) is a simple bedside test available anywhere. A is wrong: Kleihauer quantifies fetomaternal haemorrhage for anti-D dosing, not coagulopathy. B is wrong: proteinuria screens for pre-eclampsia (a risk factor for abruption) but doesn't show coagulopathy. C and E are wrong: useful in other contexts but irrelevant to the clotting question.

**10. C** · Apply NASG and/or balloon tamponade, keep IV oxytocin running, and transfer by ambulance with a health worker after calling ahead

C is right: the **NASG** and **tamponade** are temporising devices designed to keep a woman alive during transfer. Keep oxytocin running, send an escort who can do bimanual/aortic compression, and **call ahead** so blood and theatre are ready. A is wrong: this is unsafe transport with no skilled escort. B is wrong: waiting at a facility that can't transfuse or operate risks death. D is wrong: ergometrine is limited to 5 doses in 24 h and is contraindicated if hypertensive. E is wrong: IV access is essential.

**11. D** · Carboprost (15-methyl PGF<sub>2α</sub>)

D is right: **carboprost** (a prostaglandin F<sub>2α</sub> analogue) causes bronchoconstriction and is **contraindicated in asthma**. A is wrong: ergometrine is acceptable if she is normotensive without heart disease. B is wrong: TXA is safe and recommended. C and E are wrong: oxytocin and crystalloid are safe in asthma. Note that misoprostol (PGE<sub>1</sub>) does not cause bronchospasm.

**12. E** · Resuscitate with blood and return to theatre for re-laparotomy

E is right: **shock with a distended abdomen, falling Hb and little vaginal loss after CS** means intra-abdominal bleeding (a slipped uterine-angle ligature, extension of the incision into the broad ligament, bleeding vessels). Resuscitate and do a **re-laparotomy**. A, B and C are wrong: the uterus is contracted and the bleeding is not atonic or intrauterine, so uterotonics and tamponade won't stop it. D is wrong: transfusion without source control only delays the inevitable.

**13. A** · Contraction of the interlacing myometrial fibres compressing the spiral arteries

A is right: the '**living ligatures**'. Myometrial fibres contract and kink the vessels between them, which is why tone matters most. B is wrong: trophoblast invasion replaces the muscular wall of spiral arteries, so they **cannot** constrict. C is wrong: clotting is secondary and cannot control flow of 500–800 mL/min alone. D is wrong: cardiac output actually rises immediately after delivery (autotransfusion). E is wrong: fibrinolysis would **worsen** bleeding, which is why TXA helps.

**14. B** · IV ampicillin, gentamicin and metronidazole plus uterotonics, ultrasound, and evacuation of any retained products once antibiotics have started

B is right: this is **secondary PPH from endometritis ± retained products**. Resuscitate, give **broad-spectrum IV antibiotics** (ampicillin 2 g 6-hourly + gentamicin 5 mg/kg daily + metronidazole 500 mg 8-hourly) and uterotonics, scan, and evacuate retained products **after antibiotics have started**, by an experienced operator (a soft infected uterus perforates easily). A is wrong: curettage without antibiotic cover risks septicaemia and perforation. C is wrong: she is septic and bleeding, so needs admission. D is wrong: hysterectomy is a last resort. E is wrong: TXA alone doesn't treat infection or tissue.

**15. E** · B-Lynch uterine compression suture

E is right: if bimanual compression at laparotomy stops the bleeding (the 'compression test'), a **B-Lynch** brace suture is likely to work and preserves fertility. A and D are wrong: hysterectomy is for failure of conservative measures, or earlier if she is in extremis. It is not first choice in a young woman when a compression suture should work. B is wrong: internal iliac ligation is technically difficult, risks ureteric and venous injury, and is less effective than simpler steps. C is wrong: closing without definitive haemostasis risks re-bleeding and death.

# Clinical cases

Work through each step before reading the model answer.

## Case 1: Atonic PPH after a long labour at a HC IV

You are the intern on call at a HC IV in Mukono district at 02:00. A 33-year-old para 5 delivered a 4.0 kg baby 15 minutes ago after a 14-hour labour augmented with oxytocin. The midwife gave oxytocin 10 IU IM at birth and the placenta delivered by CCT. She now calls you: 'the drape is at 600 mL and still filling'.

### 1. What are your first actions in the first minute?

**Call for help** (second midwife, anaesthetic officer, theatre team), note the time, and assign roles. Assess **ABC**: lie her flat, give **oxygen**, and insert **two large-bore cannulae (14–16G)**. Take blood for Hb, **group and cross-match** and a **bedside clotting test**. Start the **E-MOTIVE bundle** immediately; PPH is already diagnosed ( $\geq 500$  mL).

### 2. What quick history and examination points will you check while resuscitating?

History from the notes: parity, labour length, **oxytocin augmentation**, big baby (all **atony** risk factors); any **hypertension** (rules out ergometrine) or **asthma** (rules out carboprost); antenatal Hb, HIV status. Exam: pulse **120**, BP **100/64** (shock index 1.2), cool peripheries. The **fundus is soft and at 2 cm above the umbilicus** = atony. Check the **placenta**: complete. Bladder distended. Brief genital inspection: small first-degree tear, not bleeding much.

### 3. Outline the bundle you will give, with doses.

- **Massage** the uterus and expel clots
- **Oxytocin 10 IU IV slowly**, then **20 IU in 1 L RL at 60 drops/min**
- **TXA 1 g IV over 10 min** (within 3 h of birth)
- **IV fluids**: warmed RL 1 L fast, then reassess
- **Catheterise** (empty bladder; hourly urine)
- **Examine** the genital tract and placenta (done)
- Plan **escalation** if bleeding continues

### 4. Fifteen minutes later the uterus keeps relaxing and loss is now 1100 mL; pulse 128. What next?

**Escalate**: **misoprostol 800 µg SL** (or ergometrine 0.2 mg IM if normotensive), a **second TXA 1 g** if bleeding continues 30 min after the first, and **bimanual compression**. Re-explore the uterus for **retained clots or tissue**. Insert a **condom catheter balloon tamponade** (300–500 mL saline) if still bleeding. **Transfuse** as soon as blood is available (it is a HC IV, so request from the blood bank). Prepare theatre and get consent for laparotomy (B-Lynch, uterine artery ligation, hysterectomy). Consider an early call to the regional hospital if you lack blood or surgical capacity.

### 5. The tamponade stops the bleeding. How will you manage her over the next 24 hours?

Continue the **oxytocin infusion** (20 IU in 1 L at 40 drops/min); give **prophylactic antibiotics** (e.g. ampicillin 2 g IV); vital signs every 15 min for 2 h, then half-hourly; **fluid balance** with urine output  $>30$  mL/h; watch the drain of the balloon. Deflate and remove the balloon after 12–24 h in theatre-ready conditions. Check **Hb at 24 h**; transfuse if symptomatic or Hb  $<7$  g/dL; oral iron and folic acid. Thromboprophylaxis once stable. Breastfeeding support (watch for Sheehan's). **Debrief** her and her husband, and document fully.

## 6. How would you present her on the morning ward round?

'Mrs K, a 33-year-old para 5, is day 0 after SVD of a 4 kg baby following a 14-h augmented labour, complicated by **severe atonic PPH of 1100 mL**. She required the E-MOTIVE bundle, a second-line uterotonic, bimanual compression and condom tamponade, and received 2 units of blood. She is now stable: pulse 92, BP 112/70, urine 50 mL/h, uterus firm, minimal loss. Hb 7.9. Plan: remove the tamponade today, continue the oxytocin infusion until then, repeat Hb, oral iron, and counsel her on family planning and on delivering in hospital next time.'

**Take-away:** Measure, don't guess; give the bundle together and early; escalate in parallel (drugs, compression, tamponade, blood, theatre) instead of waiting to see whether each step works.

## Case 2: A firm uterus but still bleeding

At Mengo Hospital, a 22-year-old primigravida has just had a vacuum delivery for prolonged second stage. The placenta delivered complete 10 minutes later after oxytocin 10 IU IM. The drape shows 700 mL and a steady trickle of bright red blood continues. Pulse 108, BP 112/72.

### 1. What is the single most useful examination finding to decide the cause?

**Palpate the fundus.** If it is **firm, central and at or below the umbilicus**, atony is unlikely. Bright red continuous bleeding with a firm uterus after an instrumental birth suggests **trauma** (cervical, vaginal or perineal tear).

### 2. What differentials would you consider and how will you distinguish them?

1. **Cervical or high vaginal laceration:** speculum, good light, sponge forceps around the whole cervix.
2. **Perineal or episiotomy extension:** visible.
3. **Vulvovaginal haematoma:** severe pain, swelling, rectal pressure.
4. **Retained fragment:** placenta checked complete, less likely.
5. **Coagulopathy:** clotting test; check that blood forms clots.
6. **Uterine rupture** (less likely in a primigravida without a scar): abdominal pain, shock out of proportion.

### 3. Examination under good light shows a 3 cm left lateral cervical tear actively bleeding. How do you manage?

Still start the bundle basics: IV access, bloods, **TXA 1 g IV** (it helps trauma bleeding too), fluids, catheter. Then **repair**: good light, assistant, analgesia or anaesthesia (move to theatre for EUA if the apex can't be seen). Grasp both sides with sponge forceps, place the **first suture above the apex**, and run continuous absorbable sutures (e.g. 2-0 polyglactin). Check haemostasis and the rest of the genital tract.

### 4. After repair, what monitoring and follow-up does she need?

Observations every 15 min for 2 h, then hourly; fundus and loss checks; urine output. **Hb** check; iron and folate. Analgesia. Document the tear and repair. Explain to the woman what happened. Consider a **PPH case review** (why was the tear not seen earlier?).

**Take-away:** A contracted uterus that still bleeds is not an oxytocin problem: look for trauma in good light, repair from above the apex, and remember that TXA also helps trauma-related PPH.

### Case 3: Secondary PPH on day 9

A 28-year-old para 2 presents to a regional referral hospital 9 days after a home birth assisted by a traditional birth attendant. She reports heavy bleeding since morning with clots, fever and lower abdominal pain. Temperature 38.8 °C, pulse 116, BP 100/64. The lochia is foul-smelling.

#### 1. What focused history will you take and why?

- **Place and conduct of birth:** TBA, prolonged labour, was the placenta complete? (retained products)
- **Onset of fever, rigors, pain:** endometritis
- **Urinary symptoms, breast pain, cough, calf pain:** other sources of fever
- **Malaria exposure:** common co-existing cause of fever in Uganda
- **HIV status/ART:** risk of severe sepsis
- Previous bleeding, contraception, breastfeeding
- **Danger symptoms:** dizziness, breathlessness, confusion

#### 2. What examination findings and investigations do you want?

Exam: pallor, signs of sepsis (RR, BP, mental state), a **tender, bulky, soft uterus** above the expected size for day 9 (subinvolution), open os with tissue, foul lochia, perineal wound state, breasts, calves. Investigations: **FBC, blood cultures, HVS/endocervical swab, malaria RDT/smear, group and cross-match**, urea/creatinine, **pelvic ultrasound** for retained products, urinalysis.  $\beta$ -hCG if bleeding persists or tissue looks molar.

#### 3. Outline management step by step.

1. Resuscitate: IV access, fluids, oxygen; transfuse if Hb low or unstable.
2. **IV antibiotics** immediately: **ampicillin 2 g 6-hourly + gentamicin 5 mg/kg daily + metronidazole 500 mg 8-hourly** (or ceftriaxone + metronidazole per local protocol).
3. **Uterotonic:** oxytocin infusion, or misoprostol 800 µg SL.
4. Treat malaria if positive.
5. Once antibiotics have started (usually within hours if bleeding is heavy): **gentle evacuation** of retained products (MVA or blunt curette) by a senior. Beware of perforation.
6. Continue antibiotics until afebrile for 48 h, then oral to complete 7 days.

#### 4. Despite treatment she remains febrile on day 3 of antibiotics. What would you think of?

Consider **pelvic abscess** (ultrasound), **septic pelvic thrombophlebitis** (fever spikes despite antibiotics → heparin), infected haematoma, **wound infection, resistant organism** (review cultures, change to e.g. meropenem where available), **malaria**, UTI/pyelonephritis, mastitis, drug fever, and rarely retained products still present or GTN.

**Take-away:** Secondary PPH is usually infection plus retained products: give antibiotics first, then evacuate gently, and look for other causes of fever (including malaria) if she doesn't settle.

## References

1. Uganda Clinical Guidelines 2023: National Guidelines for Management of Common Health Conditions. *Ministry of Health Uganda*, 2023. <https://library.health.go.ug/uganda-clinical-guidelines-2023>
2. Essential Maternal and Newborn Clinical Care Guidelines for Uganda. *Ministry of Health Uganda*, 2022. <https://library.health.go.ug/sites/default/files/resources/Essential%20Maternal%20Newborn%20Care%20Guidelines%202022%20V3.pdf>

3. WHO recommendations on the assessment of postpartum blood loss and use of a treatment bundle for postpartum haemorrhage. *World Health Organization*, 2023. <https://www.who.int/publications/i/item/9789240085398>
4. Consolidated guidelines for the prevention, diagnosis and treatment of postpartum haemorrhage: Executive summary. *WHO / FIGO / ICM (NCBI Bookshelf)*, 2025. <https://www.ncbi.nlm.nih.gov/books/NBK619236/>
5. Global health agencies issue new recommendations to help end deaths from postpartum haemorrhage. *World Health Organization news release*, 2025. <https://www.who.int/news/item/05-10-2025-global-health-agencies-issue-new-recommendations-to-help-end-deaths-from-postpartum-haemorrhage>
6. Randomized Trial of Early Detection and Treatment of Postpartum Hemorrhage (E-MOTIVE). *Gallos I, et al. New England Journal of Medicine*, 2023. <https://www.nejm.org/doi/full/10.1056/NEJMoa2303966>
7. Effect of early tranexamic acid administration on mortality, hysterectomy, and other morbidities in women with post-partum haemorrhage (WOMAN). *WOMAN Trial Collaborators. The Lancet*, 2017. [https://www.thelancet.com/journals/lancet/article/PIIS0140-6736\(17\)30638-4/fulltext](https://www.thelancet.com/journals/lancet/article/PIIS0140-6736(17)30638-4/fulltext)
8. Heat-Stable Carbetocin versus Oxytocin to Prevent Hemorrhage after Vaginal Birth (CHAMPION). *Widmer M, et al. New England Journal of Medicine*, 2018. <https://www.nejm.org/doi/full/10.1056/nejmoa1805489>
9. WHO recommendations: Uterotonics for the prevention of postpartum haemorrhage. *World Health Organization*, 2018. <https://apps.who.int/iris/bitstream/handle/10665/277276/9789241550420-eng.pdf>
10. WHO recommendation on uterine balloon tamponade for the treatment of postpartum haemorrhage. *World Health Organization (NCBI Bookshelf)*, 2021. <https://www.ncbi.nlm.nih.gov/books/NBK569828/>
11. Prevention and Management of Postpartum Haemorrhage (Green-top Guideline No. 52). *Royal College of Obstetricians and Gynaecologists*, 2016. <https://www.rcog.org.uk/guidance/browse-all-guidance/green-top-guidelines/prevention-and-management-of-postpartum-haemorrhage-green-top-guideline-no-52/>
12. FIGO recommendations on the management of postpartum hemorrhage 2022. *Escobar MF, et al. International Journal of Gynecology & Obstetrics*, 2022. <https://pmc.ncbi.nlm.nih.gov/articles/PMC9313855/>
13. Non-pneumatic anti-shock garment to stabilize women with hypovolemic shock secondary to obstetric hemorrhage. *FIGO Safe Motherhood and Newborn Health Committee. Int J Gynecol Obstet*, 2015. <https://obgyn.onlinelibrary.wiley.com/doi/full/10.1016/j.ijgo.2014.10.014>
14. Preventing and Treating Postpartum Haemorrhage: Uterotonics and Tranexamic Acid (Anaesthesia Tutorial of the Week). *World Federation of Societies of Anaesthesiologists*. <https://resources.wfsahq.org/atotw/preventing-and-treating-postpartum-haemorrhage-uterotonics-and-tranexamic-acid/>
15. Incidence and risk factors for postpartum hemorrhage in Uganda. *Ononge S, et al. Reproductive Health*, 2016. <https://pmc.ncbi.nlm.nih.gov/articles/PMC4832492/>
16. Severe postpartum haemorrhage at a large referral hospital in Uganda: A prospective observational pilot study. *PLOS One*, 2025. <https://journals.plos.org/plosone/article?id=10.1371%2Fjournal.pone.0331512>

---

A study aid, not a protocol. Check doses against the Uganda Clinical Guidelines 2023, your unit's protocol and a senior before you treat a patient.